

Universal conductivity of graphene in the ultra-relativistic regime

Igor F. Herbut^{1,2} and Vieri Mastropietro³

¹ *Department of Physics, Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6*

² *Max-Planck-Institut für Physik Komplexer Systeme,
Nöthnitzer Strasse 38, 01187 Dresden, Germany*

³ *Dipartimento di Matematica “Federigo Enriques”,
Università degli Studi di Milano, Via Cesare Saldini 50, 20133 Milano, Italy*

We calculate the zero temperature dc conductivity in graphene in the ultimate low-energy regime, when retardation effects of the electromagnetic interaction become important and when the full Lorentz symmetry emerges. In contrast to what happens with the short range or with the Coulomb long-range instantaneous interactions, the dc conductivity is now no longer equal to its non interacting value, but acquires universal corrections in powers of the fine structure constant. The coefficient of the first order correction is computed, and found to be of order one. We also present the result for the conductivity in the large- N limit, with N as the number of Dirac fermions species, to the order $1/N^2$.

INTRODUCTION

Graphene owes several of its remarkable properties to the fact that it admits an effective relativistic quantum field theory description in terms of massless Dirac fermions in two dimension. A dramatic manifestation of this fact is in the conductivity properties at half filling; recent experiments [1] found that the optical conductivity in monolayer graphene is essentially constant in a wide range of frequencies and very close to the value $(\pi/2)(e^2/h)$ (that is $1/4$ in the natural units $e = \hbar = c = 1$ which we will use), which is precisely the value found for the system of non-interacting Dirac fermions [2]. This result, however, raises a couple of natural questions: why the interactions, which are rather strong in graphene, do not produce visible corrections to the non interacting value of the conductivity? And should the dc conductivity in graphene be truly equal to its non interacting value, or there are in principle corrections which may lie inside the experimental errors?

The computation of the many body interaction effects on graphene’s conductivity is quite sensitive to regularizations and approximations, and several aspects of it have been controversial. In the case of *short range* interactions, after first perturbative computations claiming non vanishing corrections, it was finally rigorously proved [3] that there are *no interaction corrections* to the dc conductivity; all the interaction contributions to the conductivity cancel out at all orders in the renormalized expansion. The exact vanishing of interaction correction emerges as a consequence of the Ward identities and the irrelevance, in the technical renormalization group (RG) sense, of the interaction,[3, 4]. On the other hand, in the case of *long range* Coulomb interactions it has been predicted that the dc conductivity is still universal and equal to $1/4$ [5], the argument this time being based on the divergence of the Fermi velocity [6], and the *relative*

(to the kinetic energy) irrelevance of the interaction. The low frequency correction to the conductivity was found to be not particularly small, due to the slow logarithmic increase of the Fermi velocity, which is what is expected in absence of (possible) accidental cancellation. A controversy in the computation of such corrections arose in literature, [4, 5, 7–9]. Technically the reason for the controversy lies in the ambiguities produced by the ultraviolet divergences due to the continuum limit. Recently, however, the controversy has been claimed to be settled by performing a lattice computation in [10], in favor of the value originally found in [4].

On the other hand, the Fermi velocity divergence found in the Coulomb case at very low frequencies is clearly rather unphysical, and simply signals ultimate inadequacy of the usual model of instantaneous Coulomb interaction. With the increase of the Fermi velocity the retardation effects eventually become important, so that the retarded current-current interaction must be added to the Coulomb density-density interaction. Such effects have been analyzed before in [11], [12] by a RG analysis, and it was found that the flow of the Fermi velocity stops at the velocity of light c , and, maybe most importantly, that the coupling constant (i. e. the charge) in the theory is *exactly* marginal. In particular, a lattice model for graphene interacting with an electromagnetic field was considered in [12], and by iterating the RG it was proved that Lorentz symmetry spontaneously emerges, and that the system is asymptotically close to the so-called “reduced” quantum electrodynamics ($QED_{4,3}$).

The aim of this paper is to compute the corrections to the non-interacting value of the universal dc conductivity due to the full electromagnetic interaction. We show first by exact RG methods that, with the retardation effects and with the honeycomb lattice included, in units of e^2/h ,

$$\sigma = \frac{N}{8} \left(1 + \left(C_1 - \frac{N}{8} \right) \frac{e^2}{2} + O(e^4) \right), \quad (1)$$

where N is the number of four-component Dirac fermions ($N = 2$ in graphene), and C_1 is an N -independent constant. In the ultimate, low-energy regime the conductivity is therefore *different* with respect to its non interacting value of $1/4$; the dependence on the number of fermion components N in the first interaction correction immediately rules out possible cancellations in general. On the other hand, the correction to conductivity, in vacuum with the dielectric constant $\epsilon = 1$ assumed above, is still *universal*, as a consequence of the emerging Lorentz invariance, in the sense that it does not depend on the material parameters such as the Fermi velocity, but only on the fine structure constant.

As the lattice model for graphene becomes asymptotically close to $QED_{4,3}$, which is the fixed point of the RG flow, it is worthwhile computing the dc conductivity directly in this continuum model by using standard field-theoretical methods. This way an expression identical to (1) is found, with the numerical constant $C_1 = 0.0089319$. The leading $O(e^2N)$ correction is *identical* both in the lattice and in the continuum model. In graphene, the number of Dirac fermions is $N = 2$, and this correction is 0.125 and much larger than the constant C_1 . In any case, as $e^2 = 4\pi\alpha$, with $\alpha = 1/137.036$ as the fine structure constant, in vacuum we find $\sigma/\sigma_0 = 1 - 0.01064$, a small correction, and at the moment within the experimental error in [1]. The conductivity is therefore *in principle*, if not in practice, different from the non interacting value and, because of the smallness of the fine structure constant, with a new universal value only slightly reduced from the non-interacting one. Finally, we observe in passing that the prefactor multiplying the charge e^2 in the correction term is 0.11607 , of the same order in magnitude as found in the non-relativistic limit for the static Coulomb interaction [4, 5, 10], although of the opposite sign.

RENORMALIZATION GROUP COMPUTATION

We introduce the lattice $\Lambda = (n_1\vec{l}_1 + n_2\vec{l}_2)$, $n_i = 0, \dots, L-1$ and $\vec{l}_{1,2} = \frac{1}{2}(3, \pm\sqrt{3})$; the Fourier transform is $f(\vec{x}) = \frac{1}{|\Lambda|} \sum_{\vec{k} \in D} e^{i\vec{k}\vec{x}} \hat{f}_{\vec{k}}$, where $\hat{f}_{\vec{x}} = \sum_{\vec{x} \in \Lambda} e^{-i\vec{k}\vec{x}} f_{\vec{x}}$, $D = \{\vec{k} = \frac{m_1}{L}\vec{b}_1 + \frac{m_2}{L}\vec{b}_2\}$, $|\Lambda| = 3\sqrt{3}L^2/2$ and $\sum_{\vec{x}} e^{i\vec{k}\vec{x}} = |\Lambda|\delta_{\vec{k},0}$. In the limit $L \rightarrow \infty$ $\frac{1}{|\Lambda|} \sum_{\vec{k} \in D_L} \rightarrow \int_{\mathcal{B}} \frac{d\vec{k}}{(2\pi)^2}$ where the integral is over the Brillouin zone $\{\vec{k} = \alpha_1\vec{b}_1 + \alpha_2\vec{b}_2\}$, with $\alpha_{1,2} \in [0, 1)$. The Hamiltonian describing electrons on the honeycomb lattice interacting with an electromagnetic field is $H = H_e + H_a$, where

$$H_e = -t \sum_{\substack{\vec{x} \in \Lambda \\ j=1,2,3}} \sum_{\sigma=1}^N a_{\vec{x},\sigma}^+ b_{\vec{x}+\vec{\delta}_j,\sigma}^- e^{ie \int_0^1 \vec{\delta}_j \vec{a}(\vec{x}+s\vec{\delta}_j,0)} + c.c., \quad (2)$$

with $\vec{\delta}_1 = (1, 0)$, $\vec{\delta}_2 = \frac{1}{2}(-1, \sqrt{3})$, $\vec{\delta}_3 = \frac{1}{2}(-1, -\sqrt{3})$ and H_A is the free photon Hamiltonian. Note that the photons live in the three dimensional space but they interact with fermions bound to a two dimensional plane.

The physical observables are conveniently obtained in terms of the following generating functional:

$$e^{W_{L,\beta}(A,\lambda)} = \int P(d\psi) \int P(da) e^{\mathcal{V}(a+A,\psi)+(\psi,\lambda)}, \quad (3)$$

where $\psi_{\mathbf{x}}^{\pm} = (a_{\mathbf{x},\sigma}^{\pm}, b_{\mathbf{x}+\vec{\delta}_1,\sigma}^{\pm})$ are Grassman variables (denoted with a slight abuse of notation with the same symbol), $\sigma = 1, \dots, N$, $\delta_j = (0, \vec{\delta}_j)$, $P(d\psi)$ is the fermionic integration with propagator

$$g(\mathbf{x}-\mathbf{y}) = \frac{1}{\beta|\Lambda|} \sum_{\mathbf{k} \in \mathcal{D}} e^{i\mathbf{k}(\mathbf{x}-\mathbf{y})} \begin{pmatrix} ik_0 & v\Omega^*(\vec{k}) \\ v\Omega(\vec{k}) & ik_0 \end{pmatrix}^{-1} \quad (4)$$

with $\mathbf{x} = (x_0, \vec{x})$, $\mathbf{k} = (k_0, \vec{k})$, $k_0 = \frac{2\pi}{\beta}(m+1/2)$, $v = \frac{3}{2}t$, and $\Omega(\vec{k}) = \frac{2}{3} \sum_{j=1,2,3} e^{i\vec{k}(\vec{\delta}_j - \vec{\delta}_1)}$. The dispersion relation $\Omega(\vec{k})$ vanishes at the two Fermi points $\mathbf{k}_F^{\pm} = (0, \frac{2\pi}{3}, \pm \frac{2\pi}{3\sqrt{3}})$. $a_{\mu,\mathbf{x}}$ is a Gaussian variable while $P(da)$ is the photon integration with propagator $\delta_{\mu,\nu}w(\mathbf{x}-\mathbf{y})$ with

$$w(\mathbf{x}-\mathbf{y}) = \frac{1}{\beta|\Lambda|} \sum_{\mathbf{k} \in \mathcal{D}} e^{i\mathbf{k}(\mathbf{x}-\mathbf{y})} \frac{1}{2|\mathbf{k}|}, \quad (5)$$

where the Feynman gauge is assumed. Finally the interaction \mathcal{V} can be easily deduced from (2) and it has the form

$$V(a, \psi) = \sum_{\mu=1,2,3} \int d\mathbf{x} a_{\mu,\mathbf{x}} j_{\mu,\mathbf{x}} + F(a), \quad (6)$$

where $F(a)$ contains higher order term in A , $j_{\mu,\mathbf{x}} = (j_{0,\mathbf{x}}, \vec{j}_{\mathbf{x}})_{\mu}$ with

$$j_{0,\mathbf{x}} = \sum_{\sigma=1}^N [a_{\mathbf{x},\sigma}^+ a_{\mathbf{x},\sigma}^- + a_{\mathbf{x}+\vec{\delta}_1,\sigma}^+ a_{\mathbf{x}+\vec{\delta}_1,\sigma}^-] \\ \vec{j}_{\mathbf{x}} = \frac{2}{3} \sum_{j=1,2,3} \sum_{\sigma=1}^N \vec{\delta}_j (a_{\mathbf{x},\sigma}^+ b_{\mathbf{x}+\vec{\delta}_j,\sigma}^- - b_{\mathbf{x}+\vec{\delta}_j,\sigma}^+ a_{\mathbf{x},\sigma}^-). \quad (7)$$

We will denote $\lim_{\beta \rightarrow \infty} \lim_{L \rightarrow \infty} W_{L,\beta}(A, \lambda) = W_{L,\beta}(A, \lambda)$. From the identity $W_{L,\beta}(A, \lambda) = W_{L,\beta}(A + \partial\alpha, \lambda e^{i\alpha\mathbf{x}})$ we get

$$0 = \frac{\partial}{\partial\alpha} W_{L,\beta}(A + \partial\alpha, \lambda e^{i\alpha\mathbf{x}}) = 0, \quad (8)$$

and the derivatives of the above relations provide a set of Ward identities. Calling $K_{\mu,\nu}(\mathbf{p}) = \frac{\partial^2 W_{L,\beta}(A,0)}{\partial \hat{J}_{\mu,\mathbf{p}} \partial \hat{J}_{\nu,-\mathbf{p}}} |0$, $\mathbf{p} = (\omega, p)$, the dc conductivity at zero frequency is defined via Kubo formula $\sigma = \lim_{\omega \rightarrow 0} -e^2 \frac{1}{\omega} K_{i,i}(\omega, 0)$. Note that $K_{i,j}(\mathbf{p})$ is the sum

of the truncated current-current correlation and of the diamagnetic term, which is a constant in \mathbf{p} ; by the Ward identity obtained from (8) with a derivative with respect to A , we get $\lim_{p_1 \rightarrow 0} \lim_{\omega \rightarrow 0} \widehat{K}_{11}(\mathbf{p})|_{p_2=0} = \lim_{p_1 \rightarrow 0} \lim_{\omega \rightarrow 0} \frac{\omega}{p_1} \widehat{K}_{01}(\mathbf{p}) = 0$ and, as $\widehat{K}_{11}(\mathbf{p})$ is continuous at weak coupling, we can reverse the limits so that the conductivity can be written as

$$\sigma = \lim_{\omega \rightarrow 0^+} -e^2 \frac{\widehat{K}_{ii}(\omega, 0) - \widehat{K}_{ii}(0, 0)}{\omega}. \quad (9)$$

There is therefore no need of computing the diamagnetic term, but it is sufficient to compute the current-current correlation and subtract the value in $\mathbf{p} = (0, 0)$.

$K_{i,j}(\mathbf{p})$ can be computed by Wilsonian RG; for details,

see [12]. The starting point consists of writing the Grassman variables as sums of variables with momenta closer and closer to the two Fermi points $\mathbf{k}_F^\pm = (0, \frac{2\pi}{3}, \pm \frac{2\pi}{3\sqrt{3}})$, that is

$$\widehat{\psi} = \psi^{(1)} + \sum_{\varepsilon=\pm} \sum_{h=-\infty}^0 \psi_\varepsilon^{(h)} \quad (10)$$

where $\psi_\varepsilon^{(h)}$ lives on a shell of momenta distant $O(2^h)$ from \mathbf{k}_F^ε . Similarly, we can write $a^\mu = \sum_{h=-\infty}^0 a_h^\mu$, and a_h^μ has propagator $f_h(\mathbf{k})w(\mathbf{k})$ with $f_h(\mathbf{k})$ non vanishing for $2^{h-1} \leq |\mathbf{k}| \leq 2^{2+h}$. After the integration of the fields $(\psi^1, a^1), \dots, (\psi^h, a^h)$, one finds that the generating functional can be written as

$$e^{W_{L,\beta}(A,\lambda)} = e^{B^h(A,\lambda)} \int \prod_{\varepsilon=\pm 1} P(d\psi_\varepsilon^{(\leq h)}) \int P(da^{(\leq h)}) e^{V^h(\sqrt{Z_h}\psi^{(\leq h)}, a^{(\leq h)} + A, \lambda)}, \quad (11)$$

where $P(d\psi_\varepsilon^{(h)})$ has the propagator

$$g_\varepsilon^{(h)}(\mathbf{k}' + \mathbf{k}_F^\varepsilon) = \frac{1}{Z_h} \begin{pmatrix} ik_0 & v_h \Omega^*(\vec{k}) \\ v_h \Omega(\vec{k}) & ik_0 \end{pmatrix}^{-1}. \quad (12)$$

Z_h is the wave function renormalization and v_h is the effective Fermi velocity. Note also that (13) can be written as

$$g_\varepsilon^{(h)} \sim \frac{1}{Z_h} \begin{pmatrix} ik_0 & v_h(-ik'_1 + \varepsilon k_2) \\ v_h(ik'_1 + \varepsilon k_2) & ik_0 \end{pmatrix}^{-1} \quad (13)$$

so that it becomes ever closer to the Dirac propagator. By power counting, the scaling dimension of the interactions is $D = 3 - n$ if n is the number of fields; the local terms quadratic in a , describing the photon mass, are absent by exploiting the Ward identity generated by (8) (see App. E1 of [12] for an explicit computation at one loop) while the marginal terms $a^+ \partial a$ are vanishing by symmetry; in the same way the local terms $\psi^+ \psi$ are vanishing by parity, and $\psi^+ \partial \psi$ contribute to the wave function renormalization and the Fermi velocity. The effective potential is given by (at $\lambda = 0$ for definiteness):

$$V^h(\sqrt{Z_h}\psi^{\leq h}, a+A, 0) = \sum_{\mu} \sum_{\mathbf{x}} Z_{\mu,h}(a_{\mu,\mathbf{x}}^{\leq h} + A_{\mu,\mathbf{x}}) j_{\mu,\mathbf{x}}^{\leq h} + F_h, \quad (14)$$

where F_h are the *irrelevant terms*, with negative dimension $D < 0$.

From the Ward identities (8) computed at momenta at

the scale of the infrared cut-off we get

$$\frac{Z_{0,h}}{Z_h} = 1 + O(e^2), \quad \frac{Z_{i,h}}{Z_h v_h} = 1 + O(e^2). \quad (15)$$

Moreover, by symmetry $Z_h^1 = Z_h^2$ and by an explicit computation,

$$Z_h \sim 2^{-\eta h}, \quad v_h \sim 1 + A(1-v)2^{\tilde{\eta}h}, \quad (16)$$

with $\eta = \frac{e^2}{12\pi^2} + O(e^4)$, $\tilde{\eta} = \frac{2e^2}{5\pi^2} + O(e^4)$, and A as a bounded function. Therefore the wave function renormalization is diverging (i. e. there is an anomalous dimension), and by iterating the RG the Fermi velocity converges to the velocity of light.

The model considered in [11] therefore emerges naturally starting from the lattice model. Moreover, the Lorentz symmetry is restored in the infrared limit, as the Fermi velocity flows up to the velocity of light (which in our units is 1). The effective couplings $e_{h,0} = e \frac{Z_h^0}{Z_h}$ $e_{h,i} = e \frac{Z_h^1}{Z_h v_h}$ flow to a *line* of fixed points

$$e_{h,i} \rightarrow e + O(e^3), \quad (17)$$

that is, the theory is *exactly marginal*. One obtains then a renormalized expansion for the current-current correlations in terms of the effective couplings $e_{\mu,h}$, namely $\sigma = \sigma^{(0)} + \sigma^{(2)} + \dots$. This differs from perturbative expansion in that there are no infrared divergences and all coefficients are bounded. We get

$$\sigma^{(0)} = Ne^2 \lim_{\omega \rightarrow 0^+} \sum_{\varepsilon=\pm} \frac{1}{\omega} \sum_{h \leq 1} \int \frac{dk_0}{2\pi} \int_{\mathcal{B}} \frac{d\vec{k}'}{(2\pi)^2} \frac{(Z_h^{(i)})^2}{Z_h Z_h} \text{Tr} \left\{ \sigma_j g_\varepsilon^{(h)}(\mathbf{k}') \sigma_i [g_\varepsilon^{(h)}(\mathbf{k}' + (\omega, \vec{0})) - g_\varepsilon^{(h)}(\mathbf{k}')] \right\}, \quad (18)$$

where σ_j are Pauli matrices and $N = 2$ in graphene. Note the presence of a factor $\frac{1}{Z_h}$ for any fermionic line, and of a factor $Z_{i,h}$ for any vertex. The presence of such factors could radically alter the conductivity properties; for instance if we do not take into account the vertex renormalization, that is we replace $Z_{i,h}$ with unity, one would get $\sigma^{(0)} = 0$. On the contrary, thanks to the Ward identity and (15) we can replace $\frac{(Z_h^{(i)})^2}{Z_h Z_h}$ with v_h up to $O(e^2)$ terms. The integral then still appears to be non universal, as it depends of the effective Fermi velocity and on the lattice details. However it is not so; we can write integral over the Brillouin zone \mathcal{B} as sum of two terms, one $|\Omega(\vec{k})| \geq \varepsilon$ and the other $|\Omega(\vec{k})| < \varepsilon$; the former is uniformly convergent as $\omega \rightarrow 0^+$: therefore, we can exchange the integral with the limit and check that the

integral of the limit is zero simply because the integrand is odd in k_0 . In the latter we use (13) neglecting the corrections (as the size of the integral is arbitrarily small); the dependence from v_h disappears through a change of variables and we finally get $\sigma^{(0)} = e^2 \frac{N}{8} + O(Ne^4)$, that is the same result as in the non interacting case. Moreover, as shown in [13], the low frequency corrections are $O(e^2 \omega^2)$.

Let us consider now $\sigma^{(2)}$; there are three possible contributions, $\sigma^{(2)} = \sigma_a^{(2)} + \sigma_b^{(2)} + \sigma_c^{(2)}$ but only one of them, which we call $\sigma_c^{(2)}$, is proportional to N^2 . Therefore, if we find it to be non vanishing we can safely conclude that the dc conductivity is different from the non-interacting value, at least for a generic N . The value of such term is

$$\sigma_c^{(2)} = -N^2 e^2 \lim_{\omega \rightarrow 0^+} \frac{1}{2} \left\{ \frac{1}{\omega} \sum_{\varepsilon=\pm} \sum_{h(\omega) \leq h \leq 1} \int \frac{dk_0}{2\pi} \int_{\mathcal{B}} \frac{d\vec{k}'}{(2\pi)^2} \frac{Z_{i,h}}{Z_h} e_{i,h} v_h \text{Tr} \left\{ \sigma_1(\vec{k}') g_\varepsilon^{(h)}(\mathbf{k}') \sigma_i [g_\varepsilon^{(h)}(\mathbf{k}' + (\omega, \vec{0})) - g_\varepsilon^{(h)}(\mathbf{k}')] \right\} \right\} \quad (19)$$

where $h(\omega)$ is such that $2^{h(\omega)} \sim |\omega|$. In writing the above expression we have used that only the renormalized parts of the “bubble” diagrams contribute, as their local part is vanishing. Again using that $e_{i,h} = e + O(e^3)$ $Z_{i,h}/Z_h = v_h + O(e^2)$, and using that each of the two terms in (19) can be evaluated as (18) we get $\sigma_c^{(2)} = -(e^4/2)(N/8)^2 + O(e^4 N)$, and finally (1) is found. The N -dependence allows us to exclude the possibility of cancellations, and we can conclude that the dc conductivity is different from its non interacting value, with a leading correction which is universal. Note the difference with the case of Coulomb interactions: in such a case at one loop $v_h \sim 1/|h|$, while $Z_h, Z_{i,h}, e_{i,h}$ are essentially constants so that $\sigma_c^{(2)}$ vanishes. Similarly, for Hubbard interactions the photon propagator $1/2|\omega|$ should be replaced by a constant, and again $\sigma_c^{(2)}$ would be vanishing.

EFFECTIVE QED DESCRIPTION

The previous analysis shows that lattice graphene system flows, by iterating the RG, to a fixed point expressed by the the continuum $QED_{4,3}$. It is useful then to study the conductivity properties in the continuum theory in

order to get, assuming, as usual, that the value of the conductivity depends only on the fixed point of the RG,[14] more information on the subleading corrections. The action for the effective model in 2 + 1 dimensions is

$$S = \int d\mathbf{x} [\bar{\Psi}_i(\mathbf{x}) \gamma_\mu (\partial_\mu - e(a_\mu(\mathbf{x}) - A_\mu(\mathbf{x}))) \Psi_i(\mathbf{x}) + \frac{1}{2} \int d\mathbf{x} d\mathbf{y} W_{\mu\nu}^{-1}(\mathbf{x} - \mathbf{y}) a_\mu(\mathbf{x}) a_\nu(\mathbf{y})], \quad (20)$$

where Ψ is a four-component fermionic field, $i = 1, \dots, N$, $\mathbf{x} = (x_0, x_1, x_2)$, and the summation convention is assumed. Note that $\vec{x} = (x_1, x_2)$ is here a continuum variable while in the previous section was a site on the honeycomb lattice. The bare gauge field propagator is

$$W_{\mu\nu}(\mathbf{x}) = \frac{1}{2} \int \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{e^{i\mathbf{q}\mathbf{x}}}{|\mathbf{q}|} (\Pi_{\mu\nu}(\mathbf{q}) + \beta \frac{\mathbf{q}_\nu \mathbf{q}_\mu}{\mathbf{q}^2}), \quad (21)$$

with the usual transverse projector

$$\Pi_{\mu\nu}(\mathbf{q}) = \delta_{\mu\nu} - \frac{\mathbf{q}_\mu \mathbf{q}_\nu}{\mathbf{q}^2}. \quad (22)$$

The above field theory is closely related to the three dimensional quantum electrodynamics (QED_3),[15] with

one important caveat: the ‘‘Maxwell’’ term is now already at the bare level non-analytic in momentum, and proportional to $|\mathbf{q}|$, and not to the usual \mathbf{q}^2 . It can be obtained from the reduced quantum electrodynamics $QED_{4,3}$ in which the electromagnetic fields live in $3 + 1$ dimensions, but are coupled to fermions which are confined to the lower, $2 + 1$ -dimensional ‘‘brane’’, by ‘‘integrating out’’ the out-of-plane components of the vector potential.[16] This procedure in general also changes the effective gauge-fixing parameter from β' in the original $3 + 1$ -dimensional theory into β in the above expression, as in $\beta = (1 + \beta')/2$. We see that only the Feynman gauge

($\beta = 1$) remains invariant under this dimensional reduction, which makes it the most convenient one from the practical point of view. The non-analyticity of the gauge field propagator around $\mathbf{q} = 0$ can be understood as the reason for the exact marginality of the charge coupling. [17]

Fermions appear quadratically in the action and can be (formally) integrated out. If we redefine the fields as $ea_\mu \rightarrow a_\mu$, and $eA_\mu \rightarrow A_\mu$, and then shift the fluctuating field as $a_\mu - A_\mu \rightarrow a_\mu$, the result of this integration would be the action

$$\tilde{S} = \frac{1}{2} \int \frac{d\mathbf{q}}{(2\pi)^3} \left\{ \left(\frac{N}{8} |\mathbf{q}| \Pi_{\mu\nu}(\mathbf{q}) a_\mu(\mathbf{q}) a_\nu(-\mathbf{q}) + \frac{2}{e^2} |\mathbf{q}| \left(\Pi_{\mu\nu}(\mathbf{q}) + \frac{\mathbf{q}_\nu \mathbf{q}_\mu}{\beta \mathbf{q}^2} \right) (a_\mu(\mathbf{q}) + A_\mu(\mathbf{q})) (a_\nu(-\mathbf{q}) + A_\nu(-\mathbf{q})) \right\} + V(a) \right. \quad (23)$$

The first term proportional to N is the familiar one-loop polarization in the QED_3 , [19] and the second term is the quasi-Maxwell term, which now after the shift of variables also includes the external probe; $V(a)$ is a sum of monomials in the fluctuating gauge field a with degree ≥ 4 . Note that $V(a)$ does not contain the external

probe A , which after the shift appears only in the quasi-Maxwell term. This allows one to differentiate with respect to the external probe, and so to obtain the current-current correlation function in terms of the *exact* gauge field propagator, $D_{\mu\nu}^{aa} = \langle a_\mu a_\nu \rangle$:

$$\langle j_\mu(\mathbf{q}) j_\nu(-\mathbf{q}) \rangle = \frac{2}{e^2} |\mathbf{q}| \left(\Pi_{\mu\nu}(\mathbf{q}) + \frac{1}{\beta} \frac{\mathbf{q}_\nu \mathbf{q}_\mu}{\mathbf{q}^2} \right) - \frac{2}{e^2} |\mathbf{q}| \left(\Pi_{\mu\alpha}(\mathbf{q}) + \frac{1}{\beta} \frac{\mathbf{q}_\mu \mathbf{q}_\alpha}{\mathbf{q}^2} \right) D_{\alpha\beta}^{aa}(\mathbf{q}) \frac{2}{e^2} |\mathbf{q}| \left(\Pi_{\beta\nu}(\mathbf{q}) + \frac{1}{\beta} \frac{\mathbf{q}_\beta \mathbf{q}_\nu}{\mathbf{q}^2} \right). \quad (24)$$

Current conservation, on the other hand, dictates that the exact gauge-field propagator has the form:[20]

$$D_{\mu\nu}^{aa}(\mathbf{q}) = \frac{1}{|\mathbf{q}|} \left(A \Pi_{\mu\nu}(\mathbf{q}) + \beta \frac{e^2}{2} \frac{\mathbf{q}_\nu \mathbf{q}_\mu}{\mathbf{q}^2} \right), \quad (25)$$

where A is a function of the number of fermions N and of the coupling e^2 . Inserting this form into the expression for the current-current correlation function, we find that

$$\langle j_\mu(\mathbf{q}) j_\nu(-\mathbf{q}) \rangle = \sigma |\mathbf{q}| \Pi_{\mu\nu}(\mathbf{q}), \quad (26)$$

where the dc conductivity σ , in units of e^2/\hbar , is simply

$$\sigma = \frac{2}{e^2} \left(1 - \frac{2A}{e^2} \right). \quad (27)$$

Note that the current-current correlation function is completely independent of the gauge-fixing parameter β , just as one expects. We can rewrite (23) as a perturbed Gaussian action

$$\tilde{S} = \frac{1}{2} \int \frac{d\mathbf{q}}{(2\pi)^3} \left\{ \left(\frac{N}{8} + \frac{2}{e^2} \right) |\mathbf{q}| \left(\Pi_{\mu\nu}(\mathbf{q}) + \gamma \frac{\mathbf{q}_\nu \mathbf{q}_\mu}{\mathbf{q}^2} \right) a_\mu(\mathbf{q}) a_\nu(\mathbf{q}) + \frac{2}{e^2} |\mathbf{q}| \left(\Pi_{\mu\nu}(\mathbf{q}) + \frac{1}{\beta} \frac{\mathbf{q}_\nu \mathbf{q}_\mu}{\mathbf{q}^2} \right) \left((2a_\mu(\mathbf{q}) A_\nu(-\mathbf{q}) + A_\mu(\mathbf{q}) A_\nu(-\mathbf{q})) \right) \right\} + V(a). \quad (28)$$

where $\gamma = (2/\beta e^2)(1/(N/8 + 2/e^2))$. In the gaussian ap-

proximation, that is neglecting the higher-order terms

given by $V(a)$, the functional integral can be explicitly performed and the constant A is given by $A_0 = ((N/8) + (2/e^2))^{-1}$. From (27) we this way find the Ioffe-Larkin-like [21] result for the conductivity

$$\sigma^{-1} = \left[\frac{2}{e^2} \left(1 - A_0 \frac{2}{e^2} \right) \right]^{-1} = \frac{8}{N} + \frac{e^2}{2}. \quad (29)$$

which can be interpreted as the addition of the fermion's and the gauge-field's *resistivities* into the total resistivity. Expanding to the first power in the weak charge coupling e^2 yields $\sigma = (N/8)(1 - (Ne^2/16) + \dots)$, in agreement with the lattice computation.

To go beyond the gaussian approximation we can expand in powers of the effective charge in the theory $e^2/(2 + (Ne^2/8))$; one finds

$$\frac{1}{A} = \frac{2}{e^2} + \frac{N}{8} + Nx \frac{e^2}{\frac{N}{8}e^2 + 2} \dots, \quad (30)$$

where $x = (92 - 9\pi^2)/((4\pi)^2 18)$ [18]. In the weak coupling regime $e^2 \ll 1/N$, after expanding in powers of e^2 ,

$$\sigma = \frac{2}{e^2} \left(1 - \frac{1}{1 + \frac{N}{16}e^2 + \frac{Nx e^4}{4} + \dots} \right) = \frac{N}{8} \left(1 - \frac{Ne^2}{16} + e^2 4x + O(e^4) \dots \right), \quad (31)$$

so that Eq. (1) is recovered, with the numerical value of the constant

$$C_1 = 8x = \frac{23}{(3\pi)^2} - \frac{1}{4} = 0.0089319. \quad (32)$$

On the other hand in the large- N limit, $N \gg 1/e^2$, we expand in powers of $1/N$,

$$\sigma = \frac{2}{e^2} \left(1 - \frac{2}{e^2} \frac{1}{\frac{N}{8} + \frac{2}{e^2} + 8x \left(1 - \frac{16}{Ne^2} \right) + \dots} \right), \quad (33)$$

so that

$$\sigma = \frac{2}{e^2} \left(1 - \frac{16}{Ne^2} + \frac{128}{N^2 e^4} (2 + C_1 e^2) + O(N^{-3}) \right). \quad (34)$$

CONCLUSION

In conclusion, we have presented the computation of the universal zero temperature dc conductivity in the ultimate, relativistic, low-energy regime in graphene, when the Fermi velocity has reached the velocity of light. Although this regime lies beyond presently available experimental conditions, such as the temperature and the sample sizes, the issue of ultimate value of the dc conductivity is theoretically interesting, and presents an important question of principle. We find that the ultimate

value of the conductivity is universal, and dependent only on the fine-structure constant of the media surrounding the graphene sheet, but in principle different from the non-interacting value seen in the experiment. The difference from the non interacting value is of the order of the fine structure constant itself, however, and therefore of the relative size of the order of one percent.

ACKNOWLEDGEMENT

IFH was supported by the NSERC of Canada.

-
- [1] R. R. Nair, P. Blake, A. N. Grigorenko, K. S. Novoselov, T. J. Booth, T. Stauber, N. M. R. Peres, A. K. Geim, *Science* **320**, 1308 (2008)
 - [2] A. W. W. Ludwig, M. P. A. Fisher, R. Shankar, and G. Grinstein, *Phys. Rev. B* **50**, 7526 (1994).
 - [3] A. Giuliani, V. Mastropietro and M. Porta, *Phys. Rev. B* **83**, 195401 (2011); *Comm. Math. Phys.* **311**, 317 (2012).
 - [4] V. Juričić, O. Vafek and I. F. Herbut, *Phys. Rev. B* **82**, 235402 (2010); I. F. Herbut, V. Juričić, O. Vafek and M. J. Case, preprint, arXiv:0809.0725.
 - [5] I. F. Herbut, V. Juričić and O. Vafek, *Phys. Rev. Lett* **100**, 046403 (2008).
 - [6] J. González, F. Guinea and M. A. H. Vozmediano, *Phys. Rev. B* **59**, R2474 (1999).
 - [7] E. G. Mishchenko, *Europhys. Lett.* **83**, 17005 (2008).
 - [8] D. E. Sheehy and J. Schmalian, *Phys. Rev. B* **80**, 193411 (2009).
 - [9] I. Sodemann and M. M. Fogler, *Phys. Rev. B* **86**, 115408 (2012).
 - [10] B. Rosenstein, M. Lewkowicz, T. Maniv, *Phys. Rev. Lett.* **110**, 066602 (2013).
 - [11] J. Gonzalez, F. Guinea and M. A. H. Vozmediano, *Nucl. Phys. B* **424**, 595 (1994).
 - [12] A. Giuliani, V. Mastropietro and M. Porta, *Phys. Rev. B* **82**, 121418 (2010); *Ann. of Phys.* **327**, 461 (2012)
 - [13] A. Giuliani, V. Mastropietro, *Phys. Rev. B* **85**, 045420 (2012)
 - [14] I. F. Herbut, *Phys. Rev. Lett.* **79**, 3502 (1997); *ibid.* **81**, 3916 (1998).
 - [15] R. D. Pisarski, *Phys. Rev. D* **29**, 2423 (1984). T. W. Appelquist, D. Nash, L. C. R. Wijewardhana, *Phys. Rev. Lett.* **60**, 2575 (1988).
 - [16] E. V. Gorbar, V. P. Gusynin, and V. P. Miransky, *Phys. Rev. D* **64**, 105028 (2001).
 - [17] I. F. Herbut, *Phys. Rev. Lett.* **87**, 137004 (2001).
 - [18] S. Teber, *Phys. Rev. D* **86**, 025005 (2012), particularly Eq. (45a).
 - [19] See, I. F. Herbut, *Phys. Rev. Lett.* **88**, 047006 (2002); *Phys. Rev. B* **66**, 094504 (2002), and references therein.
 - [20] I. F. Herbut and Z. Tešanović, *Phys. Rev. Lett.* **76**, 4588 (1996); *ibid.* **78**, 980 (1997).
 - [21] I. F. Herbut, *Phys. Rev. Lett.* **94**, 237001 (2005).