

JAFFARD FAMILIES AND LOCALIZATIONS OF STAR OPERATIONS

DARIO SPIRITO

ABSTRACT. We generalize the concept of localization of a star operation to flat overrings; subsequently, we investigate the possibility of representing the set $\text{Star}(R)$ of star operations on R as the product of $\text{Star}(T)$, as T ranges in a family of overrings of R with special properties. We then apply this method to study the set of star operations on a Prüfer domain R , in particular the set of stable star operations and the star-class groups of R .

1. INTRODUCTION

Recently, the study of star operations, initiated by the works of Krull [26] and Gilmer [16, Chapter 32], has focused on studying the whole set $\text{Star}(R)$ of star operations on R , and in particular its cardinality. Using as a starting point the characterization of domains with $|\text{Star}(R)| = 1$ due to Heinzer [19], Houston, Mimouni and Park have devoted a series of papers [21, 22, 23, 24] to this study, obtaining, among other results, a characterization of Prüfer domains with two star operations [21, Theorem 3.3] and the precise determination of $|\text{Star}(R)|$ on some classes of one-dimensional Noetherian domains [22, 24]. Their work is based – at least partly – on the concept of *localization* of finite-type star operations to localizations of the ring.

The purpose of this paper is to generalize the concept of localization of a star operation $*$, by avoiding (when possible) the hypothesis that $*$ is of finite type and by considering, instead of localizations, flat overrings of the base ring R . In particular, we will prove that, if R admits a family of overrings with certain properties (precisely, a *Jaffard family* [13, Section 6.3]) then $\text{Star}(R)$ can be represented as a cartesian product of $\text{Star}(T)$, as T ranges in this family, and that this representation preserves the main properties of the star operations.

We then specialize to the case of Prüfer domain, when this approach is complemented by the possibility, in certain cases, to link star operations on R with star operations on a quotient of R . This method allows one to obtain a better grasp of several properties, like being a stable

Date: October 5, 2016.

2010 Mathematics Subject Classification. 13A15, 13C20, 13F05, 13F30, 13G05.

Key words and phrases. Star operations, overrings, Jaffard families, Prüfer domains, class group.

operation (Proposition 6.11), and to describe the star-class group of R in terms of the class groups of some localizations of R .

2. PRELIMINARIES AND NOTATION

Let R be an integral domain with quotient field K , and denote by $\mathcal{F}(R)$ the set of fractional ideals of R . A *star operation* on R is a map $*$: $\mathcal{F}(R) \rightarrow \mathcal{F}(R)$, $I \mapsto I^*$ such that, for every $I, J \in \mathcal{F}(R)$ and $x \in K$,

- (a) $I \subseteq I^*$;
- (b) $(I^*)^* = I^*$;
- (c) if $I \subseteq J$, then $I^* \subseteq J^*$;
- (d) $R^* = R$;
- (e) $(xI)^* = x \cdot I^*$.

The set of star operations on R is denoted by $\text{Star}(R)$. An ideal I is a **-ideal* if $I = I^*$.

Similarly, a *semistar operation* on R is a map $*$: $\mathbf{F}(R) \rightarrow \mathbf{F}(R)$ (where $\mathbf{F}(R)$ is the set of R -submodules of K) satisfying the previous properties, except for $R^* = R$; if $*$ verifies also the latter, then it is said to be a *(semi)star operation*. We indicate the sets of semistar and (semi)star operations by $\text{SStar}(R)$ and $(\text{S})\text{Star}(R)$, respectively. A *semiprime operation* is a map c , from the set of integral ideals of R to itself, that satisfies the first four properties of star operations and, moreover, such that $xI^* \subseteq (xI)^*$ for every $x \in R$.

A star operation is said to be:

- *of finite type* if, for every I ,

$$I^* = \bigcup \{J^* \mid J \subseteq I, J \text{ is finitely generated}\};$$

- *semifinite* if any proper *-ideal I is contained in a prime *-ideal;
- *stable* if $(I \cap J)^* = I^* \cap J^*$ for all ideals I, J ;
- *spectral* if it is in the form $I^* = \bigcap \{IR_P \mid P \in \Delta\}$ for some $\Delta \subseteq \text{Spec}(R)$; equivalently, $*$ is spectral if and only if it is stable and semifinite [1, Theorem 4];
- *endlich arithmetisch brauchbar* (*eab* for short) if, for every nonzero finitely generated ideals F, G, H such that $(FG)^* \subseteq (FH)^*$, we have $G^* \subseteq H^*$; if this property holds for arbitrary nonzero fractional ideals G, H (but F still finitely generated) then $*$ is said to be *arithmetisch brauchbar* (*ab* for short);
- *Noetherian* if any set $\{I_\alpha \mid \alpha \in A\}$ of proper *-ideals has a maximum, or equivalently if and only if every ascending chain of *-closed ideals stabilizes. (More commonly, under this hypothesis R is said to be **-Noetherian* [33].)

The set of star operations has a natural order, such that $*_1 \leq *_2$ if and only if $I^{*1} \subseteq I^{*2}$ for every ideal I , or equivalently if and only if every $*_2$ -closed ideal is also $*_1$ -closed. Under this order, $\text{Star}(R)$

becomes a complete lattice, where the minimum is the identity (usually denoted by d) and the maximum the v -operation (or *divisorial closure*) $I \mapsto (R : (R : I))$.

If R is an integral domain, an *overring* of R is a ring T contained between R and its quotient field K . A family Θ of overrings of R is *locally finite* (or *of finite character*) if every $x \in K \setminus \{0\}$ (or, equivalently, every $x \in R \setminus \{0\}$) is a nonunit in only finitely many $T \in \Theta$. The ring R itself is said to be of finite character if $\{R_M : M \in \text{Max}(R)\}$ is a family of finite character.

A flat overring of R is an overring that is flat as a R -module. If T is a flat overring, then $(I_1 \cap \cdots \cap I_n)T = I_1T \cap \cdots \cap I_nT$ for every $I_1, \dots, I_n \in \mathbf{F}(R)$, and $(I : J)T = (IT : JT)$ for every $I, J \in \mathbf{F}(R)$ with J finitely generated [27, Theorem 7.4] (see also [12, Proposition 2]).

3. EXTENDABLE STAR OPERATIONS

The starting point is the notion of localization of a star operation, originally defined in [21]. We shall adopt a more general and more abstract approach.

Definition 3.1. *Let R be an integral domain and T a flat overring of R . We say that a star operation $* \in \text{Star}(R)$ is extendable to T if the map*

$$(1) \quad \begin{aligned} *_{T}: \mathcal{F}(T) &\longrightarrow \mathcal{F}(T) \\ IT &\longmapsto I^*T \end{aligned}$$

is well-defined (where I is a fractional ideal of R).

Remark 3.2.

- (1) If T is flat over R , then every fractional ideal of T is an extension of a fractional ideal of R (since, if J is an integral ideal of T , $J = (J \cap R)T$); therefore, $*_{T}$ is (potentially) defined on all of $\mathcal{F}(T)$.
- (2) If T is flat over R and P is a prime of R such that $PT \neq T$, then PT is a prime ideal of T . Indeed, let Q be a minimal prime of PT . By the previous point, $Q = (Q \cap R)T$; suppose $P \subsetneq Q \cap R$. By [29, Theorem 2], $T_Q = R_{Q \cap R}$, and thus PT_Q is not minimal over $QT_Q = (Q \cap R)T_Q$, a contradiction. Note that the equality $T_Q = R_{Q \cap R}$ also shows that there is at most one $Q \in \text{Spec}(T)$ over any $P \in \text{Spec}(R)$.
- (3) When $T = S^{-1}R$ is a localization of R and $*$ is of finite type, Definition 3.1 coincides with the definition of $*_S$ given in [21, Proposition 2.4].
- (4) When $T = R_P$ for some $P \in \text{Spec}(R)$, we will sometimes denote $*_{T}$ with $*_P$.

The following proposition shows the basic properties of extendability.

Proposition 3.3. *Let R be an integral domain, let $*$ \in $\text{Star}(R)$ and let T be a flat overring of R .*

- (a) *If $*$ is extendable to T , then $*_T$ is a star operation.*
- (b) *$*$ is extendable to T if and only if $I^*T = J^*T$ whenever $IT = JT$.*
- (c) *The identity star operation d is always extendable, and d_T is the identity on T .*
- (d) *If $*$ is of finite type, then it is extendable to T , and $*_T$ is of finite type.*

Note that, if T is a localization of R , point (d) is proved in [21, Proposition 2.4].

Proof. (a) and (c) are obvious, while (b) is just a reformulation of Definition 3.1.

For (d), by symmetry it is enough to show that $J^*T \subseteq I^*T$, or equivalently that $1 \in (I^*T : J^*T)$. Since $*$ is of finite type,

$$\begin{aligned} (I^*T : J^*T) &= \left(I^*T : \left(\sum_{\substack{L \subseteq J \\ L \text{ finitely generated}}} L^* \right) T \right) = \left(I^*T : \sum_{\substack{L \subseteq J \\ L \text{ fin. gen.}}} L^*T \right) = \\ &= \bigcap_{\substack{L \subseteq J \\ L \text{ fin. gen.}}} (I^*T : L^*T) \supseteq \bigcap_{\substack{L \subseteq J \\ L \text{ fin. gen.}}} (I^* : L^*)T. \end{aligned}$$

By properties of star operations, $(I^* : L^*) = (I^* : L)$; since L is finitely generated and T is flat, it follows that, for every L ,

$$(I^* : L^*)T = (I^* : L)T = (I^*T : LT)$$

which contains 1 since $LT \subseteq JT = IT \subseteq I^*T$. Hence, $1 \in (I^*T : J^*T)$, as requested. \square

Example 3.4. Not every star operation is extendable: let R be an almost Dedekind domain which is not Dedekind (i.e., a one-dimensional non-Noetherian domain such that R_M is a discrete valuation ring for every $M \in \text{Max}(R)$), and let P be a non-finitely generated prime ideal of R . Then P is not divisorial [15, Lemma 4.1.8], and thus the v -operation is not extendable to R_P , since otherwise $(PR_P)^{v_P} = P^v R_P = R_P$, while the unique star operation on R_P is the identity.

Beside being of finite type, extension preserves the main properties of a star operation.

Proposition 3.5. *Let R be a domain and T be a flat overring of R ; suppose $*$ \in $\text{Star}(R)$ is extendable to T . If $*$ is stable (respectively, spectral, Noetherian) then so is $*_T$.*

Proof. Suppose $*$ is stable, and let $I_1 := J_1T$, $I_2 := J_2T$ be ideals of T , where J_1 and J_2 are ideals of R . Then,

$$\begin{aligned} (I_1 \cap I_2)^{*T} &= (J_1T \cap J_2T)^{*T} = [(J_1 \cap J_2)T]^{*T} = \\ &= (J_1 \cap J_2)^{*T} = (J_1^* \cap J_2^*)T = J_1^*T \cap J_2^*T = I_1^{*T} \cap I_2^{*T} \end{aligned}$$

and thus $*_T$ is stable.

If $*$ is spectral, it is stable, and thus so is $*_T$. Let now I be a proper $*_T$ -closed ideal of T , and let $J := I \cap R$; then, $JT = (I \cap R)T = I$, and thus $J^* \subseteq I^{*T} \cap R = I \cap R = J$, so that J is a $*$ -ideal. By definition, there is a $\Delta \subseteq \text{Spec}(R)$ such that $*$ = $*_\Delta$; hence,

$$J = J^* = \bigcap_{P \in \Delta} JR_P = \bigcap_{P \in \Delta} (I \cap R)R_P = \bigcap_{P \in \Delta} IR_P \cap R_P.$$

In particular, there is a $P \in \Delta$ such that $1 \notin IR_P = ITR_P$; hence, there is a $Q \in \text{Spec}(TR_P)$ such that $ITR_P \subseteq Q$. We claim that $Q_0 := Q \cap T$ is a prime $*_T$ -ideal containing I . Indeed, $I \subseteq ITR_P \cap T \subseteq Q \cap T = Q_0$; moreover, since $Q \cap R = Q_0 \cap R \subseteq P$, $Q_0 = LT$ for some prime ideal L of T contained in P (Remark 3.22), and thus

$$Q_0^{*T} = L^*T \subseteq (LR_P \cap R)T = LT = Q_0.$$

Therefore, $*_T$ is also semifinite, and by [1, Theorem 4] it is spectral.

Suppose $*$ is Noetherian, and let $\{I_\alpha T \mid \alpha \in A\}$ be an ascending chain of $*_T$ -ideals. Then, $\{I_\alpha^* \mid \alpha \in A\}$ is an ascending chain of $*$ -ideals, which has to stabilize at $I_{\bar{\alpha}}$. Hence, the original chain stabilizes at $I_{\bar{\alpha}}T$, and $*_T$ is Noetherian. \square

Extendability works well with the order structure of $\text{Star}(R)$.

Proposition 3.6. *Let R be an integral domain and T be a flat overring of R . Let $*_1, *_2, \{*_\lambda \mid \lambda \in \Lambda\}$ be star operations that are extendable to T .*

- (a) *If $*_1 \leq *_2 \in \text{Star}(R)$, then $(*_1)_T \leq (*_2)_T$.*
- (b) *$*_1 \wedge *_2$ is extendable to T and $(*_1 \wedge *_2)_T = (*_1)_T \wedge (*_2)_T$.*
- (c) *If each $*_\lambda$ is of finite type, then $\sup_\lambda *_\lambda$ is extendable to T and $(\sup_\lambda *_\lambda)_T = \sup_\lambda (*_\lambda)_T$.*

Proof. (a) If $*_1 \leq *_2$, then $I^{*1} \subseteq I^{*2}$ for every fractional ideal I , and thus $(I^{*1}T) \subseteq (I^{*2}T)$. Using the definition of $*_T$, we get $(*_1)_T \leq (*_2)_T$.

(b) Let I be an ideal of R . By definition, $I^{*1 \wedge *2} = I^{*1} \cap I^{*2}$, so that

$$\begin{aligned} (IT)^{(*1 \wedge *2)_T} &= (I^{*1 \wedge *2})T = (I^{*1} \cap I^{*2})T = \\ &= I^{*1}T \cap I^{*2}T = (IT)^{(*1)_T} \cap (IT)^{(*2)_T} = (IT)^{(*1)_T \wedge (*2)_T} \end{aligned}$$

and thus $(*_1 \wedge *_2)_T = (*_1)_T \wedge (*_2)_T$.

(c) Let $* := \sup_\lambda *_\lambda$. Since each $*_\lambda$ is of finite type, so is $*$ [2, p.1628], and thus $*$ is extendable to T by Proposition 3.3(d). Moreover, again by [2, p.1628], $I^* = \sum I^{*1 \circ \dots \circ *n}$, as $(*_1, \dots, *_n)$ ranges among the finite

strings of elements of $\{*\lambda \mid \lambda \in \Lambda\}$ (here $*_1 \circ \dots \circ *_n$ indicates simply the composition of $*_1, \dots, *_n$); therefore,

$$I^*T = \left(\sum I^{*1 \circ \dots \circ *_n} \right) T = \sum I^{*1 \circ \dots \circ *_n} T.$$

We claim that $I^{*1 \circ \dots \circ *_n} T = (IT)^{(*1)T \circ \dots \circ (*n)T}$; we proceed by induction. The case $n = 1$ is just the definition of the extension; suppose the claim holds for $m < n$. Then,

$$I^{*1 \circ \dots \circ *_n} T = (I^{*1})^{*2 \circ \dots \circ *_n} T = (I^{*1} T)^{(*2)T \circ \dots \circ (*n)T} = (IT)^{(*1)T \circ \dots \circ (*n)T}$$

as claimed. Thus,

$$I^*T = \sum (IT)^{(*1)T \circ \dots \circ (*n)T} = (IT)^{\sup_{\lambda} (*\lambda)T}$$

the last equality coming from [2, p.1628] and Proposition 3.3(d). Hence, $* = \sup_{\lambda} (*\lambda)_T$. \square

Extendability is also transitive:

Proposition 3.7. *Let R be a domain and $T_1 \subseteq T_2$ be two flat overrings of R . If $* \in \text{Star}(R)$ is extendable to T_1 and $*_{T_1}$ is extendable to T_2 , then $*$ is extendable to T_2 , and $*_{T_2} = (*_{T_1})_{T_2}$.*

Proof. Note first that if T_2 is flat over R then it is flat over T_1 , and thus it makes sense to speak of the extendability of $*_{T_1}$. For every ideal I of R , we have

$$I^*T_2 = (I^*T_1)T_2 = (IT_1)^{*_{T_1}}T_2 = (IT_1T_2)^{(*_{T_1})_{T_2}} = (IT_2)^{(*_{T_1})_{T_2}}$$

and thus if $IT_2 = JT_2$ then $I^*T_2 = J^*T_2$, so that $*$ is extendable to T_2 . The previous calculation also shows that $*_{T_2} = (*_{T_1})_{T_2}$. \square

Proposition 3.8. *Let R be an integral domain and T be a flat overring of R . Let $\Delta := \{M \cap R \mid M \in \text{Max}(T)\}$. If $* \in \text{Star}(R)$ is extendable to R_P , for every $P \in \Delta$, then it is extendable to T .*

Proof. Let I, J be ideals of R such that $IT = JT$. Let $P \in \Delta$ and let M be the (necessarily unique – see Remark 3.2(2)) maximal ideal of T such that $M \cap R = P$. Then, $T_M = R_P$, and since $*$ is extendable to R_P we have $I^*R_P = J^*R_P$. It follows that

$$I^*T = \bigcap_{P \in \Delta} I^*R_P = \bigcap_{P \in \Delta} J^*R_P = J^*T,$$

and thus $*$ is extendable to T . \square

Corollary 3.9. *Let R be a domain, and let $* \in \text{Star}(R)$. The following are equivalent:*

- (i) $*$ is extendable to R_P , for every $P \in \text{Spec}(R)$;
- (ii) $*$ is extendable to every flat overring of R .

Note that condition (i) of the above corollary cannot be replaced by the version that considers only maximal ideals of T : indeed, if (R, M) is local, then clearly every star operation is extendable to R_M , but it would be implausible that every star operation is extendable to every localization. We can build an explicit counterexample tweaking slightly [21, Remark 2.5(3)]. Let $R := \mathbb{Z}_{p\mathbb{Z}} + X\mathbb{Q}(\sqrt{2})[[X]]$ (where p is a prime number). Then, R is a two-dimensional local domain, with maximal ideal $M := p\mathbb{Z}_{p\mathbb{Z}} + X\mathbb{Q}(\sqrt{2})[[X]]$; let $P := X\mathbb{Q}(\sqrt{2})[[X]]$. We claim that the v -operation is not extendable to $R_P = \mathbb{Q} + P$. Let $A := X(\mathbb{Q} + P)$ and $B := XR$: then, $AR_P = BR_P = A$, but $A^v R_P = P$ while $B^v R_P = BR_P \neq P$.

4. JAFFARD FAMILIES

The concept of Jaffard family was introduced and studied in [13, Section 6.3].

Definition 4.1. *Let R be a domain and Θ be a set of overrings of R such that the quotient field of R is not in Θ . We say that Θ is a Jaffard family on R if, for every integral ideal I of R ,*

- $R = \bigcap_{T \in \Theta} T$;
- Θ is locally finite;
- $I = \prod_{T \in \Theta} (IT \cap R)$;
- if $T \neq S$ are in Θ , then $(IT \cap R) + (IS \cap R) = R$.

We say that an overring T of R is a Jaffard overring of R if T belongs to a Jaffard family of R .

Note that, by the second axiom, if $I \neq (0)$ then $IT = T$ for all but finitely many $T \in \Theta$, so that the product $I = \prod_{T \in \Theta} (IT \cap R)$ is finite.

The next propositions collect the properties of Jaffard families that we will be using.

Proposition 4.2 [13, Theorem 6.3.1]. *Let R be an integral domain with quotient field K , and let Θ be a Jaffard family on R . For each $T \in \Theta$, let $\Theta^\perp(T) := \bigcap \{U \in \Theta \mid U \neq T\}$.*

- (a) Θ is complete (i.e., $I = \bigcap \{IT \mid T \in \Theta\}$ for every ideal I of R).
- (b) For each $P \in \text{Spec}(R)$, $P \neq (0)$, there is a unique $T \in \Theta$ such that $PT \neq T$.
- (c) For each $T \in \Theta$, both T and $\Theta^\perp(T)$ are flat over R .
- (d) For each $T \in \Theta$, we have $T \cdot \Theta^\perp(T) = K$.

Proposition 4.3. *Let Θ be a family of flat overrings of the domain R , and let K be the quotient field of R . Then, Θ is a Jaffard family if and only if it is complete, locally finite and $TS = K$ for all $T, S \in \Theta$, $T \neq S$.*

Proof. If Θ is a Jaffard family, the properties follow by the definition and Proposition 4.2. Conversely, suppose Θ verifies the three properties, let $I \neq (0)$ be an ideal of R and let $T \neq S$ be members of Θ . If $IT \cap R$ and $IS \cap R$ are not coprime, then there would be a prime P of R containing both; since Θ is complete, it would follow that both $IT \cap R$ and $IS \cap R$ survive in some $A \in \Theta$. In particular, without loss of generality, $A \neq T$; however,

$$(IT \cap R)A = ITA \cap A = IK \cap A = A,$$

a contradiction. Therefore, $(IT \cap R) + (IS \cap R) = R$. Moreover, $I = \bigcap \{IT \cap R \mid T \in \Theta\} = (IT_1 \cap R) \cap \cdots \cap (IT_n \cap R)$ by local finiteness; since the $IT_i \cap R$ are coprime, their intersection is equal to their product, and thus $I = (IT_1 \cap R) \cdots (IT_n \cap R)$. \square

Remark 4.4. Any Jaffard family Θ defines a partition on $\text{Max}(R)$, where each class is composed by the $M \in \text{Max}(R)$ such that $MT \neq T$ for some fixed $T \in \Theta$. In particular, $T = \bigcap R_M$, as M ranges in the class relative to T ; hence, different Jaffard families define different partitions. In particular, a local domain has only one Jaffard family, namely $\{R\}$, and a semilocal domain has only a finite number of Jaffard families.

However, not every partition of $\text{Max}(R)$ can arise in this way. For example, let Θ be a Jaffard family and let $M, N \in \text{Max}(R)$; by Proposition 4.2(b), there are unique overrings $T, U \in \Theta$ such that $MT \neq T$ and $NU \neq U$. If there is a nonzero prime $P \subseteq M \cap N$, then $PT \neq T$ and $PU \neq U$; therefore, again by Proposition 4.2(b), it must be $T = U$.

A *h-local domain* is an integral domain R such that $\text{Max}(R)$ is locally finite and such that every prime ideal P is contained in only one maximal ideal. In this case, $\{R_M \mid M \in \text{Max}(R)\}$ is a Jaffard family of R ; conversely, if $\{R_M \mid M \in \text{Max}(R)\}$ is a Jaffard family, then $\text{Max}(R)$ is locally finite (by definition) and each prime is contained in only one maximal ideal (by Proposition 4.2(b)), and thus R is *h-local*. Many properties of the Jaffard families can be seen as generalizations of the corresponding properties of *h-local domains*; the following proposition is an example (compare [28, Proposition 3.1]).

Proposition 4.5. *Let R be a domain and T be a Jaffard overring of R . Then:*

- (a) *for every family $\{X_\alpha : \alpha \in A\}$ of R -submodules of K with nonzero intersection, we have $(\bigcap_{\alpha \in A} X_\alpha)T = \bigcap_{\alpha \in A} X_\alpha T$;*
- (b) *if $\{I_\alpha : \alpha \in A\}$ is a family of integral ideals of R with nonzero intersection such that $(\bigcap_{\alpha \in A} I_\alpha)T \neq T$, then $I_{\bar{\alpha}}T \neq T$ for some $\bar{\alpha} \in A$.*

Proof. (a) Let Θ be a Jaffard family of R such that $T \in \Theta$. Then, by the flatness of T ,

$$\begin{aligned} \left(\bigcap_{\alpha \in A} X_\alpha \right) T &= \left(\bigcap_{\alpha \in A} \bigcap_{U \in \Theta} X_\alpha U \right) T = \left(\bigcap_{U \in \Theta} \bigcap_{\alpha \in A} X_\alpha U \right) T = \\ &= \left(\bigcap_{U \in \Theta \setminus \{T\}} \bigcap_{\alpha \in A} X_\alpha U \right) T \cap \bigcap_{\alpha \in A} X_\alpha T = K \cap \bigcap_{\alpha \in A} X_\alpha T \end{aligned}$$

since $\bigcap_{U \in \Theta \setminus \{T\}} \bigcap_{\alpha \in A} X_\alpha U$ is a $\Theta^\perp(T)$ -module, and thus its product with T is equal to K by Proposition 4.2(d).

(a \implies b). Suppose $(\bigcap_{\alpha \in A} I_\alpha) T \neq T$. Since $(\bigcap_{\alpha \in A} I_\alpha) T \subseteq T$, then 1 is not contained in the left hand side. By (a), 1 is not contained in $\bigcap_{\alpha \in A} I_\alpha T$, i.e., there is a $\bar{\alpha}$ such that $1 \notin I_{\bar{\alpha}} T$, and thus $I_{\bar{\alpha}} T \neq T$. \square

5. JAFFARD FAMILIES AND STAR OPERATIONS

The reason why we introduced Jaffard families is that they provide a way to decompose $\text{Star}(R)$ as a product of spaces of star operations of overrings of T . Before reaching this objective (Theorem 5.4) we show that weaker properties can lead to a decomposition of at least a subset of $\text{Star}(R)$.

Proposition 5.1. *Let R be an integral domain with quotient field K . Let Θ be a set of flat overrings of R such that $R = \bigcap \{T \mid T \in \Theta\}$ and such that $AB = K$ whenever $A, B \in \Theta$ and $A \neq B$. Then, there is an injective order-preserving map*

$$\begin{aligned} \rho_\Theta: \prod_{T \in \Theta} \text{Star}(T) &\longrightarrow \text{Star}(R) \\ (*^{(T)})_{T \in \Theta} &\longmapsto \bigwedge_{T \in \Theta} *^{(T)}, \end{aligned}$$

where $\bigwedge_{T \in \Theta} *^{(T)}$ is the map such that

$$I \mapsto \bigcap_{T \in \Theta} (IT)^{*^{(T)}}$$

for every fractional ideal I of R .

Proof. Let $(*_T)_{T \in \Theta} \in \prod_{T \in \Theta} \text{Star}(T)$, and let $* := \rho_\Theta((*_T)_{T \in \Theta})$. Since $\bigcap_{T \in \Theta} T = R$, the map $*$ is a star operation; moreover, it is clear that if $*_1^{(T)} \leq *_2^{(T)}$ for all T then $\rho_\Theta(*_1^{(T)}) \leq \rho_\Theta(*_2^{(T)})$. Hence, ρ_Θ is well-defined and order-preserving; we need to show that it is injective.

Suppose it is not; then, $* := \rho_\Theta(*_1^{(T)}) = \rho_\Theta(*_2^{(T)})$ for some families of star operations such that $*_1^{(U)} \neq *_2^{(U)}$ for some $U \in \Theta$. There is an

integral ideal J of U such that $J^{*1^{(U)}} \neq J^{*2^{(U)}}$; let $I := J \cap R$. Since U is flat, for both $i = 1$ and $i = 2$ we have

$$I^*U = \left[\bigcap_{T \in \Theta} (IT)^{*i^{(T)}} \right] U = (IU)^{*i^{(U)}} U \cap \left[\bigcap_{T \in \Theta \setminus \{U\}} (IT)^{*i^{(T)}} \right] U.$$

If $T \neq U$, then, since T is flat,

$$(IT)^{*i^{(T)}} = ((J \cap R)T)^{*i^{(T)}} = (JT \cap T)^{*i^{(T)}}.$$

However, $JT = JUT = K$ since $UT = K$ (by hypothesis); therefore, $(IT)^{*i^{(T)}} = T$, and (since $I \subseteq U$)

$$\begin{aligned} I^*U &= (IU)^{*i^{(U)}} U \cap \left[\bigcap_{T \in \Theta \setminus \{U\}} T \right] U = (IU)^{*i^{(U)}} U \cap \left[\bigcap_{T \in \Theta} T \right] U = \\ &= (IU)^{*i^{(U)}} \cap RU = (IU)^{*i^{(U)}} = J^{*i^{(U)}} \end{aligned}$$

for both $i = 1$ and $i = 2$. However, this contradicts the choice of J ; hence, ρ_Θ is injective. \square

If Θ is a Jaffard family, the previous proposition can be strengthened. We need two lemmas.

Lemma 5.2. *Let R be a domain with quotient field K , and let Θ be a Jaffard family on R . For every $U \in \Theta$, let J_U be a U -submodule of K , and define $J := \bigcap_{U \in \Theta} J_U$. If $J \neq (0)$, then for every $T \in \Theta$ we have $JT = J_T$.*

Proof. By Proposition 4.5(a), we have

$$JT = \left(\bigcap_{U \in \Theta} J_U \right) T = \bigcap_{U \in \Theta} J_U T.$$

If $U \neq T$, then $J_U T = J_U UT = J_U K = K$; therefore, $JT = J_T T = J_T$. \square

The next lemma can be seen as a generalization of [13, Theorem 6.2.2(2)] and [6, Lemma 2.3].

Lemma 5.3. *Let R be an integral domain, T be a Jaffard overring of R , and let $I, J \in \mathbf{F}(R)$ such that $(I : J) \neq (0)$. Then, $(I : J)T = (IT : JT)$.*

Proof. It is enough to note that $(I : J) = \bigcap_{j \in J} j^{-1}I \neq (0)$, and apply Proposition 4.5(a). \square

Theorem 5.4. *Let R be an integral domain and let Θ be a Jaffard family on R . Then, every $*$ \in $\text{Star}(R)$ is extendable to every $T \in \Theta$, and the maps*

$$\begin{aligned} \lambda_{\Theta}: \text{Star}(R) &\longrightarrow \prod_{T \in \Theta} \text{Star}(T) & \rho_{\Theta}: \prod_{T \in \Theta} \text{Star}(T) &\longrightarrow \text{Star}(R) \\ * &\longmapsto (*_T)_{T \in \Theta} & \text{and} & & (*^{(T)})_{T \in \Theta} &\longmapsto \bigwedge_{T \in \Theta} *^{(T)} \end{aligned}$$

(where $\bigwedge_{T \in \Theta} *^{(T)}$ is defined as in Proposition 5.1) are order-preserving bijections between $\text{Star}(R)$ and $\prod\{\text{Star}(T) \mid T \in \Theta\}$.

Proof. We first show that every $*$ \in $\text{Star}(R)$ is extendable. Let $T \in \Theta$ and let I, J be ideals of R such that $IT = JT$. Then, using Lemma 5.3, we have

$$(I^*T : J^*T) = (I^* : J^*)T = (I^* : J)T = (I^*T : JT)$$

and, since $JT = IT \subseteq I^*T$, we have $1 \in (I^*T : J^*T)$, so that $J^*T \subseteq I^*T$. Symmetrically, $I^*T \subseteq J^*T$, and hence $J^*T = I^*T$. By Proposition 3.3(b), $*_T$ is well-defined, and $*$ is extendable to T ; in particular, λ_{Θ} is well-defined.

Moreover, for every $*$ \in $\text{Star}(R)$, we have

$$I^* = \bigcap_{T \in \Theta} I^*T = \bigcap_{T \in \Theta} (IT)^{*T}$$

using the completeness of Θ in the first equality and the definition of extension in the second. Thus, $*$ $=$ $\rho_{\Theta} \circ \lambda_{\Theta}(*)$, i.e., $\rho_{\Theta} \circ \lambda_{\Theta}$ is the identity. It follows that λ_{Θ} is injective and ρ_{Θ} is surjective. But ρ_{Θ} is injective by Proposition 5.1, so λ_{Θ} and ρ_{Θ} must be bijections. \square

The second part of the following corollary is a generalization of [22, Theorem 2.3].

Corollary 5.5. *Let R be a one-dimensional integral domain.*

- (a) $|\text{Star}(R)| \geq \prod\{|\text{Star}(R_M)| : M \in \text{Max}(R)\}$;
- (b) *if R is of finite character (for example, if R is Noetherian), then $|\text{Star}(R)| = \prod\{|\text{Star}(R_M)| : M \in \text{Max}(R)\}$.*

Proof. If $M \neq N$ are maximal ideals of R , then $R_MR_N = K$, since both M and N have height 1. By Proposition 5.1, there is an injective map from $\text{Star}(R)$ to the product $\prod \text{Star}(R_M)$, which in particular implies the first inequality.

If, moreover, R is one-dimensional and of finite character, then $\{R_M \mid M \in \text{Max}(R)\}$ is a Jaffard family, and the claim follows by applying Theorem 5.4. \square

The bijections ρ_{Θ} and λ_{Θ} respect the properties of star operations; see the following Proposition 5.10 for the eab case.

Theorem 5.6. *Let R be a domain and Θ be a Jaffard family on R , and let $*$ \in $\text{Star}(R)$. Then, $*$ is of finite type (respectively, semifinite, stable, spectral, Noetherian) if and only if $*_T$ is of finite type (resp., semifinite, stable, spectral, Noetherian) for every $T \in \Theta$.*

Proof. By Propositions 3.3(d) and 3.5, if $*$ is of finite type, stable, spectral or Noetherian so is $*_T$. If $*$ is semifinite, let I be a $*_T$ -closed ideal of T , and let $J := I \cap R$. Then $JT = I$, and $J^* \subseteq I^{*T} \cap R = J$, so that there is a prime ideal $Q \supseteq J$ such that $Q^* = Q$. For every $U \in \Theta$, $U \neq T$, we have $JU = U$; hence $QU = U$, and thus $QT \neq T$; moreover, since R is flat, QT is prime (Remark 3.2(2)). Therefore, $(QT)^{*T} = Q^*T = QT$ is a proper prime $*_T$ -ideal containing I , and $*_T$ is semifinite.

Let now $* := \rho_{\Theta}(*^{(T)})$.

If each $*^{(T)}$ is of finite type, then $*$ is of finite type by [2].

Suppose each $*^{(T)}$ is semifinite and $I = I^*$ is a proper ideal of R . Then, $1 \notin I$, so there is a $T \in \Theta$ such that $(IT)^{*^{(T)}} \neq T$, and thus there is a prime ideal P of T containing IT such that $P = P^{*^{(T)}}$. If $Q := P \cap R$, then

$$Q^* \subseteq (QT)^{*^{(T)}} \cap R \subseteq P^{*^{(T)}} \cap R = Q,$$

so that Q is a $*$ -prime ideal of R containing I .

If each $*^{(T)}$ is stable, then, given ideal I, J of R , we have

$$(I \cap J)^* = \bigcap_{T \in \Theta} ((I \cap J)T)^{*^{(T)}} = \bigcap_{T \in \Theta} (IT)^{*^{(T)}} \cap \bigcap_{T \in \Theta} (JT)^{*^{(T)}} = I^* \cap J^*.$$

Hence, $*$ is stable. The case of spectral star operation follows since $*$ is spectral if and only if it is stable and semifinite [1, Theorem 4].

Suppose now $*^{(T)}$ is Noetherian for every $T \in \Theta$ and let $\{I_{\alpha} : \alpha \in A\}$ be an ascending chain of $*$ -ideals. If $I_{\alpha} = (0)$ for every α we are done. Otherwise, there is a $\bar{\alpha}$ such that $I_{\bar{\alpha}} \neq (0)$, and thus $I_{\bar{\alpha}}$ (and, consequently, every I_{α} for $\alpha > \bar{\alpha}$) survives in only a finite number of elements of Θ , say T_1, \dots, T_n . For each $i \in \{1, \dots, n\}$, the set $\{I_{\alpha}T_i\}$ is an ascending chain of $*^{(T_i)}$ -ideals, and thus there is a α_i such that $I_{\alpha}T_i = I_{\alpha_i}T_i$ for every $\alpha \geq \alpha_i$.

Let thus $\tilde{\alpha} := \max\{\bar{\alpha}, \alpha_i : 1 \leq i \leq n\}$. For every $\beta \geq \tilde{\alpha}$, we have $I_{\beta}T_i = I_{\alpha_i}T_i = I_{\tilde{\alpha}}T_i$, while, if $T \neq T_i$ for every i , then $I_{\beta}T = T = I_{\tilde{\alpha}}T$ since $\beta \geq \bar{\alpha}$. Therefore, $I_{\beta} = \bigcap_{T \in \Theta} I_{\beta}T = \bigcap_{T \in \Theta} I_{\tilde{\alpha}}T = I_{\tilde{\alpha}}$ and the chain $\{I_{\alpha}\}$ stabilizes. \square

Corollary 5.7. *Let R be a domain and Θ be a Jaffard family on R . If every $T \in \Theta$ is Noetherian, so is R .*

Proof. A domain A is Noetherian if and only if the identity star operation $d^{(A)}$ is Noetherian. If every $T \in \Theta$ is Noetherian, each d_T is a Noetherian star operation, and thus (by Theorem 5.6) $\rho_{\Theta}(d_T)$ is

Noetherian. However, by Theorem 5.4, $\rho_{\Theta}(d_T) = d_R$, and thus R is a Noetherian domain. \square

Lemma 5.8. *Let R be an integral domain and let T be a Jaffard overring of R . For all nonzero integral ideals I, J of T ,*

$$(I \cap R)(J \cap R) = IJ \cap R.$$

Proof. Let Θ be a Jaffard family containing T . Since Θ is complete, it is enough to show that they are equal when localized on every $U \in \Theta$. We have

$$(I \cap R)(J \cap R)U = (IU \cap U)(JU \cap U) = \begin{cases} IJ & \text{if } U = T \\ U & \text{if } U \neq T \end{cases}$$

while

$$(IJ \cap R)U = IJU \cap U = \begin{cases} IJ & \text{if } U = T \\ U & \text{if } U \neq T \end{cases}$$

and thus $(I \cap R)(J \cap R) = IJ \cap R$. \square

Lemma 5.9. *Let R be an integral domain, T a Jaffard overring of R , and let I be a finitely generated integral ideal of T . Then, $I \cap R$ is finitely generated (over R).*

Proof. Let $S := \Theta^{\perp}(T)$, where Θ is a Jaffard family to which T belongs. Then, by Proposition 4.2, $(I \cap R)S = IS \cap S = ITS \cap S = S$, and thus there are $i_1, \dots, i_n \in I \cap R$, $s_1, \dots, s_n \in S$ such that $1 = i_1s_1 + \dots + i_ns_n$; let $I_0 := (i_1, \dots, i_n)$.

Let x_1, \dots, x_m be the generators of I in T . Since $(I \cap R)T = IT = I$, for every x_i there are $j_{1i}, \dots, j_{n_i i} \in I \cap R$, $t_{1i}, \dots, t_{n_i i} \in T$ such that $x_i = j_{1i}t_{1i} + \dots + j_{n_i i}t_{n_i i}$; let $I_i := (j_{1i}, \dots, j_{n_i i})$. Then, $J := I_0 + I_1 + \dots + I_n$ is a finitely generated ideal contained in $I \cap R$ (since it is generated by elements of $I \cap R$) such that $(I \cap R)T \subseteq JT$ and $(I \cap R)S \subseteq JS$; thus, $I \cap R \subseteq J$. Therefore, $I \cap R = J$ is finitely generated, as claimed. \square

Proposition 5.10. *Let R be an integral domain and let Θ be a Jaffard family on R . $A * \in \text{Star}(R)$ is eab (resp., ab) if and only if $*_T$ is eab (resp., ab) for every $T \in \Theta$.*

Proof. (\implies). Suppose $(IJ)^{*T} \subseteq (IL)^{*T}$ for some finitely generated ideals I, J, L of T (which we can suppose contained in T). Since

$$(IJ \cap R)^{*T} = ((IJ \cap R)T)^{*T} = (IJ)^{*T}$$

(and the same happens for IL), we have $(IJ \cap R)^{*T} \subseteq (IL \cap R)^{*T}$, and so

$$(IJ \cap R)^{*T} \cap R \subseteq (IL \cap R)^{*T} \cap R.$$

However, both $IJ \cap R$ and $IL \cap R$ survive (among the ideals of Θ) only in T , so that

$$(IJ \cap R)^{*T} \cap R = (IJ \cap R)^{*} = ((I \cap R)(J \cap R))^*$$

by Lemma 5.8, and thus

$$((I \cap R)(J \cap R))^* \subseteq ((I \cap R)(L \cap R))^*.$$

Since I is finitely generated, by Lemma 5.9 so is $I \cap R$; the same happens for $J \cap R$ and $L \cap R$. Hence, since $*$ is eab, $(J \cap R)^* \subseteq (L \cap R)^*$, and thus

$$J^{*T} = (J \cap R)^*T \subseteq (L \cap R)^*T = L^{*T}.$$

Hence, $*_T$ is eab.

(\Leftarrow). Suppose $(IJ)^* \subseteq (IL)^*$. Then, $(IJ)^*T \subseteq (IL)^*T$, i.e., $(IJT)^{*T} \subseteq (ILT)^{*T}$ for every $T \in \Theta$. Since $*_T$ is eab, this implies that $(JT)^{*T} \subseteq (LT)^{*T}$ for every $T \in \Theta$; since $H^* = \bigcap_{T \in \Theta} (HT)^{*T}$, it follows that $J^* \subseteq L^*$, and $*$ is eab.

The same reasoning applies for the ab case. \square

Following [20], we say that an ideal A is m -canonical if $I = (A : (A : I))$ for every fractional ideal I of R . The following proposition can be seen as a generalization of [20, Theorem 6.7] to domains that are not necessarily integrally closed.

Proposition 5.11. *Let R be a domain. Then R admits an m -canonical ideal if and only if R is h -local, R_M admits an m -canonical ideal for every $M \in \text{Max}(R)$ and $|\text{Star}(R_M)| \neq 1$ for only a finite number of maximal ideals of M .*

Proof. Suppose A is m -canonical. Then R is h -local by [20, Proposition 2.4]; moreover, if I is a R_M -fractional ideal, then $I = JR_M$ for some R -fractional ideal, and thus

$$\begin{aligned} (AR_M : (AR_M : I)) &= (AR_M : (AR_M : JR_M)) = \\ &= (AR_M : (A : J)R_M) = (A : (A : J))R_M = JR_M = I \end{aligned}$$

applying Lemma 5.3 (which is applicable since R h -local implies that R_M is a Jaffard overring of R). If $AR_M = R_M$, it follows that R_M is an m -canonical ideal for R_M , and thus that the v -operation on R_M is the identity, or equivalently that $|\text{Star}(R_M)| = 1$; hence, if $|\text{Star}(R_M)| \neq 1$ then $AR_M \neq R_M$. But this can happen only for a finite number of M , since R is h -local and thus of finite character.

Conversely, suppose that the three hypotheses hold. For every $M \in \text{Max}(R)$, let J_M be an m -canonical ideal of R_M , and define

$$I_M := \begin{cases} R_M & \text{if } |\text{Star}(R_M)| = 1 \\ J_M & \text{if } |\text{Star}(R_M)| > 1 \end{cases}$$

Note that, if $|\text{Star}(R_M)| = 1$, then R_M is m -canonical for R_M , and thus I_M is m -canonical for every M .

The ideal $J := \bigcap_{P \in \text{Max}(R)} I_P$ of R is nonzero, and by Lemma 5.2 $JR_M = I_M$ for every maximal ideal M . If L is an ideal of R then, for

every maximal ideal M ,

$$(J : (J : L))R_M = (JR_M : (JR_M : LR_M)) = (I_M : (I_M : LR_M)) = LR_M,$$

so that

$$(J : (J : L)) = \bigcap_{M \in \text{Max}(R)} (J : (J : L))R_M = \bigcap_{M \in \text{Max}(R)} LR_M = L.$$

Therefore, J is an m -canonical ideal of R . \square

Remark 5.12. The results in Sections 3 and 5 can be generalized in two different directions.

On the one hand, we can consider, instead of star operations, other classes of closure operations, for example semiprime or semistar operations. In both cases, the definitions of extendability and the results in Section 3 carry over without modifications, noting that the equalities $(I^c : J^c) = (I^c : J)$ and $(I^* : J^*) = (I^* : J)$ holds when c and $*$ are, respectively, a semiprime or a semistar operation.

However, the behaviour of these two classes differs when we come to Jaffard families. In one case there is no problem: with the obvious modifications, all result of Section 5 hold for the set $\text{Sp}(R)$ of semiprime operations. For example, this means that we can analyze the structure of the semiprime operation on a Dedekind domain D almost directly from the structure of $\text{Sp}(V)$, for V a discrete valuation ring, shortening the analysis done in [32, Section 3].

The case of semistar operations is much more delicate: indeed, the result corresponding to Theorem 5.4 is *not* true for $\text{SStar}(R)$, meaning that a semistar operation on R may not be extendable to a Jaffard overring T of R . For example, let $*$ be the semistar operation defined by

$$I^* = \begin{cases} I & \text{if } I \in \mathcal{F}(R) \\ K & \text{otherwise.} \end{cases}$$

If $T \neq R$ is a Jaffard overring of R , then it is not a fractional ideal of R (for otherwise $T \cdot \Theta^\perp(T) = K$ would imply $\Theta^\perp(T) = K$); however, we have $RT = TT$, while

$$R^*T = T \neq K = T^*T.$$

Hence, $*$ is not extendable to T . The exact point in which the proof of Theorem 5.4 fails is the possibility of using Lemma 5.3, because the equality $IT = JT$ does not imply that $(I : J) \neq (0)$. However, if we restrict to finite-type semistar operations, the analogue of Theorem 5.4 does hold: indeed, a proof analogous to the one of Proposition 3.3(d) shows that finite-type operations are extendable, and thus the proof of Theorem 5.4 continues without problems.

A second way of generalizing these results is by considering, beyond the order structure, also a *topological* structure on $\text{Star}(R)$: mimicking

the definition of the Zariski topology on $\text{SStar}(R)$ given in [11], we can define a topology on $\text{Star}(R)$ by declaring open the sets of the form

$$V_I := \{ * \in \text{Star}(R) \mid 1 \in I^* \},$$

as I ranges among the fractional ideals of R . In particular, Theorem 5.4 can be interpreted at the topological level: if Θ is a Jaffard family of R , then λ_Θ and ρ_Θ are homeomorphisms between $\text{Star}(R)$ and the space $\prod_{T \in \Theta} \text{Star}(T)$ endowed with the product topology.

6. APPLICATION TO PRÜFER DOMAINS

Theorem 5.4 allows one to split the study of the set $\text{Star}(R)$ of star operations on R into the study of the sets $\text{Star}(T)$, as T ranges among the members of a Jaffard family Θ . Obviously, this result isn't quite useful if we don't know how to find Jaffard families, or if studying $\text{Star}(T)$ is as complex as studying $\text{Star}(R)$. The purpose of this section is to show that, in the case of (some classes of) Prüfer domains, we can resolve the first question, and we can at least make some progress on the second, proving more explicit results on $\text{Star}(R)$. We shall employ a method similar to the one used in [23, Sections 3-5]

Let now R be a Prüfer domain with quotient field K . We say that two maximal ideals M, N are *dependent* if $R_M R_N \neq K$, or equivalently if $M \cap N$ contains a nonzero prime ideal. Since the spectrum of R is a tree, being dependent is an equivalence relation; we indicate the equivalence classes by Δ_λ , as λ ranges in Λ . We also define $T_\lambda := \bigcap \{ R_P \mid P \in \Delta_\lambda \}$. We call the set $\Theta := \{ T_\lambda \mid \lambda \in \Lambda \}$ the *standard decomposition* of R .

Lemma 6.1. *Let R be a finite-dimensional Prüfer domain. Then, $\Delta \subseteq \text{Max}(R)$ is an equivalence class with respect to dependence if and only if $\Delta = V(P) \cap \text{Max}(R)$ for some height-one prime P of R .*

Proof. Suppose $\Delta = V(P) \cap \text{Max}(R)$. If $M, N \in \Delta$, then $P \subseteq M \cap N$; conversely, since P has height 1, $M \in \Delta$ and $Q \subseteq M \cap N$ imply that $P \subseteq Q$ (since the spectrum of R is a tree).

On the other hand, suppose $\Delta = \Delta_\lambda$ for some λ , and let $M, N \in \Delta$. Since $\text{Spec}(R)$ is a tree and $\dim(R) < \infty$, both M and N contain a unique height-one prime, respectively (say) P_M and P_N ; if $P_M \neq P_N$, then $M \cap N$ cannot contain a nonzero prime, and thus M and N are not dependent, against the hypothesis $M, N \in \Delta$. Therefore, the height-1 prime contained in the members of Δ is unique, and $\Delta = V(P) \cap \text{Max}(R)$. \square

Proposition 6.2. *Let R be a Prüfer domain, and suppose that*

- (a) $\text{Max}(R)$ is a Noetherian space; or
- (b) R is semilocal.

Then, the standard decomposition Θ of R is a Jaffard family of R .

Proof. Since R is Prüfer, every overring of R is flat [15, Theorem 1.1.1], and this in particular applies to the $T \in \Theta$.

We claim that, under both hypotheses, if $T = T_\lambda \in \Theta$, then $\text{Spec}(T) = \{PT \mid P \subseteq M \text{ for some } M \in \Delta_\lambda\}$. Indeed, in both cases every Δ_λ is compact: if $\text{Max}(R)$ is Noetherian this is immediate, while if R is semilocal they are finite and thus compact. Hence, the semistar operation $*_\Delta$ is of finite type [14, Corollary 4.6], and $R^{*\Delta} = T$; since the unique finite-type (semi)star operation on a Prüfer domain is the identity (since all finitely generated ideals are invertible), it follows that $*_\Delta$ is just the map $I \mapsto IT$, and thus $QT = T$ if Q is not contained in any $M \in \Delta$. Therefore, no prime ideal P of R survives in two different members of Θ ; thus, $PT_\lambda T_\mu = T_\lambda T_\mu$ if $\lambda \neq \mu$ are in Λ . Hence, $T_\lambda T_\mu = K$.

We need to show that Θ is locally finite. If R is semilocal then Θ is finite, and in particular locally finite; suppose $\text{Max}(R)$ is Noetherian. For every $x \in R$, $x \neq 0$, the ideal xR has only a finite number of minimal primes (this follows, for example, from the proof of [7, Chapter 4, Corollary 3, p.102] or [5, Chapter 6, Exercises 5 and 7]); in particular, since each prime survives in only one $T \in \Theta$, the family Θ is of finite character.

Hence, in both case Θ is a Jaffard family by Proposition 4.3. \square

Remark 6.3.

- (1) If R is a Prüfer domain that is both of finite character and finite-dimensional, then $\text{Spec}(R)$ (and so $\text{Max}(R)$) is Noetherian. Indeed, if I is a nonzero radical ideal of R , then $V(I)$ is finite, and thus every ascending chain of radical ideals must stop; by [5, Chapter 6, Exercise 5], this implies Noetherianity.
- (2) The standard decomposition Θ of R is the “finest” Jaffard family of R , in the sense that the partition of $\text{Max}(R)$ determined by Θ (see Remark 4.4) is the finer partition that can be induced by a Jaffard family; this follows exactly from the definition of the dependence relation.
- (3) In general, the standard decomposition of R need not be a Jaffard family of R . For example, let R be an almost Dedekind domain which is not Dedekind. Since R is one-dimensional, no two maximal ideals are dependent, and thus each T_λ has the form R_M for some maximal ideal M . However, Θ is not a Jaffard family, since it is not locally finite (if it were, R would be a Dedekind domain). Indeed, Example 3.4 shows that not every star operation is extendable to every R_M .

6.1. Cutting the branch. Let R be a finite-dimensional Prüfer domain whose standard decomposition Θ is a Jaffard family. By Lemma 6.1, every $T \in \Theta$ will have a nonzero prime ideal P contained in all its

maximal ideals; moreover, by Remark 6.3(2), T does not admit a further decomposition. On the other hand, it may be possible that T/P has a nontrivial standard decomposition that is still a Jaffard family; thus, if we could relate $\text{Star}(T)$ with $\text{Star}(T/P)$, we could (in principle) simplify the study of $\text{Star}(T)$.

Lemma 6.4. *Let R be a Prüfer domain whose Jacobson radical $\text{Jac}(R)$ contains a nonzero prime ideal. Then, there is a prime ideal $Q \subseteq \text{Jac}(R)$ such that $\text{Jac}(R/Q)$ does not contain nonzero prime ideals.*

Proof. Let $\Delta := \{P \in \text{Spec}(R), P \subseteq \text{Jac}(R)\}$. By hypothesis, Δ contains nonzero prime ideals. Let $Q := \bigcup_{P \in \Delta} P$.

Since R is treed, Δ is a chain; hence, Q is itself a prime ideal, and it is contained in every maximal ideal of R . Suppose $\text{Jac}(R/Q)$ contains a nonzero prime ideal \overline{Q} . Then, $\overline{Q} = Q'/P$ for some prime ideal Q' of R , and Q' is contained in every maximal ideal of R . It follows that $Q \subsetneq Q' \subseteq \text{Jac}(R)$, against the construction of Q . \square

Suppose now that R is a Prüfer domain with quotient field K , and suppose there is a nonzero prime ideal P contained in every maximal ideal of R . Then, we have a quotient map $\phi : R_P \rightarrow R_P/PR_P = k$ that, for every star operation $*$ on R , induces a semistar operation $*_\phi$ on $D := R/P$ defined by

$$I^{*\phi} := \phi(\phi^{-1}(I)^*),$$

such that $D^{*\phi} = D$. Conversely, if \sharp is a star operation on D , then we can construct a star operation \sharp^ϕ on R : indeed, if I is a fractional ideal of R , then I is either divisorial (and so we define $I^{\sharp^\phi} := I$) or there is an $\alpha \in K$ such that $R \subseteq \alpha I \subseteq R_P$ [23, Proposition 2.2(5)]: in the latter case, we define

$$I^{\sharp^\phi} := \alpha^{-1}\phi^{-1}(\phi(\alpha I)^\sharp).$$

Proposition 6.5. *Let R, P, D, ϕ as above. Then, the maps*

$$\begin{array}{ccc} \text{Star}(R) \longrightarrow (\text{S})\text{Star}(R/P) & \text{and} & (\text{S})\text{Star}(R/P) \longrightarrow \text{Star}(R) \\ * \longmapsto *_\phi & & * \longmapsto *^\phi \end{array}$$

are well-defined order-preserving bijections.

Proof. The fact that they are well-defined and bijections follow from [23, Lemmas 2.3 and 2.4]; the fact that they are order-preserving is immediate from the definitions. \square

6.2. h -local Prüfer domains. If R is both a Prüfer domain and a h -local domain, then its standard decomposition $\Theta := \{R_M \mid M \in \text{Max}(R)\}$ is composed by valuation domains, and star operations behave particularly well. We start by re-proving [21, Theorem 3.1] using our general theory.

Proposition 6.6. *Let R be an h -local Prüfer domain, and let \mathcal{M} be the set of nondivisorial maximal ideals of R . Then, $|\text{Star}(R)| = 2^{|\mathcal{M}|}$.*

Proof. By Theorem 5.4, there is an order-preserving bijection between $\text{Star}(R)$ and $\prod\{\text{Star}(R_M) \mid M \in \text{Max}(R)\}$, and a maximal ideal M is divisorial (in R) if and only if MR_M is divisorial (in R_M). Since R_M is a valuation domain, $|\text{Star}(R_M)|$ is equal to 1 if MR_M is divisorial, and to 2 if MR_M is not; the claim follows. \square

It is noted in the proof of [28, Theorem 3.10] that, if R is an h -local Prüfer domain and I, J are divisorial ideals of R , then $I + J$ is also divisorial. We can extend this result to arbitrary star operations; we shall see a similar result in Proposition 7.8.

Proposition 6.7. *Let R be an h -local Prüfer domain, let $*$ \in $\text{Star}(R)$ and let I, J be $*$ -closed ideals. Then, $I + J$ is $*$ -closed.*

Proof. Since R is h -local, $I + J$ is $*$ -closed if and only if $(I + J)R_M$ is $*_M$ -closed for every $M \in \text{Max}(R)$. However, since R_M is a valuation domain, either $IR_M \subseteq JR_M$ or $JR_M \subseteq IR_M$; hence, $(I + J)R_M = IR_M + JR_M$ is equal either to IR_M or to JR_M , both of which are $*_M$ -closed. \square

This result does not hold if we drop the hypothesis that R is h -local: in fact, let $R = \mathbb{Z} + X\mathbb{Q}[[X]]$ and let $R_p := \mathbb{Z}[1/p] + X\mathbb{Q}[[X]]$ for each prime number p . Consider the star operation

$$* : I \mapsto (R : (R : I)) \cap (R_2 : (R_2 : I)) \cap (R_3 : (R_3 : I)).$$

Then, R_2 and R_3 are $*$ -closed; we claim that $R_2 + R_3$ is not. Indeed, if T is equal to R, R_2 or R_3 , then $(T : (R_2 + R_3)) = X\mathbb{Q}[[X]]$, and thus $(R_2 + R_3)^* = \mathbb{Q}[[X]]$; however, $R_2 + R_3 = (\mathbb{Z}[1/2] + \mathbb{Z}[1/3]) + X\mathbb{Q}[[X]]$ does not contain rationals with denominator not divisible by 2 or 3 (for example, $1/5 \notin R_2 + R_3$), and thus $R_2 + R_3 \neq \mathbb{Q}[[X]]$.

The following can be seen as a sort of converse to Proposition 6.7.

Proposition 6.8. *Let R be a Prüfer domain and suppose that R is either:*

- (a) *semilocal; or*
- (b) *locally finite and finite-dimensional.*

Then, the following are equivalent:

- (i) *R is h -local;*
- (ii) *for every $*$ \in $\text{Star}(R)$, $I \in \mathcal{F}(R) \setminus \mathcal{F}^*(R)$ and $J \in \mathcal{F}(R)$, at least one of $I \cap J$ and $I + J$ is not $*$ -closed;*
- (iii) *for every $I \in \mathcal{F}(R) \setminus \mathcal{F}^v(R)$ and $J \in \mathcal{F}(R)$, at least one of $I \cap J$ and $I + J$ is not divisorial.*

Proof. (i \implies ii) For every $M \in \text{Max}(R)$, $(I + J)R_M = IR_M + JR_M = \max\{IR_M, JR_M\}$, while $(I \cap J)R_M = IR_M \cap JR_M = \min\{IR_M, JR_M\}$. Since I is not $*$ -closed, and $\{R_M \mid M \in \text{Max}(R)\}$ is a Jaffard family of

R , there is a maximal ideal N such that IR_N is not $*_N$ -closed; however, at least one of $(I + J)R_N$ and $(I \cap J)R_N$ is equal to IR_N , and thus at least one is not $*_N$ -closed. Therefore, at least one between $I + J$ and $I \cap J$ is not $*$ -closed.

(ii \implies iii) is obvious.

(iii \implies i) Consider the standard decomposition Θ of R ; then, (iii) holds for every member of Θ but, if R is not h -local, there must be a $T \in \Theta$ that is not local. By Lemma 6.4, there is a prime ideal P of T such that $\text{Jac}(T/P)$ does not contain nonzero primes. Let Λ be the standard decomposition of $D := T/P$, let $Z \in \Lambda$ and define $Z' := \bigcap_{W \in \Lambda \setminus \{Z\}} W = \Lambda^\perp(Z)$. We have $Z \cap Z' = D$ and, for every maximal ideal M of D , either $ZD_M = K$ or $Z'D_M = K$. Therefore, $Z + Z' = \bigcap_{M \in \text{Max}(T)} (Z + Z')D_M = K$.

By Proposition 6.5, the v -operation on T correspond to a (semi)star operation on D such that $A^* = K$ if A is not a fractional ideal of D ; therefore, both $\phi^{-1}(Z)$ and $\phi^{-1}(Z')$ are not divisorial, but both $\phi^{-1}(Z \cap Z') = T$ and $\phi^{-1}(Z + Z') = T_P$ are (where $\phi : T \rightarrow D$ is the quotient map). This is a contradiction, and R must be h -local. \square

6.3. Stability. Recall that a star operation $*$ is *stable* if it distributes over finite intersections, i.e., if $(I \cap J)^* = I^* \cap J^*$. In this section, we study stable operations on Prüfer domains; we start with an analogue of Proposition 6.5.

Proposition 6.9. *Preserve the notation and the hypotheses of Proposition 6.5. There is a bijection between $\text{Star}_{st}(R)$ and $\text{Star}_{st}(R/P)$.*

Proof. We first show that the bijections of Proposition 6.5 become bijections on the subsets of stable operations; let thus $*$ be a semistar operation in the first set and \sharp be the corresponding operation on $(S)\text{Star}(R/P)$. Let $\phi : R \rightarrow R/P$ be the quotient map.

Suppose that $*$ is stable and let $I, J \in \mathbf{F}(R/P)$. Then, since ϕ is a bijection between the ideal comprised between P and R_P and $\mathbf{F}(R/P)$,

$$\begin{aligned} (I \cap J)^\sharp &= \phi[\phi^{-1}(I \cap J)^*] = \phi[(\phi^{-1}(I) \cap \phi^{-1}(J))^*] = \\ &= \phi[\phi^{-1}(I)^* \cap \phi^{-1}(J)^*] = \phi(\phi^{-1}(I)^*) \cap \phi(\phi^{-1}(J)^*) = \\ &= I^\sharp \cap J^\sharp. \end{aligned}$$

Therefore, \sharp is stable.

Conversely, suppose \sharp is stable and let $I, J \in \mathcal{F}(R)$. If I and J are divisorial, so is $I \cap J$; hence, $(I \cap J)^* = I \cap J = I^* \cap J^*$. Suppose (without loss of generality) that $I \neq I^v$. Then, there is an α such that $P \subseteq \alpha I \subseteq R_P$. Moreover, since R is Prüfer and P is contained in every maximal ideal of R , every fractional ideal must be comparable with both P and R_P : more precisely, if \mathbf{v} is the valuation relative to R_P , and L is an ideal, then either $\inf \mathbf{v}(L) = 0$ (so that $P \subseteq L \subseteq R_P$), $\inf \mathbf{v}(L)$ exist and has a sign (if positive, $L \subseteq P$, if negative, $R_P \subseteq L$) or $\inf \mathbf{v}(L)$ has no infimum (so that if $\mathbf{v}(L)$ contains negative values

then $R_P \subseteq L$, while $L \subseteq P$ in the other case). Therefore, we can distinguish three cases:

- $\alpha J \subseteq P$: then, $\alpha J \subseteq \alpha I$, and thus $(I \cap J)^* = J^* = I^* \cap J^*$;
- $R_P \subseteq \alpha J$: then, $\alpha I \subseteq \alpha J$, and thus $(I \cap J)^* = I^* = I^* \cap J^*$;
- $P \subseteq \alpha J \subseteq R_P$. Let $I_0 := \alpha I$ and $J_0 := \alpha J$. Then,

$$\begin{aligned} (I_0 \cap J_0)^* &= \phi^{-1}(\phi(I_0 \cap J_0)^\#) = \phi^{-1}(\phi(I_0)^\# \cap \phi(J_0)^\#) = \\ &= \phi^{-1}(\phi(I_0)^\#) \cap \phi^{-1}(\phi(J_0)^\#) = I_0^* \cap J_0^*. \end{aligned}$$

Hence,

$$\begin{aligned} (I \cap J)^* &= \alpha^{-1}(\alpha(I \cap J)^*) = \alpha^{-1}(I_0 \cap J_0)^* = \\ &= \alpha^{-1}(I_0^* \cap J_0^*) = \alpha^{-1}I_0^* \cap \alpha^{-1}J_0^* = I^* \cap J^*. \end{aligned}$$

In all cases, $*$ distributes over finite intersection, and thus $*$ is stable.

Therefore, there is an order-preserving bijection between $\text{Star}_{st}(R)$ and $(S)\text{Star}_{st}(R/P)$. However, for every domain D , the restriction map $(S)\text{Star}_{st}(D) \rightarrow \text{Star}_{st}(D)$ is a bijection (see [9, Discussion after Proposition 3.10] or [10, Proposition 3.4]), and thus $\text{Star}_{st}(R)$ corresponds bijectively with $\text{Star}_{st}(R/P)$. The claim follows. \square

We say that a star (or semistar) operation $*$ *distributes over arbitrary intersections* if, whenever $\{I_\alpha\}_{\alpha \in A}$ is a family of ideals with nonzero intersection, we have $(\bigcap_{\alpha \in A} I_\alpha)^* = \bigcap_{\alpha \in A} I_\alpha^*$.

Lemma 6.10. *If V is a valuation domain, the v -operation distributes over arbitrary intersections.*

Proof. Let $\mathcal{A} := \{I_\alpha\}_{\alpha \in A}$ be a family of ideals of V with nonzero intersection. If \mathcal{A} has a minimum $I_{\bar{\alpha}}$, then $I_{\bar{\alpha}}^v \subseteq I_\beta^v$ for every $\beta \in A$, and thus $(\bigcap_{\alpha \in A} I_\alpha)^v = I_{\bar{\alpha}}^v = \bigcap_{\alpha \in A} I_\alpha^v$.

Suppose \mathcal{A} does not have a minimum: since $(\bigcap_{\alpha \in A} I_\alpha)^v \subseteq I_\alpha^v$ for every $\alpha \in A$, we have $(\bigcap_{\alpha \in A} I_\alpha)^v \subseteq \bigcap_{\alpha \in A} I_\alpha^v$.

Let $x \in \bigcap_{\alpha \in A} I_\alpha^v$: if $x \in \bigcap_{\alpha \in A} I_\alpha$ then $x \in (\bigcap_{\alpha \in A} I_\alpha)^v$. On the other hand, if $x \notin \bigcap_{\alpha \in A} I_\alpha$, then there is an $\bar{\alpha}$ such that $x \in I_{\bar{\alpha}}^v \setminus I_{\bar{\alpha}}$, i.e., $\mathbf{v}(x) = \inf \mathbf{v}(I_{\bar{\alpha}})$ (where \mathbf{v} is the valuation associated to V and $\mathbf{v}(J) := \{\mathbf{v}(j) \mid j \in J\}$). However, since \mathcal{A} has no minimum, there are $\beta, \gamma \in A$ such that $I_\alpha \supsetneq I_\beta \supsetneq I_\gamma$; in particular, $\mathbf{v}(x) > \inf \mathbf{v}(I_\gamma)$, and thus $x \notin I_\gamma^v$, which is absurd. Therefore, $x \in \bigcap_{\alpha \in A} I_\alpha$. \square

The following proposition may also be proved, in a slightly more generalized setting, using a different, more direct, approach; see [31].

Proposition 6.11. *Let R be a Prüfer domain and suppose that R is either:*

- (a) *semilocal; or*
- (b) *locally finite and finite-dimensional.*

Then, every stable star operation $*$ on R is in the form

$$(2) \quad I \mapsto \bigcap_{P \in \text{Max}(R)} (IR_P)^{*(P)},$$

where each $*(P) \in \text{Star}(R_P)$. In particular, $\text{Star}_{st}(R)$ is order-isomorphic to $\prod\{\text{Star}(R_P) \mid P \in \text{Max}(R)\}$.

Proof. For any ring A , let \mathcal{M}_A be the set of maximal ideals of A that are not divisorial.

Suppose first that R is semilocal, and let Δ be the set of star operations defined as in (2). By Lemma 6.10, every star operation in Δ is stable; moreover, a maximal ideal P is $*$ -closed if and only if $*(P)$ is the identity, and thus $|\Delta| = 2^{|\mathcal{M}_R|}$. Since $\text{Star}(R)$ is finite [23, Theorem 5.3], it is enough to show that the cardinalities of Δ and $\text{Star}_{st}(R)$ are equal.

We proceed by induction on $n := |\text{Max}(R)|$; if $n = 1$ the claim follows from Lemma 6.10. Suppose it holds up to $n - 1$.

Let Θ be the standard decomposition of R . If Θ is not trivial, then by the inductive hypothesis the claim holds for every member of Θ ; by Theorem 5.4, $M \in \text{Max}(R)$ is divisorial over R if and only if MT is divisorial over T (where $T \in \Theta$ is such that $MT \neq T$), and thus $|\mathcal{M}_R| = \sum_{T \in \Theta} |\mathcal{M}_T|$. Since, by Theorem 5.6, we have $\text{Star}_{st}(R) \simeq \prod\{\text{Star}_{st}(T) \mid T \in \Theta\}$, it follows that the claim holds also for R .

Suppose Θ is trivial: then, $\text{Jac}(R)$ must contain a nonzero prime ideal P (and, by Lemma 6.4, we can suppose P is maximal with these properties). By Proposition 6.9, $|\text{Star}_{st}(R)| = |\text{Star}_{st}(R/P)|$; moreover, by Proposition 6.5 \mathcal{M}_R and $\mathcal{M}_{R/P}$ have the same cardinality. By the maximality of P , R/P has a nontrivial standard decomposition; by induction, the claim holds for every member of the decomposition, and thus, with the same reasoning as above, we see that $|\text{Star}_{st}(R/P)| = 2^{|\mathcal{M}_{R/P}|}$. Putting all together we have $|\text{Star}_{st}(R)| = 2^{|\mathcal{M}_R|}$ and so $\text{Star}_{st}(R) = \Delta$ holds for every semilocal Prüfer domain.

If R is locally finite and finite-dimensional, then $\text{Star}_{st}(R) = \prod\{\text{Star}_{st}(T) \mid T \in \Theta\}$, where Θ is the standard decomposition of R . Each $T \in \Theta$ is semilocal, and thus we can apply the previous part of the proof; the claim follows. \square

Proposition 6.12. *Let R be a Prüfer domain and suppose that R is either:*

- (a) *semilocal; or*
- (b) *locally finite and finite-dimensional.*

Then, the following are equivalent:

- (i) *R is h -local;*
- (ii) *every star operation on R distributes over arbitrary intersections;*
- (iii) *every star operation on R distributes over finite intersections;*

- (iv) the v -operation on R distributes over arbitrary intersections;
- (v) the v -operation on R distributes over finite intersections;
- (vi) for every fractional ideal I of R , $I^v = \bigcap \{(IR_M)^{v(R_M)} \mid M \in \text{Max}(R)\}$.

Proof. (i \implies ii) follows from Theorem 5.4, Lemma 5.2 and Lemma 6.10, since $\{R_M \mid M \in \text{Max}(R)\}$ is a Jaffard family if R is h -local. (ii \implies iii \implies v) and (ii \implies iv \implies v) are clear, while (v \iff vi) follows from Proposition 6.11; we only have to show that (v \implies i).

Suppose (v) holds and let Θ be the standard decomposition of R . If R is not h -local, then a branch $T \in \Theta$ is not local; the hypotheses on R guarantee that there is a nonzero prime ideal of T contained in every maximal ideal. Therefore, we can apply Lemma 6.4 and find a prime ideal Q such that $\text{Jac}(T/Q)$ contains no prime ideals. By Proposition 6.5, there is an order-preserving bijection between $\text{Star}(T)$ and (S) $\text{Star}(T/Q)$, where the v -operation on T corresponds to the semistar operation $*$ which is the trivial extension of the v -operation on T/Q .

Since $\text{Jac}(T/Q)$ does not contain nonzero primes, T/Q admits a non-trivial Jaffard family Λ ; let $Z \in \Lambda$, and define $Z' := \bigcap_{W \in \Lambda \setminus \{Z\}} W = \Lambda^\perp(Z)$. Then, Z and Z' are not fractional ideals of T/Q , and thus $Z^* = Z'^* = F$, where F is the quotient field of T/Q ; on the other hand, $Z \cap Z' = T/Q$ and thus $(Z \cap Z')^* = T/Q$.

If $\pi : T_Q \longrightarrow T_Q/QT_Q$ is the canonical quotient, it follows that $\pi^{-1}(Z)^v = \pi^{-1}(Z')^v = T_Q$, while $\pi^{-1}(Z \cap Z')^v = \pi^{-1}(T/Q)^v = T^v = T$. Since T is not local, $T \neq T_Q$, and thus v does not distribute over finite intersections, against the hypothesis. \square

7. THE CLASS GROUP

Let $*$ be a star operation on R . An ideal I of R is $*$ -invertible if $(I(R : I))^* = R$; the set of $*$ -invertible $*$ -ideals, indicated with $\text{Inv}^*(R)$, is a group under the natural “ $*$ -product” $I \times_* J \mapsto (IJ)^*$ [25, 17, 34, 18]. Any $*$ -invertible $*$ -ideal is divisorial [34, Theorem 1.1 and Observation C] and, if $*_1 \leq *_2$, there is a natural inclusion $\text{Inv}^{*1}(R) \subseteq \text{Inv}^{*2}(R)$.

Proposition 7.1. *Let R be an integral domain and Θ be a Jaffard family on R . The map*

$$\begin{aligned} \Gamma : \text{Inv}^*(R) &\longrightarrow \bigoplus_{T \in \Theta} \text{Inv}^{*T}(T) \\ I &\longmapsto (IT)_{T \in \Theta} \end{aligned}$$

is well-defined and a group isomorphism.

Proof. Define a map

$$\begin{aligned} \widehat{\Gamma} : \mathcal{F}(R) &\longrightarrow \prod_{T \in \Theta} \mathcal{F}(T) \\ I &\longmapsto (IT)_{T \in \Theta} \end{aligned}$$

For every $*$ -ideal I , $\widehat{\Gamma}(I) = (IT)$ is a sequence such that IT is $*_T$ -closed. Moreover, if I is $*$ -invertible, then $(I(R : I))^* = R$ and thus $(I(R : I)T)^{*T} = T$, so that IT is $*_T$ -invertible. Thus $\widehat{\Gamma}(\text{Inv}^*(R)) \subseteq \prod_{T \in \Theta} \text{Inv}^{*T}(T)$, and indeed $\widehat{\Gamma}(\text{Inv}^*(R)) \subseteq \bigoplus_{T \in \Theta} \text{Inv}^{*T}(T)$ since Θ is locally finite, by Theorem 5.4. Hence, Γ is well-defined, since it is the restriction of $\widehat{\Gamma}$ to $\text{Inv}^*(R)$.

It is straightforward to verify that Γ is a group homomorphism, and since $I = \bigcap_{T \in \Theta} IT$, we have that Γ (or even $\widehat{\Gamma}$) is injective.

We need only to show that Γ is surjective. Let $(I_T) \in \bigoplus_{T \in \Theta} \text{Inv}^{*T}(T)$, and define $I := \bigcap I_T$. Since $I_T = T$ for all but a finite number of elements of Θ , say T_1, \dots, T_n , there are $d_1, \dots, d_n \in R$ such that $d_i I_{T_i} \subseteq T_i$. Defining $d := d_1 \cdots d_n$, we have $dI_T \subseteq T$ for every T , and thus $dI \subseteq \bigcap_{T \in \Theta} T = R$, so that I is indeed a fractional ideal of R . Moreover, since I_T is $*_T$ -closed, $I_T \cap R$ is $*$ -closed, and thus I , being the intersection of a family of $*$ -closed ideals, is $*$ -closed. It is also $*$ -invertible, since

$$(I(R : I))^* = \bigcap_{T \in \Theta} (I(R : I)T)^{*T} = \bigcap_{T \in \Theta} (IT(T : IT))^{*T} = \bigcap_{T \in \Theta} T = R.$$

Therefore, $(I_T) = \Gamma(I) \in \Gamma(\text{Inv}^*(R))$, and thus Γ is an isomorphism. \square

The set of nonzero principal fractional ideals forms a subgroup of $\text{Inv}^*(R)$, denoted by $\text{Prin}(I)$. The quotient between $\text{Inv}^*(R)$ and $\text{Prin}(R)$ is called the $*$ -class group of R [3], and it is denoted by $\text{Cl}^*(R)$. If $*_1 \leq *_2$, there is an injective homomorphism $\text{Cl}^{*1}(R) \subseteq \text{Cl}^{*2}(R)$. Of particular interest are the class group of the identity star operation (usually called the *Picard group* of R , denoted by $\text{Pic}(R)$) and the t -class group, which is linked to the factorization properties of the group (see for example [30, 8, 34]). The quotient between $\text{Cl}^*(R)$ and $\text{Pic}(R)$ is called the $*$ -local class group of R , and it is indicated by $G_*(R)$ [3].

Theorem 7.2. *Let R be an integral domain and let Θ be a Jaffard family on R . Then, the map*

$$\begin{aligned} \Lambda: G_*(R) &\longrightarrow \bigoplus_{T \in \Theta} G_{*T}(T) \\ [I] &\longmapsto ([IT])_{T \in \Theta} \end{aligned}$$

is well-defined and a group isomorphism.

Proof. By Proposition 7.1, there are two isomorphisms $\Gamma^* : \text{Inv}^*(R) \longrightarrow \bigoplus_{T \in \Theta} \text{Inv}^{*T}(T)$ and $\Gamma^d : \text{Inv}^d(R) \longrightarrow \bigoplus_{T \in \Theta} \text{Inv}^{dT}(T)$.

Consider the chain of maps

$$\text{Inv}^*(R) \xrightarrow{\Gamma^*} \bigoplus_{T \in \text{Max}(T)} \text{Inv}^{*T}(T) \xrightarrow{\pi} \bigoplus_{T \in \text{Max}(T)} \frac{\text{Inv}^{*T}(T)}{\text{Inv}^{dT}(T)}$$

where π is the componentwise quotient; then, the kernel of π is exactly $\bigoplus_{T \in \Theta} \text{Inv}^{dT}(T)$. However, Γ^* and Γ^d coincide on $\text{Inv}^d(R) \subseteq \text{Inv}^*(R)$; hence,

$$\ker(\pi \circ \Gamma^*) = (\Gamma^d)^{-1}(\ker \pi) = \text{Inv}^d(R).$$

Therefore, there is an isomorphism $\frac{\text{Inv}^*(R)}{\text{Inv}^d(R)} \simeq \bigoplus_{T \in \text{Max}(T)} \frac{\text{Inv}^{*T}(T)}{\text{Inv}^{dT}(T)}$. However, for an arbitrary domain A and an arbitrary $\sharp \in \text{Star}(A)$, we have $\text{Prin}(A) \subseteq \text{Inv}^d(A) \subseteq \text{Inv}^\sharp(A)$, and thus

$$\frac{\text{Inv}^\sharp(A)}{\text{Inv}^d(A)} \simeq \frac{\text{Inv}^\sharp(A)/\text{Prin}(A)}{\text{Inv}^d(A)/\text{Prin}(A)} \simeq \frac{\text{Cl}^\sharp(A)}{\text{Pic}(A)} = G_\sharp(A)$$

so that Λ becomes an isomorphism between $G_*(R)$ and $\bigoplus_{T \in \Theta} G_{*T}(T)$, as claimed. \square

7.1. The class group of a Prüfer domain. If $*$ is a (semi)star operation, we can define the $*$ -class group by mirroring the definition of the case of star operations: we say that I is $*$ -invertible if $(I(R : I))^* = R$, and we define $\text{Cl}^*(R)$ as the quotient between the group of the $*$ -invertible $*$ -ideals (endowed with the $*$ -product) and the subgroup of principal ideals. Since $(R : I) = (0)$ if $I \in \mathbf{F}(R) \setminus \mathcal{F}(R)$, every $*$ -invertible ideal is a fractional ideal, and thus $\text{Cl}^*(R)$ coincides with $\text{Cl}^{*'}(R)$, where $*' := *|_{\mathcal{F}(R)}$ is the restriction of $*$.

The first result of this section is that Proposition 6.5 can be extended to the class group.

Proposition 7.3. *Let R be a Prüfer domain and let P be a nonzero prime ideal of R contained in every maximal ideal. Suppose also that $P \notin \text{Max}(R)$. Let $*$ \in $\text{Star}(R)$ and let \sharp be the corresponding (semi)star operation on $D := R/P$. Then, $\text{Cl}^*(R)$ is naturally isomorphic to $\text{Cl}^\sharp(D)$.*

Proof. Let $\pi : R_P \rightarrow F = Q(D)$ be the quotient map, and let I be a fractional ideal of R contained between P and R_P . We claim that $\pi((R : I)) = (D : \pi(I))$. In fact, if $y \in \pi((R : I))$ then $y = \pi(x)$ for some $x \in (R : I)$, and thus $y\pi(I) = \pi(x)\pi(I) = \pi(xI) \subseteq \pi(R) = D$, and thus $x \in (D : \pi(I))$. Conversely, if $y \in (D : \pi(I))$ and $y = \pi(x)$ then $y\pi(I) \subseteq D$, i.e., $\pi(xI) \subseteq D$. By the correspondence between R -submodules of R_P and D -submodules of F we have $xI \subseteq R$ and $y \in \pi((R : I))$.

Let $J = \pi(I)$ be a \sharp -invertible ideal of D . Then, $(J(D : J))^\sharp = D$, and thus

$$\begin{aligned} R &= \pi^{-1}((J(D : J))^\sharp) = \pi^{-1}(J(D : J))^* = \\ &= (\pi^{-1}(J)\pi^{-1}(D : J))^* = (I(R : I))^*. \end{aligned}$$

Therefore, I is $*$ -invertible, and there is an injective map $\theta : \text{Inv}^\sharp(D) \longrightarrow \text{Inv}^*(R)$. It is also straightforward to see that θ is a group homomorphism.

The well-definedness of the map $* \mapsto *_\phi$ implies that, if J, J' are D -submodules of F , and $I := \pi^{-1}(J)$, $I' := \pi^{-1}(J')$, then $J = zJ'$ for some $z \in F$ if and only if $I = wI'$ for some $w \in K$. Therefore, θ induces an injective map $\bar{\theta} : \text{Cl}^\sharp(D) \longrightarrow \text{Cl}^*(R)$, that is clearly is a group homomorphism.

Let now I be a $*$ -invertible ideal of R . Then, I is v -invertible, and thus $(I : I) = R$ [16, Proposition 34.2(2)]. In particular, I is not a R_P -module, and thus the set $\mathbf{v}(I)$ has an infimum α , where \mathbf{v} is the valuation associated to R_P . If a is an element of valuation α , then $P \subsetneq a^{-1}I \subsetneq R_P$; hence, $a^{-1}I = \phi^{-1}(\phi(a^{-1}I))$ and $[I] = \bar{\theta}([\pi(a^{-1}I)])$, and in particular $[I]$ is in the image of $\bar{\theta}$. Since I was arbitrary, $\bar{\theta}$ is surjective and $\text{Cl}^\sharp(D) \simeq \text{Cl}^*(R)$. \square

Theorem 7.4. *Let R be a Prüfer domain, and suppose that R is either:*

- (a) *semilocal; or*
- (b) *locally finite and finite-dimensional.*

Consider a star operation $$ on R . Then,*

$$G_*(R) \simeq \bigoplus_{\substack{M \in \text{Max}(R) \\ M \neq M^*}} \text{Cl}^v(R_M).$$

Proof. We start by considering the case of R semilocal, and we proceed by induction on the number n of maximal ideals of R . Note that, in this case, $\text{Pic}(R) = (0)$ and so $G_*(R) = \text{Cl}^*(R)$. If $n = 1$, the conclusion is trivial, since $* \neq v$ if and only if $M \neq M^*$.

Suppose $n > 1$ and let Θ be the standard decomposition of R (which is a Jaffard family by Proposition 6.2). By Theorem 7.2, and using the fact that $\text{Pic}(R) = (0) = \text{Pic}(T)$ for every $T \in \Theta$, we have $\text{Cl}^*(R) \simeq \bigoplus_{T \in \Theta} \text{Cl}^{*T}(T)$. Moreover, since a maximal ideal M of R is $*$ -closed if and only if MT is $*_T$ -closed, by induction it suffices to prove the theorem when the standard decomposition of R is $\{R\}$.

In this case, $\text{Jac}(R)$ contains nonzero primes, and by Lemma 6.4 we can find a prime ideal $Q \subseteq \text{Jac}(R)$ such that $\text{Jac}(R/Q)$ does not contain nonzero prime ideals. Let $A := R/Q$.

The standard decomposition Θ' of A is nontrivial, and thus every $B \in \Theta'$ is a semilocal Prüfer domain with less than n maximal ideals. Moreover, by Proposition 7.3, $\text{Cl}^*(R) \simeq \text{Cl}^\sharp(A)$, where \sharp is the restriction to $\mathcal{F}(A)$ of the (semi)star operation corresponding to $*$. Therefore, by the inductive hypothesis,

$$\text{Cl}^\sharp(A) \simeq \bigoplus_{B \in \Theta'} \text{Cl}^v(B) \simeq \bigoplus_{\substack{B \in \Theta' \\ N \in \text{Max}(B) \\ N \neq N^\sharp B}} \bigoplus \text{Cl}^v(B_N) \simeq \bigoplus_{\substack{N \in \text{Max}(A) \\ N \neq N^\sharp}} \text{Cl}^v(A_N).$$

Thus,

$$\mathrm{Cl}^*(R) \simeq \mathrm{Cl}^\sharp(A) \simeq \bigoplus_{\substack{N \in \mathrm{Max}(A) \\ N \neq N^\sharp}} \mathrm{Cl}^v(A_N).$$

However, if M is the maximal ideal of R which corresponds to the maximal ideal N of A , then $R_M/QR_M \simeq A_N$, and thus by [4, Theorem 3.5] we have $\mathrm{Cl}^v(R_M) \simeq \mathrm{Cl}^v(A_N)$; the claim follows.

Suppose now R is finite-dimensional and of finite character, and let Θ be the standard decomposition of R . By Lemma 6.1, there is a bijective correspondence between Θ and the height 1 prime ideals of R , and every $T \in \Theta$ is semilocal. Hence, by Proposition 6.2 and by the previous case,

$$G_*(R) \simeq \bigoplus_{T \in \Theta} G_{*T}(T) \simeq \bigoplus_{T \in \Theta} \mathrm{Cl}^{*T}(T) \simeq \bigoplus_{T \in \Theta} \bigoplus_{\substack{M \in \mathrm{Max}(T) \\ M \neq M^{*T}}} \mathrm{Cl}^v(T_M).$$

The conclusion now follows since $T_M = R_N$ (where $N := M \cap R$) and $N = N^*$ if and only if $M = M^{*T}$. \square

Corollary 7.5. *Let R be a Bézout domain, and suppose that R is either:*

- (a) *semilocal; or*
- (b) *finite-dimensional and of finite character.*

Let $$ be a star operation on R . Then,*

$$\mathrm{Cl}^*(R) \simeq \bigoplus_{\substack{M \in \mathrm{Max}(R) \\ M \neq M^*}} \mathrm{Cl}^v(R_M).$$

Proof. It is enough to note that $\mathrm{Pic}(R) = 0$ if R is a Bézout domain, so that $G_*(R) = \mathrm{Cl}^*(R)$ for every $*$ \in $\mathrm{Star}(R)$, and then apply Theorem 7.4. \square

Corollary 7.6. *Let R be a Bézout domain, and suppose that R is either*

- (a) *semilocal; or*
- (b) *finite-dimensional and of finite character.*

Let S be a multiplicatively closed subset of R . Then, there is a natural surjective group homomorphism $\mathrm{Cl}^v(R) \rightarrow \mathrm{Cl}^v(S^{-1}R)$, $[I] \mapsto [S^{-1}I]$.

Proof. Let $\Delta := \{M \in \mathrm{Max}(R) : M \cap S = \emptyset\}$. Then, for every $M \in \Delta$, $R_M = (S^{-1}R)_{S^{-1}M}$, and thus the isomorphism of Theorem 7.4 reduces to a surjective map $\mathrm{Cl}^v(R) \rightarrow \bigoplus_{M \in \Delta} \mathrm{Cl}^v(R_M) \simeq \mathrm{Cl}^v(S^{-1}R)$, where the last equality comes from the fact that the maximal ideals of $S^{-1}R$ are the extensions of the ideals belonging to Δ . \square

Therefore, under each case of Theorem 7.4, the determination of $G_*(R)$ is reduced to the calculation of $\mathrm{Cl}^v(V)$, where V is a valuation domain. In the case where the maximal ideal M of V is *branched* (that is, if there is a M -primary ideal of V different from R , or equivalently

if there is a prime ideal $P \subsetneq M$ such that there are no prime ideal properly contained between P and M [16, Theorem 17.3]), this group has been calculated in [4, Corollaries 3.6 and 3.7]. Indeed, if P is the prime ideal directly below M , and H is the value group of V/P (represented as a subgroup of \mathbb{R}), then

$$\mathrm{Cl}^v(V) \simeq \begin{cases} 0 & \text{if } G \simeq \mathbb{Z} \\ \mathbb{R}/H & \text{otherwise.} \end{cases}$$

In particular, we have the following.

Corollary 7.7. *Let R be a Bézout domain, and suppose that R is either:*

- (a) *semilocal; or*
- (b) *finite-dimensional and of finite character.*

For every $ \in \mathrm{Star}(R)$, $\mathrm{Cl}^*(R)$ is an injective group (equivalently, an injective \mathbb{Z} -module).*

Proof. By Corollary 7.5 and the previous discussion, $\mathrm{Cl}^*(R) \simeq \bigoplus \mathbb{R}/H_\alpha$, for a family $\{H_\alpha : \alpha \in A\}$ of additive subgroups of \mathbb{R} . Each \mathbb{R}/H_α is a divisible group, and thus so is their direct sum; however, a divisible group is injective, and thus so is $\mathrm{Cl}^*(R)$. \square

We end with a result similar in spirit to Proposition 6.7.

Proposition 7.8. *Let R be a Prüfer domain and suppose that R is either:*

- (a) *semilocal; or*
- (b) *finite-dimensional and of finite character.*

Let $ \in \mathrm{Star}(R)$. If $I, J \in \mathrm{Inv}^*(R)$, then $I + J \in \mathrm{Inv}^*(R)$.*

Proof. Suppose first that R is semilocal, and proceed by induction on $n := |\mathrm{Max}(R)|$. If $n = 1$, then R is a valuation domain and $I + J$ is equal either to I or to J , and the claim is proved.

Suppose the claim is true up to rings with $n - 1$ maximal ideals, let $|\mathrm{Max}(R)| = n$ and consider the standard decomposition Θ of R . By Proposition 7.1, $I + J \in \mathrm{Inv}^*(R)$ if and only if $(I + J)T \in \mathrm{Inv}^{*T}(T)$ for every $T \in \Theta$; therefore, if Θ is not trivial, we can use the inductive hypothesis. Suppose Θ is trivial: then, $\mathrm{Jac}(R)$ contains nonzero prime ideals, and by Lemma 6.4 there is a nonzero prime ideal $Q \subseteq \mathrm{Jac}(R)$ such that $\mathrm{Jac}(R/Q)$ does not contain nonzero primes. By Proposition 7.3, I/Q and J/Q are \sharp -invertible \sharp -ideals of R/Q (where \sharp is the (semi)star operation induced by $*$), and in particular I/Q and J/Q are fractional ideals of R/Q .

By construction, R/Q admits a nontrivial Jaffard family Λ : for every $U \in \Lambda$, $(I/Q)U$ and $(J/Q)U$ are \sharp_U -invertible \sharp_U -ideals, and thus by the inductive hypothesis so is $(I/Q)U + (J/Q)U = ((I + J)/Q)U$. Hence

$(I + J)/Q$ is a \sharp -invertible \sharp -ideal, and so $I + J$ is a $*$ -invertible $*$ -ideal, i.e., $I + J \in \text{Inv}^*(R)$.

If now R is locally finite and finite-dimensional, we see that if Θ is the standard decomposition of R then every $T \in \Theta$ is semilocal. The ideal $I + J$ is $*$ -invertible if and only if $(I + J)T$ is $*_T$ -invertible for every $T \in \Theta$; however, since IT and JT are $*_T$ -invertible $*_T$ -ideals, the previous part of the proof shows that so is $IT + JT = (I + J)T$. Therefore, $I + J \in \text{Inv}^*(R)$. \square

8. ACKNOWLEDGEMENTS

The author would like to thank the referee for his/her careful reading of the manuscript and his/her suggestions.

REFERENCES

- [1] D. D. Anderson. Star-operations induced by overrings. *Comm. Algebra*, 16(12):2535–2553, 1988.
- [2] D. D. Anderson and David F. Anderson. Examples of star operations on integral domains. *Comm. Algebra*, 18(5):1621–1643, 1990.
- [3] David F. Anderson. A general theory of class groups. *Comm. Algebra*, 16(4):805–847, 1988.
- [4] David F. Anderson, Marco Fontana, and Muhammad Zafrullah. Some remarks on Prüfer $*$ -multiplication domains and class groups. *J. Algebra*, 319(1):272–295, 2008.
- [5] M. F. Atiyah and I. G. Macdonald. *Introduction to commutative algebra*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969.
- [6] Silvana Bazzoni and Luigi Salce. Warfield domains. *J. Algebra*, 185(3):836–868, 1996.
- [7] Nicolas Bourbaki. *Commutative algebra. Chapters 1–7*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 1989. Translated from the French, Reprint of the 1972 edition.
- [8] Alain Bouvier and Muhammad Zafrullah. On some class groups of an integral domain. *Bull. Soc. Math. Grèce (N.S.)*, 29:45–59, 1988.
- [9] Carmelo A. Finocchiaro, Marco Fontana, and Dario Spirito. New distinguished classes of spectral spaces: a survey. In *Multiplicative Ideal Theory and Factorization Theory: Commutative and Non-Commutative Perspectives*. Springer Verlag, 2016.
- [10] Carmelo A. Finocchiaro, Marco Fontana, and Dario Spirito. Spectral spaces of semistar operations. *J. Pure Appl. Algebra*, 220(8):2897–2913, 2016.
- [11] Carmelo A. Finocchiaro and Dario Spirito. Some topological considerations on semistar operations. *J. Algebra*, 409:199–218, 2014.
- [12] Carmelo A. Finocchiaro and Dario Spirito. Topology, intersections and flat overrings. *Proc. Amer. Math. Soc.*, 144(10):4125–4133, 2016.
- [13] Marco Fontana, Evan Houston, and Thomas Lucas. *Factoring ideals in integral domains*, volume 14 of *Lecture Notes of the Unione Matematica Italiana*. Springer, Heidelberg; UMI, Bologna, 2013.
- [14] Marco Fontana and James A. Huckaba. Localizing systems and semistar operations. In *Non-Noetherian commutative ring theory*, volume 520 of *Math. Appl.*, pages 169–197. Kluwer Acad. Publ., Dordrecht, 2000.

- [15] Marco Fontana, James A. Huckaba, and Ira J. Papick. *Prüfer domains*, volume 203 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker Inc., New York, 1997.
- [16] Robert Gilmer. *Multiplicative ideal theory*. Marcel Dekker Inc., New York, 1972. Pure and Applied Mathematics, No. 12.
- [17] Malcolm Griffin. Some results on v -multiplication rings. *Canad. J. Math.*, 19:710–722, 1967.
- [18] Franz Halter-Koch. *Ideal systems. An introduction to multiplicative ideal theory*, volume 211 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker, Inc., New York, 1998.
- [19] William Heinzer. Integral domains in which each non-zero ideal is divisorial. *Mathematika*, 15:164–170, 1968.
- [20] William J. Heinzer, James A. Huckaba, and Ira J. Papick. m -canonical ideals in integral domains. *Comm. Algebra*, 26(9):3021–3043, 1998.
- [21] Evan G. Houston, Abdeslam Mimouni, and Mi Hee Park. Integral domains which admit at most two star operations. *Comm. Algebra*, 39(5):1907–1921, 2011.
- [22] Evan G. Houston, Abdeslam Mimouni, and Mi Hee Park. Noetherian domains which admit only finitely many star operations. *J. Algebra*, 366:78–93, 2012.
- [23] Evan G. Houston, Abdeslam Mimouni, and Mi Hee Park. Integrally closed domains with only finitely many star operations. *Comm. Algebra*, 42(12):5264–5286, 2014.
- [24] Evan G. Houston and Mi Hee Park. A characterization of local noetherian domains which admit only finitely many star operations: The infinite residue field case. *J. Algebra*, 407:105–134, 2014.
- [25] Paul Jaffard. *Les systèmes d'idéaux*. Travaux et Recherches Mathématiques, IV. Dunod, Paris, 1960.
- [26] Wolfgang Krull. *Idealtheorie*. Springer-Verlag, Berlin, 1935.
- [27] Hideyuki Matsumura. *Commutative ring theory*, volume 8 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1986. Translated from the Japanese by M. Reid.
- [28] Bruce Olberding. Globalizing local properties of Prüfer domains. *J. Algebra*, 205(2):480–504, 1998.
- [29] Fred Richman. Generalized quotient rings. *Proc. Amer. Math. Soc.*, 16:794–799, 1965.
- [30] Pierre Samuel. Sur les anneaux factoriels. *Bull. Soc. Math. France*, 89:155–173, 1961.
- [31] Dario Spirito. Towards a classification of stable semistar operations on a Prüfer domain. *submitted*.
- [32] Janet C. Vassilev. Structure on the set of closure operations of a commutative ring. *J. Algebra*, 321(10):2737–2753, 2009.
- [33] Muhammad Zafrullah. Ascending chain conditions and star operations. *Comm. Algebra*, 17(6):1523–1533, 1989.
- [34] Muhammad Zafrullah. Putting t -invertibility to use. In *Non-Noetherian commutative ring theory*, volume 520 of *Math. Appl.*, pages 429–457. Kluwer Acad. Publ., Dordrecht, 2000.

E-mail address: spirito@mat.uniroma3.it

DIPARTIMENTO DI MATEMATICA E FISICA, UNIVERSITÀ DEGLI STUDI “ROMA TRE”, ROMA, ITALY