

The Picard group of the universal moduli stack of principal bundles on pointed smooth curves II

ROBERTO FRINGUELLI AND FILIPPO VIVIANI

Abstract. In this paper, which is a sequel of [14], we investigate, for any reductive group G over an algebraically closed field k , the Picard group of the universal moduli stack $\text{Bun}_{G,g,n}$ of G -bundles over n -pointed smooth projective curves of genus g . In particular, we give new functorial presentations of the Picard group of $\text{Bun}_{G,g,n}$, we study the restriction homomorphism onto the Picard group of the moduli stack of principal G -bundles over a fixed smooth curve, we determine the Picard group of the rigidification of $\text{Bun}_{G,g,n}$ by the center of G as well as the image of the obstruction homomorphism of the associated gerbe. As a consequence, we compute the divisor class group of the moduli space of semistable G -bundles over n -pointed smooth projective curves of genus g .

Mathematics Subject Classification (2020): 14H60 (primary); 14D20, 14C22, 20G07, 14H10 (secondary).

Contents

1	Introduction	368
2	Preliminaries	375
3	The universal moduli stack $\text{Bun}_{G,g,n}$ and its Picard group	387
4	The restriction homomorphism	404
5	The rigidification $\text{Bun}_{G,g,n}^{\delta} // \mathcal{Z}(G)$ and its Picard group	412
6	The universal moduli space $M_{G,g,n}^{ss}$ and its divisor class group	423
7	Examples	427
	References	446

The first author acknowledges PRIN2017 CUP E84I19000500006, and the MIUR Excellence Department Project awarded to the Department of Mathematics, University of Rome Tor Vergata, CUP E83C18000100006. The second author was funded by the projects PRIN2017 “Advances in Moduli Theory and Birational Classification” and PRIN2020 “Curves, Ricci Flat Varieties and their Interactions” funded by MIUR (Italy), and by the projects EXPL/MAT-PUR/1162/2021 funded by FCT (Portugal), and he is a member of the Center of Mathematics of the University of Coimbra – UID/MAT/00324/2019 funded by FCT (Portugal).

Received September 22, 2020; accepted in revised form November 9, 2021.

Published online March 2023.

1. Introduction

The aim of this paper, which is a sequel of the paper [14], is to study the Picard group of the *universal moduli stack of (principal) G -bundles* $\text{Bun}_{G,g,n}$, which parametrizes G -bundles, where G is a connected and smooth linear algebraic group over $k = \bar{k}$, over families of (connected, smooth and projective) k -curves of genus $g \geq 0$ endowed with $n \geq 0$ pairwise disjoint ordered sections. We refer the reader to [14] for the motivation behind this investigation as well as for its relationship with previous results in the literature.

Recall (see Theorem 3.1) that the stack $\text{Bun}_{G,g,n}$ is an algebraic stack, locally of finite type and smooth over the moduli stack $\mathcal{M}_{g,n}$ of n -marked curves of genus g and its connected components (which are integral and smooth over k) are in functorial bijection with the fundamental group $\pi_1(G)$. We will denote the connected components and the restriction of the forgetful morphism by

$$\Phi_G^\delta : \text{Bun}_{G,g,n}^\delta \rightarrow \mathcal{M}_{g,n} \quad \text{for any } \delta \in \pi_1(G).$$

We proved in [14, Theorem A] that if $\text{red} : G \rightarrow G^{\text{red}}$ is the reductive quotient of G , *i.e.*, the quotient of G by its unipotent radical, then for any $\delta \in \pi_1(G) \xrightarrow[\cong]{\pi_1(\text{red})} \pi_1(G^{\text{red}})$ the pull-back homomorphism

$$\text{red}_\#^* : \text{Pic} \left(\text{Bun}_{G^{\text{red}},g,n}^\delta \right) \xrightarrow{\cong} \text{Pic} \left(\text{Bun}_{G,g,n}^\delta \right)$$

is an isomorphism. Hence, throughout this paper, we will restrict to the case of a *reductive group* G . We fix a maximal torus $\iota : T_G \hookrightarrow G$ and let \mathscr{W}_G be the Weyl group of G .

Since the Picard group of $\mathcal{M}_{g,n}$ is well-known up to torsion (and completely known if $\text{char}(k) \neq 2$ by [13]) and the pull-back morphism

$$\left(\Phi_G^\delta \right)^* : \text{Pic}(\mathcal{M}_{g,n}) \rightarrow \text{Pic} \left(\text{Bun}_{G,g,n}^\delta \right)$$

is injective since Φ_G^δ is fpqc and cohomologically flat in degree zero (see Theorem 3.1 for the definition), we can focus our attention on the relative Picard group

$$\text{RPic} \left(\text{Bun}_{G,g,n}^\delta \right) := \text{Pic} \left(\text{Bun}_{G,g,n}^\delta \right) / \left(\Phi_G^\delta \right)^* \left(\text{Pic}(\mathcal{M}_{g,n}) \right).$$

Throughout this paper, we will mainly restrict to the case of *positive genus*; the case $g = 0$ is easier, see Remarks 4.7 and 5.10.

The relative Picard group $\text{RPic}(\text{Bun}_{G,g,n}^\delta)$ was described in [14, Theorem C]: it is generated by the image of a functorial transgression homomorphism

$$\tau_G^\delta : \text{Sym}^2(\Lambda^*(T_G))^{\mathscr{W}_G} \cong \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{W_G} \hookrightarrow \text{RPic} \left(\text{Bun}_{G,g,n}^\delta \right),$$

where $\text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{W_G}$ is the lattice of W_G -invariant even symmetric bilinear forms on the lattice $\Lambda(T_G)$ of cocharacters of a maximal torus T_G in G , and by the image of the pull-back homomorphism

$$\text{ab}_\#^* : \text{RPic}(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}) \hookrightarrow \text{RPic}(\text{Bun}_{G,g,n}^{\delta}),$$

where $\text{ab} : G \twoheadrightarrow G^{\text{ab}}$ is the maximal Abelian quotient and $\delta^{\text{ab}} := \pi_1(\text{ab})(\delta) \in \pi_1(G^{\text{ab}})$. Moreover, the image of $\text{ab}_\#^*$ coincides with the subgroup generated by the tautological line bundles, see [14, Theorem B].

The first result of this paper is a new description of $\text{RPic}(\text{Bun}_{G,g,n}^{\delta})$ in terms of three functorial exact sequences.

Theorem 1.1 (see Corollary 3.4, Theorem 3.6, Theorem 3.12). *Assume that $g \geq 1$. Let G be a reductive group and fix $\delta \in \pi_1(G)$. Then the relative Picard group of $\text{Bun}_{G,g,n}^{\delta}$ sits into the following functorial commutative diagram with exact rows*

$$\begin{array}{ccccc} \Lambda^*(G^{\text{ab}}) \otimes H_{g,n} & \xrightarrow{j_G^\delta} & \text{RPic}(\text{Bun}_{G,g,n}^{\delta}) & \xrightarrow{\omega_G^\delta \oplus \gamma_G^\delta} & \text{NS}(\text{Bun}_{G,g,n}^{\delta}) \\ \downarrow & & \parallel & & \downarrow \text{res}_G^{\text{NS}} \\ \Lambda^*(G^{\text{ab}}) \otimes \widehat{H}_{g,n} & \xrightarrow{i_G^\delta} & \text{RPic}(\text{Bun}_{G,g,n}^{\delta}) & \xrightarrow{\gamma_G^\delta} & \text{Bil}^{s,\mathcal{D}\text{-ev}}(\Lambda(T_G))^{W_G} \\ \downarrow & & \parallel & & \downarrow \text{res}_{\mathcal{D}} \\ \text{RPic}(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}) & \xrightarrow{\text{ab}_\#^*} & \text{RPic}(\text{Bun}_{G,g,n}^{\delta}) & \xrightarrow{\theta_G^\delta} & \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}}(G)) | \Lambda(T_{G^{\text{ss}}}))^{W_G} \end{array} \quad (1.1)$$

where the left vertical morphisms are injective and the right vertical morphisms are surjective.

Moreover, the image of $\omega_G^\delta \oplus \gamma_G^\delta$ is equal to

$$\begin{aligned} & \text{Im}(\omega_G^\delta \oplus \gamma_G^\delta) \\ &= \begin{cases} \text{NS}(\text{Bun}_{G,g,n}^{\delta}) & \text{if } n \geq 1 \\ \left\{ ([\chi], b) \in \text{NS}(\text{Bun}_{G,g,n}^{\delta}) : \begin{array}{l} [\chi(x) - b(\delta \otimes x)] + (g-1)b(x \otimes x) \\ \text{is divisible by } 2g-2, \text{ for any } x \in \Lambda(T_G) \end{array} \right\} & \text{if } n = 0. \end{cases} \end{aligned}$$

In the above diagram:

- $\Lambda^*(G^{\text{ab}})$ is the lattice of characters of G^{ab} ;
- The groups $\widehat{H}_{g,n}$ and $H_{g,n}$ are defined by

$$\begin{aligned} \widehat{H}_{g,n} &:= \begin{cases} \mathbb{Z} \oplus \mathbb{Z}^n & \text{if } g \geq 2 \\ \mathbb{Z}^n & \text{if } g = 1 \end{cases} \quad \text{and} \\ H_{g,n} &:= \begin{cases} \{(m, \zeta) \in \mathbb{Z} \oplus \mathbb{Z}^n : (2g-2)m + |\zeta| = 0\} & \text{if } g \geq 2 \\ \{\zeta \in \mathbb{Z}^n : |\zeta| = 0\} & \text{if } g = 1; \end{cases} \end{aligned}$$

- $\text{Bil}^{s, \mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathcal{W}_G}$ is the lattice of \mathcal{W}_G -invariant symmetric bilinear forms on $\Lambda(T_G)$ which are even on $\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_{\mathcal{D}(G)})$, where $T_{\mathcal{D}(G)}$ is the maximal torus of the derived subgroup $\mathcal{D}(G)$ of G , see Corollary 2.6;
- $\text{Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}_G}$ is the lattice of \mathcal{W}_G -invariant even symmetric bilinear forms on $\Lambda(T_{\mathcal{D}(G)})$ which are integral on $\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_{G^{\text{ss}}})$, where $T_{G^{\text{ss}}}$ is the maximal torus of the semisimplification G^{ss} of G , see Proposition 2.4;
- $\text{NS}(\text{Bun}_{G, g, n}^\delta) \subset \Lambda^*(T_G) / \Lambda^*(T_{G^{\text{ad}}}) \oplus \text{Bil}^{s, \mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathcal{W}_G}$, where $G^{\text{ad}} = G / \mathcal{Z}(G)$ with $\mathcal{Z}(G)$ center of G , is introduced in Definition 3.9 and further studied in Definition/Lemma 3.10 and Proposition 3.11;
- The homomorphisms $j_G^\delta, \omega_G^\delta, \gamma_G^\delta, i_G^\delta$ and θ_G^δ are defined in, respectively, Theorem 3.12, Definition/Lemma 3.7, Definition/Lemma 3.5, Theorem 3.6 and Corollary 3.4.

The second result of this paper is a description of the restriction homomorphism

$$\text{res}_G^\delta(C) : \text{RPic}(\text{Bun}_{G, g, n}^\delta) \rightarrow \text{Pic}(\text{Bun}_G^\delta(C)) \tag{1.2}$$

for any $(C, p_1, \dots, p_n) \in \mathcal{M}_{g, n}(k)$, where $\text{Bun}_G(C)$ is the k -stack of G -bundles on C , i.e., the stack over k whose S -points $\text{Bun}_G(C)(S)$ is the groupoid of G -bundles on $C_S := C \times_k S$ for any k -scheme S , and it is canonically identified with the fiber of $\Phi_{G, g, n}^\delta$ over $(C, p_1, \dots, p_n) \in \mathcal{M}_{g, n}(k)$. The Picard group $\text{Pic}(\text{Bun}_G^\delta(C))$ has been described by Biswas-Hoffman in [3], see Theorem 4.1 and Proposition 4.2.

Theorem 1.2 (see Theorem 4.3, Theorem 4.5, Corollary 4.6). *Assume that $g \geq 1$ and let $(C, p_1, \dots, p_n) \in \mathcal{M}_{g, n}(k)$ be a geometric point. Let G be a reductive group and fix $\delta \in \pi_1(G)$.*

- (1) *The restriction homomorphism (1.2) sits into the following functorial commutative diagram with exact rows:*

$$\begin{CD} \Lambda^*(G^{\text{ab}}) \otimes H_{g, n} @<j_G^\delta<< \text{RPic}(\text{Bun}_{G, g, n}^\delta) @>\omega_G^\delta \oplus \gamma_G^\delta>> \text{NS}(\text{Bun}_{G, g, n}^\delta) \\ @V \text{res}_G^\delta(C)^\circ VV @VV \text{res}_G^\delta(C) V @VV \text{res}_G^\delta(C)^{\text{NS}} V \\ \text{Hom}(\pi_1(G), J_C(k)) @<j_G^\delta(C)<< \text{Pic}(\text{Bun}_G^\delta(C)) @>c_G^\delta(C)>> \text{NS}(\text{Bun}_G^\delta(C)); \end{CD} \tag{1.3}$$

- (2) *The composition $\overline{\text{res}_G^\delta(C)} := c_G^\delta(C) \circ \text{res}_G^\delta(C)$ sits into the following functorial commutative diagram with exact rows:*

$$\begin{CD} \text{RPic}(\text{Bun}_{G^{\text{ab}}, g, n}^{\delta^{\text{ab}}}) @<\text{ab}_\#^* << \text{RPic}(\text{Bun}_{G, g, n}^\delta) @>\theta_G^\delta >> \text{Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}_G} \\ @V \overline{\text{res}_G^{\delta^{\text{ab}}}} VV @VV \overline{\text{res}_G^\delta} V @VV r_G V \\ \text{NS}(\text{Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C)) @<\text{ab}^*. \text{NS}(C)<< \text{NS}(\text{Bun}_G^\delta(C)) @>P>> \text{Bil}^{s, \text{-ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}_G}. \end{CD} \tag{1.4}$$

In particular, if $\text{id}_{J_C} : \mathbb{Z} \xrightarrow{\cong} \text{End}(J_C)$ is an isomorphism (which is true if k is uncountable and $(C, p_1, \dots, p_n) \in \mathcal{M}_{g,n}(k)$ is very general), then $\text{coker}(\text{res}_G^\delta(C))$ sits into a canonical short exact sequence

$$0 \rightarrow \text{coker} \left(\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} \oplus \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}} \right) \rightarrow \text{coker} \left(\overline{\text{res}_G^\delta(C)} \right) \rightarrow \text{coker}(r_G) \rightarrow 0,$$

where

$$\text{coker} \left(\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} \oplus \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}} \right) = \begin{cases} 0 & \text{if } n > 0 \\ \left(\frac{\mathbb{Z}}{(2g-2)\mathbb{Z}} \right)^{\dim G^{\text{ab}}} & \text{if } n = 0. \end{cases}$$

The group $\text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}_G}$ has been defined after Theorem 1.1 while

- $\text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}_G}$ is the lattice of \mathcal{W}_G -invariant symmetric bilinear forms on $\Lambda(T_{\mathcal{D}(G)})$ which are integral on $\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_{G^{\text{ss}}})$ and even on $\Lambda(T_G) \otimes \Lambda(T_G)$, where T_G is the maximal torus of the simply-connected cover G of the semisimplification G of G , see Definition-Lemma 2.11.

It follows that the cokernel of r_G is a 2-elementary Abelian group of rank bounded by the number of the simple factors of the semisimple part \mathfrak{g}^{ss} of the Lie algebra \mathfrak{g} of G (see Definition/Lemma 2.11 and Corollary 2.7). In Section 7, we compute $\text{coker}(r_G)$ for all reductive groups G such that \mathfrak{g}^{ss} is a simple Lie algebra.

The third result of this paper is the computation of the relative Picard group of the rigidification

$$\nu_G^\delta : \text{Bun}_{G,g,n}^\delta \rightarrow \text{Bun}_{G,g,n}^\delta // \mathcal{Z}(G) := \mathfrak{Bun}_{G,g,n}^\delta,$$

of the stack $\text{Bun}_{G,g,n}^\delta$ by the center $\mathcal{Z}(G)$ of G , which acts functorially on any G -bundle. This is also closely related to the divisor class group of the adequate moduli space $M_{G,g,n}^{\delta,ss}$ of the locus $\text{Bun}_{G,g,n}^{\delta,ss} \subseteq \text{Bun}_{G,g,n}^\delta$ parametrizing semistable G -bundles over n -marked curves of genus g , or equivalently of its image $\mathfrak{Bun}_{G,g,n}^{\delta,ss}$ in the $\mathcal{Z}(G)$ -rigidification $\mathfrak{Bun}_{G,g,n}^\delta$. See Section 6 for a discussion of the properties of the loci $\text{Bun}_{G,g,n}^{\delta,ss}$ and $\mathfrak{Bun}_{G,g,n}^{\delta,ss}$, as well as for the state of the art on the existence of the adequate moduli space $M_{G,g,n}^{\delta,ss}$.

Theorem 1.3 (see Theorems 5.5 and 6.6). *Assume that $g \geq 1$. Let G be a reductive group and fix $\delta \in \pi_1(G)$.*

- (1) *The relative Picard group of the rigidification $\mathfrak{Bun}_{G,g,n}^\delta := \text{Bun}_{G,g,n}^\delta // \mathcal{Z}(G)$ sits into the exact sequence*

$$0 \rightarrow \Lambda^*(G^{\text{ab}}) \otimes H_{g,n} \xrightarrow{\overline{j}_G^\delta} \text{RPic} \left(\text{Bun}_{G,g,n}^\delta \right) \xrightarrow{\overline{\gamma}_G^\delta} \text{NS} \left(\mathfrak{Bun}_{G,g,n}^\delta \right), \quad (1.5)$$

and the image of $\overline{\gamma_G^\delta}$ is equal to

$$\text{Im}(\overline{\gamma_G^\delta}) = \begin{cases} \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) & \text{if } n \geq 1 \\ \left\{ b \in \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) : \begin{array}{l} 2g - 2|b(\delta \otimes x) + (g - 1)b(x \otimes x) \\ \text{for any } x \in \Lambda(T_G) \end{array} \right\} & \text{if } n = 0; \end{cases} \quad (1.6)$$

(2) Assume that there exists an adequate moduli space $\pi : \mathfrak{Bun}_{G,g,n}^{\delta,ss} \rightarrow M_{G,g,n}^{\delta,ss}$ (e.g., $\text{char}(k) = 0$ and $n = 0$ or $n > 2g + 2$). Suppose that $g + n \geq 3$ (i.e., $M_{g,n}$ is generically a variety) and that one of the following conditions hold:

- (i) G is a torus;
- (ii) G is not a torus, $\text{char}(k) > 0$, $g \geq 4$;
- (iii) G is not a torus, $\text{char}(k) = 0$, $g \geq 2$, with the exception of the case $g = 2$ and G having a non-trivial homomorphism into PGL_2 .

Then there are isomorphisms

$$\text{Cl}(M_{G,g,n}^{\delta,ss}) \xrightarrow{\cong} \text{Pic}(\mathfrak{Bun}_{G,g,n}^{\delta,ss}) \xrightarrow[\text{res}^{-1}]{\cong} \text{Pic}(\mathfrak{Bun}_{G,g,n}^\delta),$$

where res is the restriction homomorphism and the first isomorphism is obtained by pull-back along π .

The group $\text{NS}(\mathfrak{Bun}_{G,g,n}^\delta)$ is introduced in Definition 5.3 and further studied in Proposition 5.4. From the above Theorem 1.3, one easily recovers [24, Theorem B(i) and 1.5] (see also [18]) if $G = \mathbb{G}_m$, $n = 0$, $g \geq 2$ and $\text{char}(k) = 0$; [11, Theorem B(i) and Theorem A.2] (see also [19]) if $G = GL_r$, $n = 0$, $g \geq 2$ and $\text{char}(k) = 0$.

The final result of the paper deals with the triviality of the $\mathcal{L}(G)$ -gerbe v_G^δ . From the Leray spectral sequence associated to the $\mathcal{L}(G)$ -gerbe v_G^δ and the multiplicative group \mathbb{G}_m , we get the exact sequence

$$\begin{aligned} \text{Pic}(\mathfrak{Bun}_{G,g,n}^\delta) &\xrightarrow{(v_G^\delta)^*} \text{Pic}(\text{Bun}_{G,g,n}^\delta) \xrightarrow{\text{wt}_G^\delta} \Lambda^*(\mathcal{L}(G)) \\ &\xrightarrow{\text{obs}_G^\delta} H^2(\mathfrak{Bun}_{G,g,n}^\delta, \mathbb{G}_m) \xrightarrow{(v_G^\delta)^*} H^2(\text{Bun}_{G,g,n}^\delta, \mathbb{G}_m). \end{aligned} \quad (1.7)$$

For a geometric interpretation of the weight homomorphism wt_G^δ and of the obstruction homomorphism obs_G^δ , see Section 5. In particular, $\text{Im}(\text{obs}_G^\delta) \cong \text{coker}(\text{wt}_G^\delta)$ is an obstruction to the triviality of the $\mathcal{L}(G)$ -gerbe v_G^δ .

Theorem 1.4 (see **Theorem 5.7**). *Assume that $g \geq 1$. Let G be a reductive group and fix $\delta \in \pi_1(G)$.*

(1) *If $n > 0$ then*

$$\text{coker} \left(\text{wt}_G^\delta \right) \xrightarrow{\cong} \text{coker} \left(\text{ev}_{\mathcal{D}(G)}^\delta \right);$$

(2) *If $n = 0$ then the cokernel of wt_G^δ sits in an exact sequence*

$$\begin{aligned} 0 \rightarrow \text{coker} \left(\overline{\gamma}_G^\delta \right) \xrightarrow{\partial_G^\delta} \text{Hom} \left(\Lambda(G^{\text{ab}}), \frac{\mathbb{Z}}{(2g-2)\mathbb{Z}} \right) \xrightarrow{\widetilde{\Lambda}_{\text{ab}}^*} \text{coker}(\text{wt}_G^\delta) \\ \xrightarrow{\widetilde{\Lambda}_{\mathcal{D}}^*} \text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta) \rightarrow 0. \end{aligned}$$

For a definition of the homomorphism $\text{ev}_{\mathcal{D}(G)}^\delta$, see Subsection 2.2. In Section 7, we compute $\text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta)$ for all reductive groups G such that \mathfrak{g}^{ss} is a simple Lie algebra, together with its quotient $\text{coker}(\widetilde{\text{ev}}_{\mathcal{D}(G)}^\delta)$ (see Definition/Lemma 2.11(ii), which is an obstruction to the triviality of the $\mathcal{L}(G)$ -gerbe

$$v_G^\delta(C) : \text{Bun}_G^\delta(C) \rightarrow \mathfrak{Bun}_G^\delta(C) := \text{Bun}_G^\delta(C) // \mathcal{L}(G),$$

for any $(C, p_1, \dots, p_n) \in \mathcal{M}_{g,n}(k)$, as shown by Biswas-Hoffmann [5], see Theorem 5.1.

From Theorem 1.4, one easily recovers [24, Theorem 6.4] if $G = \mathbb{G}_m, n = 0, g \geq 3$ and $\text{char}(k) = 0$; [11, Corollary 3.3.2(i)] if $G = \text{GL}_r, n = 0, g \geq 3$ and $\text{char}(k) = 0$; [10, Theorem B(i)] if $G = \text{GL}_r, g \geq 3$ and $\text{char}(k) = 0$.

The computation of the image of the obstruction homomorphism obs_G^δ carried out in Theorem 1.4 will be a crucial ingredient in our upcoming work [12], where we will compute the (cohomological) Brauer groups of $\text{Bun}_{G,g,n}^\delta, \mathfrak{Bun}_{G,g,n}^\delta$ and $M_{G,g,n}^{\delta,ss}$, extending the work of Pirisi and the first author [10] from $G = \text{GL}_r$ to an arbitrary reductive group G . This is also closely related to the works of Biswas-Hogadi [4] and Biswas-Holla [6], where the (cohomological) Brauer group of $\text{Bun}_G(C)$ (and of its good moduli space) has been computed for a fixed curve C and a complex semisimple group G .

Notation 1.5. We denote by $k = \bar{k}$ an algebraically closed field of arbitrary characteristic. All the schemes and algebraic stacks that we will appear in this paper will be locally of finite type over k (hence locally Noetherian).

Notation 1.6. A *curve* is a connected, smooth and projective scheme of dimension one over k . The genus of a curve C is $g(C) := \dim H^0(C, \omega_C)$.

A *family of curves* $\pi : \mathcal{C} \rightarrow S$ is a proper and flat morphism of stacks whose geometric fibers are curves. If all the geometric fibers of π have the same genus g , then we say that $\pi : \mathcal{C} \rightarrow S$ is a family of curves of genus g (or a family of curves with relative genus g) and we set $g(\mathcal{C}/S) := g$. We will denote by ω_π the relative canonical line bundle of π . Note that any family of curves $\pi : \mathcal{C} \rightarrow S$ with S connected is a family of genus g curves for some $g \geq 0$.

Notation 1.7. Given two integers $g, n \geq 0$, we will denote by $\mathcal{M}_{g,n}$ the stack (over k) whose fiber over a scheme S is the groupoid of families $(\pi : \mathcal{C} \rightarrow S, \underline{\sigma} = \{\sigma_1, \dots, \sigma_n\})$ of n -pointed curves of genus g over S , i.e., $\pi : \mathcal{C} \rightarrow S$ is a family of curves of genus g and $\{\sigma_1, \dots, \sigma_n\}$ are (ordered) sections of π that are fiberwise disjoint.

It is well known that the stack $\mathcal{M}_{g,n}$ is an irreducible algebraic stack, smooth and separated over k , and of dimension $3g - 3 + n$. Moreover, $\mathcal{M}_{g,n}$ is a DM(=Deligne-Mumford) stack if and only if $3g - 3 + n > 0$.

We will denote by $(\pi_{g,n} = \pi : \mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}, \underline{\sigma})$ the universal n -pointed curve over $\mathcal{M}_{g,n}$.

Notation 1.8. A linear algebraic group over k is a group scheme of finite type over k that can be realized as a closed algebraic subgroup of GL_n , or equivalently it is an affine group scheme of finite type over k . We will be dealing almost always with linear algebraic groups that are smooth (which is always the case if $\mathrm{char}(k) = 0$) and connected.

Given a linear algebraic group G , a principal G -bundle over an algebraic stack S is a G -torsor over S , where G acts on the right.

Notation 1.9. In the paper, we introduce several groups and morphisms. To help the reader, we make a table of the main objects together with a reference to their definitions.

ACKNOWLEDGEMENTS. We are grateful to Giulio Codogni, Martina Lanini and Johan Martens for useful conversations. We would like to thank the referee for carefully reading the paper and for giving us constructive comments which helped to improve the readability of the paper.

Symbol	Definition
$\mathcal{D}(G), \mathcal{R}(G), G^{\text{ss}}, G^{\text{ab}}, G, G^{\text{ad}}$	(2.1), (2.3)
Weil group \mathcal{W}_G , Fundamental group $\pi_1(G)$, Center $\mathcal{Z}(G)$	(2.5), (2.15), (2.16)
Maximal tori $T_G, T_{\mathcal{D}(G)}, T_{G^{\text{ss}}}, T_G, T_{G^{\text{ad}}}$	(2.6)
Cocharacter lattices $\Lambda(-)$ and character lattices $\Lambda^*(-)$	(2.8), (2.11)
$\Lambda(T_G)_{\mathbb{Q}}^{\text{ab}}, \Lambda(T_G)_{\mathbb{Q}}^{\text{ss}}, \Lambda^*(T_G)_{\mathbb{Q}}^{\text{ab}}, \Lambda^*(T_G)_{\mathbb{Q}}^{\text{ss}}$	(2.9), (2.12)
$\text{Bil}^s(-)$ and $\text{Bil}^{s,\text{ev}}(-)$	(2.18)
$\text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathcal{W}_G} \subseteq \text{Bil}^s(\Lambda(T_G))^{\mathcal{W}_G}$	Section 2.2
$\text{Bil}^{s,(\text{ev})}(\Lambda(G^{\text{ab}})) \xrightarrow{B_{\text{ab}}^*} \text{Bil}^{s,(\text{ev})}(\Lambda(T_G))^{\mathcal{W}_G}$	Proposition 2.4
$\text{Bil}^{s,(\text{ev})}(\Lambda(T_G))^{\mathcal{W}_G} \xrightarrow{\text{res}_{\mathcal{D}}} \text{Bil}^{s,(\text{ev})}(\Lambda(T_{\mathcal{D}(G)}) \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}_G}$	Proposition 2.4
$\text{Bil}^s(\Lambda(G^{\text{ab}})) \xrightarrow{B_{\text{ab}}^*} \text{Bil}^{s,\mathcal{D}-\text{ev}}(\Lambda(T_G))^{\mathcal{W}_G}$	Corollary 2.6
$\text{Bil}^{s,\mathcal{D}-\text{ev}}(\Lambda(T_G))^{\mathcal{W}_G} \xrightarrow{\text{res}_{\mathcal{D}}} \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}_G}$	Corollary 2.6
$\text{ev}_G^\delta : \text{Bil}^{s,\mathcal{D}-\text{ev}}(\Lambda(T_G))^{\mathcal{W}_G} \rightarrow \Lambda^*(T_G)/\Lambda^*(T_{G^{\text{ad}}})$	Definition/Lemma 2.8
$\text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}_G}$	Definition/Lemma 2.11
$\text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}_G} \xrightarrow{r_G} \text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}_G}$	Definition/Lemma 2.11(i)
$\text{ev}_{\mathcal{D}(G)}^\delta : \text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}_G} \rightarrow \Lambda^*(T_{\mathcal{D}(G)})/\Lambda^*(T_{G^{\text{ad}}})$	Definition/Lemma 2.11(ii)
$\tau_G^\delta : \text{Sym}^2(\Lambda^*(T_G))^{\mathcal{W}_G} \cong \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathcal{W}_G} \rightarrow \text{RPic}(\text{Bun}_{G,g,n}^\delta)$	Theorem 3.3
$\theta_G^\delta : \text{RPic}(\text{Bun}_{G,g,n}^\delta) \rightarrow \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}_G}$	Corollary 3.4
$\gamma_G^\delta : \text{RPic}(\text{Bun}_{G,g,n}^\delta) \rightarrow \text{Bil}^{s,\mathcal{D}-\text{ev}}(\Lambda(T_G))^{\mathcal{W}_G}$	Definition/Lemma 3.5
$\widetilde{H}_{g,n}$ and $i_G^\delta : \Lambda^*(G^{\text{ab}}) \otimes \widetilde{H}_{g,n} \rightarrow \text{RPic}(\text{Bun}_{G,g,n}^\delta)$	Theorem 3.6
$\omega_G^\delta : \text{RPic}(\text{Bun}_{G,g,n}^\delta) \rightarrow \Lambda^*(T_G)/\Lambda^*(T_{G^{\text{ad}}})$	Definition/Lemma 3.7
$\text{NS}(\text{Bun}_{G,g,n}^\delta) \subseteq \Lambda^*(T_G)/\Lambda^*(T_{G^{\text{ad}}}) \oplus \text{Bil}^{s,\mathcal{D}-\text{ev}}(\Lambda(T_G))^{\mathcal{W}_G}$	Definition 3.9
$\phi^{*,\text{NS}}$	Definition/Lemma 3.10
$\text{res}_G^{\text{NS}} : \text{NS}(\text{Bun}_{G,g,n}^\delta) \rightarrow \text{Bil}^{s,\mathcal{D}-\text{ev}}(\Lambda(T_G))^{\mathcal{W}_G}$	Proposition 3.11
$H_{g,n}$ and $j_G^\delta : \Lambda^*(G^{\text{ab}}) \otimes H_{g,n} \rightarrow \text{RPic}(\text{Bun}_{G,g,n}^\delta)$	Theorem 3.12
$\text{Hom}(\pi_1(G), J_C(k)) \xrightarrow{j_G^\delta(C)} \text{Pic}(\text{Bun}_G^\delta(C)) \xrightarrow{c_G^\delta(C)} \text{NS}(\text{Bun}_G^\delta(C))$	Theorem 4.1
$p : \text{NS}(\text{Bun}_G^\delta(C)) \rightarrow \text{Bil}^{s,\text{ev}}(\Lambda(T_{G^{\text{sc}}}))^{\mathcal{W}_G}$	Proposition 4.2
$\text{res}_G^\delta(C), \text{res}_G^\delta(C)^\circ, \text{res}_G^\delta(C)^{\text{NS}}, \text{res}_G^\delta(C)$	Theorem 4.3, (4.7)
$\text{Pic}(\text{Bun}_{G,g,n}^\delta) \xrightarrow{\text{wt}_G^\delta} \Lambda^*(\mathcal{Z}(G)) \xrightarrow{\text{obs}_G^\delta} H^2(\mathfrak{Bun}_{G,g,n}^\delta, \mathbb{G}_m)$	(5.2)
$\text{Pic}(\text{Bun}_G^\delta(C)) \xrightarrow{\text{wt}_G^\delta(C)} \Lambda^*(\mathcal{Z}(G)) \xrightarrow{\text{obs}_G^\delta(C)} H^2(\mathfrak{Bun}_G^\delta(C), \mathbb{G}_m)$	(5.4)
$\text{wt}_G^\delta(C) : \text{Pic}(\text{Bun}_G^\delta(C)) \xrightarrow{c_G^\delta(C)} \text{NS}(\text{Bun}_G^\delta(C)) \xrightarrow{\overline{\text{wt}}_G^\delta(C)} \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})}$	Theorem 5.1
$\text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) \subseteq \text{Bil}^{s,\mathcal{D}-\text{ev}}(\Lambda(T_G))^{\mathcal{W}_G}$	Definition 5.3
$\text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) \xrightarrow{v_G^{\delta,\text{NS}}} \text{NS}(\text{Bun}_{G,g,n}^\delta) \xrightarrow{\omega_G^{\delta,\text{NS}}} \Lambda^*(T_G)/\Lambda^*(T_{G^{\text{ad}}})$	Proposition 5.4
$\Lambda^*(G^{\text{ab}}) \otimes H_{g,n} \xrightarrow{j_G^\delta} \text{RPic}(\mathfrak{Bun}_{G,g,n}^\delta) \xrightarrow{\gamma_G^\delta} \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta)$	Theorem 5.5
$\partial_G^\delta : \text{coker}(\gamma_G^\delta) \rightarrow \text{Hom}\left(\Lambda(G^{\text{ab}}), \frac{\mathbb{Z}}{(2g-2)\mathbb{Z}}\right)$	Theorem 5.7(2)

2. Preliminaries

2.1. Reductive groups

In this subsection we will collect some results on the structure of reductive groups, that will be used in what follows.

A **reductive group** (over $k = \bar{k}$) is a smooth and connected linear algebraic group (over k) which does not contain non-trivial connected normal unipotent alge-

braic subgroups. To any reductive group G , we can associate a cross-like diagram of reductive groups

$$\begin{array}{ccccc}
 & & \mathcal{D}(G) & & \\
 & & \downarrow \mathcal{D} & \searrow & \\
 \mathcal{R}(G) & \hookrightarrow & G & \twoheadrightarrow & G^{\text{ss}} \\
 & \searrow \mathcal{R} & \downarrow \text{ab} & & \\
 & & G^{\text{ab}} & &
 \end{array} \tag{2.1}$$

where

- $\mathcal{D}(G) := [G, G]$ is the derived subgroup of G ;
- $G^{\text{ab}} := G/\mathcal{D}(G)$ is called the abelianization of G ;
- $\mathcal{R}(G)$ is the radical subgroup of G , which is equal (since G is reductive) to the connected component $\mathcal{Z}(G)^o$ of the center $\mathcal{Z}(G)$;
- $G^{\text{ss}} := G/\mathcal{R}(G)$ is called the semisimplification of G .

In the above diagram, the horizontal and vertical lines are short exact sequences of reductive groups, the morphisms $\mathcal{D}(G) \twoheadrightarrow G^{\text{ss}}$ and $\mathcal{R}(G) \twoheadrightarrow G^{\text{ab}}$ are central isogenies of, respectively, semisimple groups and tori with the same kernel which is equal to the finite multiplicative (algebraic) group

$$\mu := \mathcal{D}(G) \cap \mathcal{R}(G) \subset G. \tag{2.2}$$

Since the two semisimple groups $\mathcal{D}(G)$ and G^{ss} are isogenous, they share the same simply-connected cover, that we will denote by G , and the same adjoint quotient, that we will denote by G^{ad} . Hence we have the following tower of central isogenies of semisimple groups:

$$G \twoheadrightarrow \mathcal{D}(G) \twoheadrightarrow G^{\text{ss}} \twoheadrightarrow G^{\text{ad}}. \tag{2.3}$$

The Lie algebra \mathfrak{g} of G splits as

$$\mathfrak{g} = \mathfrak{g}^{\text{ab}} \oplus \mathfrak{g}^{\text{ss}}, \tag{2.4}$$

where \mathfrak{g}^{ab} is the Abelian Lie algebra of the tori $\mathcal{R}(G)$ and G^{ab} , whose dimension is called the Abelian rank of G , and \mathfrak{g}^{ss} is the semisimple Lie algebra of each of the semisimple groups in (2.3), whose rank is called the semisimple rank of G . The semisimple Lie algebra \mathfrak{g}^{ss} decomposes as a direct sum of simple Lie algebras of classical type (*i.e.*, type $A_n, B_n, C_n, D_n, E_6, E_7, E_8, F_4$ or G_2). If G is a semisimple group such that its Lie algebra $\mathfrak{g} = \mathfrak{g}^{\text{ss}}$ is simple, then G is said to be almost-simple.

Remark 2.1. It follows from the universal property of the maximal Abelian quotient G^{ab} and from the universal property of the universal cover G that the morphisms

$$G \rightarrow \mathcal{D}(G) \xrightarrow{\mathcal{D}} G \rightarrow G^{\text{ab}}$$

are covariantly functorial with respect to homomorphisms of reductive groups.

On the other hand, the morphisms

$$\mathcal{R}(G) \xrightarrow{\mathcal{R}} G \xrightarrow{\text{ss}} G^{\text{ss}} \rightarrow G^{\text{ad}}$$

are not functorial with respect to arbitrary homomorphisms of reductive groups, e.g., the inclusion of a maximal torus $T \hookrightarrow G$ does not factor, in general, through $\mathcal{R}(G)$ or, equivalently, does not map to zero in G^{ss} or G^{ad} .

Recall now that all maximal tori of G are conjugate and let us fix one such *maximal torus*, that we call T_G . We will denote by B_G a Borel subgroup of G that contains T_G and by $\mathcal{N}(T_G)$ the normalizer of T_G in G , so that

$$\mathcal{W}_G := \mathcal{N}(T_G)/T_G \tag{2.5}$$

is the Weyl group of G .

The maximal torus T_G induces compatible maximal tori of every semisimple group appearing in (2.3), that we will call, respectively, $T_G, T_{\mathcal{D}(G)}, T_{G^{\text{ss}}}$ and $T_{G^{\text{ad}}}$. These tori fit into the following commutative diagram:

$$\begin{array}{ccccc}
 & & T_G & & \\
 & & \searrow & & \\
 & & T_{\mathcal{D}(G)} & & \\
 & & \downarrow & \searrow & \\
 \mathcal{R}(G) \hookrightarrow & T_G & \twoheadrightarrow & T_{G^{\text{ss}}} & \\
 & \downarrow & & \searrow & \\
 & G^{\text{ab}} & & & T_{G^{\text{ad}}}
 \end{array} \tag{2.6}$$

where the horizontal and vertical lines are short exact sequences of tori, and the diagonal arrows are (central) isogenies of tori. Using the canonical realization (2.5) of the Weyl group (and the similar ones for the semisimple groups in (2.3)), diagram (2.6) induces canonical isomorphisms of Weyl groups

$$\mathcal{W}_G \cong \mathcal{W}_{\mathcal{D}(G)} \cong \mathcal{W}_G \cong \mathcal{W}_{G^{\text{ss}}} \cong \mathcal{W}_{G^{\text{ad}}}. \tag{2.7}$$

By taking the *cocharacter lattices* $\Lambda(-) := \text{Hom}(\mathbb{G}_m, -)$ of the tori in the diagram (2.6), we get the \mathcal{W}_G -equivariant commutative diagram of lattices

$$\begin{array}{ccccc}
 & & \Lambda(T_G) & & \\
 & & \swarrow & & \\
 & & \Lambda(T_{\mathcal{D}(G)}) & & \\
 & & \downarrow \Lambda_{\mathcal{D}} & \searrow & \\
 \Lambda(\mathcal{R}(G)) & \xrightarrow{\Lambda_{\mathcal{R}}} & \Lambda(T_G) & \xrightarrow{\Lambda_{ss}} & \Lambda(T_{G^{ss}}) \\
 & \searrow & \downarrow \Lambda_{ab} & & \searrow \\
 & & \Lambda(G^{ab}) & & \Lambda(T_{G^{ad}})
 \end{array} \tag{2.8}$$

where the horizontal and vertical lines are short exact sequences and the diagonal arrows are finite index inclusions. The above diagram induces a canonical splitting

$$\Lambda(T_G)_{\mathbb{Q}} = \Lambda(T_G)_{\mathbb{Q}}^{ab} \oplus \Lambda(T_G)_{\mathbb{Q}}^{ss}, \tag{2.9}$$

where $\Lambda(T_G)_{\mathbb{Q}}^{ab}$ and $\Lambda(T_G)_{\mathbb{Q}}^{ss}$ are the unique subgroups of $\Lambda(T_G)_{\mathbb{Q}}$ such that

$$\begin{aligned}
 \Lambda(\mathcal{R}(G)) &= \Lambda(T_G) \cap \Lambda(T_G)_{\mathbb{Q}}^{ab} & \text{and} & & \Lambda(T_{\mathcal{D}(G)}) &= \Lambda(T_G) \cap \Lambda(T_G)_{\mathbb{Q}}^{ss}, \\
 \Lambda(G^{ab}) &= p_1(\Lambda(T_G)) & \text{and} & & \Lambda(T_{G^{ss}}) &= p_2(\Lambda(T_G)),
 \end{aligned} \tag{2.10}$$

where p_1 and p_2 are the two projections onto the two factors of (2.9).

In a similar way, by taking the *character lattices* $\Lambda^*(-) := \text{Hom}(-, \mathbb{G}_m)$ (which we will interpret as spaces of integral functionals on $\Lambda(-)$), we get the \mathcal{W}_G -equivariant commutative diagram of lattices

$$\begin{array}{ccccc}
 & & \Lambda^*(T_G) & & \\
 & & \swarrow & & \\
 & & \Lambda^*(T_{\mathcal{D}(G)}) & & \\
 & & \uparrow \Lambda^*_{\mathcal{D}} & \swarrow & \\
 \Lambda^*(\mathcal{R}(G)) & \xleftarrow{\Lambda^*_{\mathcal{R}}} & \Lambda^*(T_G) & \xleftarrow{\Lambda^*_{ss}} & \Lambda^*(T_{G^{ss}}) \\
 & \swarrow & \uparrow \Lambda^*_{ab} & & \swarrow \\
 & & \Lambda^*(G^{ab}) & & \Lambda^*(T_{G^{ad}})
 \end{array} \tag{2.11}$$

where the horizontal and vertical lines are short exact sequences and the diagonal arrows are finite index inclusions. The above diagram induces a canonical splitting

$$\Lambda^*(T_G)_{\mathbb{Q}} = \Lambda^*(T_G)_{\mathbb{Q}}^{ab} \oplus \Lambda^*(T_G)_{\mathbb{Q}}^{ss}, \tag{2.12}$$

where $\Lambda^*(T_G)^{\text{ab}}$ and $\Lambda^*(T_G)^{\text{ss}}$ are the unique subgroups of $\Lambda^*(T_G)_{\mathbb{Q}}$ such that

$$\begin{aligned} \Lambda^*(G^{\text{ab}}) &= \Lambda^*(T_G) \cap \Lambda^*(T_G)^{\text{ab}}_{\mathbb{Q}} & \text{and} & \quad \Lambda^*(T_{G^{\text{ss}}}) = \Lambda^*(T_G) \cap \Lambda^*(T_G)^{\text{ss}}_{\mathbb{Q}}, \\ \Lambda^*(\mathcal{R}(G)) &= p_1(\Lambda^*(T_G)) & \text{and} & \quad \Lambda^*(T_{\mathcal{D}(G)}) = p_2(\Lambda^*(T_G)), \end{aligned} \tag{2.13}$$

where p_1 and p_2 are the two projections onto the two factors of (2.12).

Remark 2.2. There are natural identifications

$$\begin{aligned} \Lambda(T_G) &\cong \Lambda_{\text{coroots}}(\mathfrak{g}^{\text{ss}}) & \text{and} & \quad \Lambda(T_{G^{\text{ad}}}) \cong \Lambda_{\text{coweights}}(\mathfrak{g}^{\text{ss}}), \\ \Lambda^*(T_G) &\cong \Lambda_{\text{weights}}(\mathfrak{g}^{\text{ss}}) & \text{and} & \quad \Lambda^*(T_{G^{\text{ad}}}) \cong \Lambda_{\text{roots}}(\mathfrak{g}^{\text{ss}}), \end{aligned}$$

where $\Lambda_{\text{roots}}(\mathfrak{g}^{\text{ss}})$ (respectively $\Lambda_{\text{coroots}}(\mathfrak{g}^{\text{ss}})$) is the lattice of roots (respectively coroots) of the semisimple Lie algebra \mathfrak{g}^{ss} , and $\Lambda_{\text{weights}}(\mathfrak{g}^{\text{ss}})$ (respectively $\Lambda_{\text{coweights}}(\mathfrak{g}^{\text{ss}})$) is the lattice of weights (respectively coweights) of \mathfrak{g}^{ss} . The two diagrams (2.8) and (2.11), together with the root system of the semisimple Lie algebra \mathfrak{g}^{ss} , are equivalent to the root data of the reductive group G (see [23, Section 19]), and hence they determine completely the reductive group G .

Remark 2.3. It follows from Remark 2.1 that the morphism of lattices

$$\begin{aligned} \Lambda(T_G) &\hookrightarrow \Lambda(T_{\mathcal{D}(G)}) \xrightarrow{\Lambda_{\mathcal{D}}} \Lambda(T_G) \xrightarrow{\Lambda_{\text{ab}}} \Lambda(G^{\text{ab}}) \\ \left(\text{respectively } \Lambda^*(G^{\text{ab}}) \xrightarrow{\Lambda_{\text{ab}}^*} \Lambda^*(T_G) \xrightarrow{\Lambda_{\mathcal{D}}^*} \Lambda^*(T_{\mathcal{D}(G)}) \hookrightarrow \Lambda^*(T_G) \right) \end{aligned} \tag{2.14}$$

are covariantly (respectively contravariantly) functorial with respect to homomorphisms of reductive groups $\phi : G \rightarrow H$ provided that we choose (and this is always possible) the maximal tori T_G and T_H of, respectively, G and H in such a way that $\phi(T_G) \subseteq T_H$.

On the other hand, the morphisms of lattices appearing in (2.8) and in (2.11) and different from the ones in (2.14) are not functorial.

The **fundamental group** $\pi_1(G)$ of G is canonically isomorphic to $\Lambda(T_G)/\Lambda(T_G)$ and it fits into the following (covariantly functorial) short exact sequence of finitely generated Abelian groups:

$$\pi_1(\mathcal{D}(G)) = \frac{\Lambda(T_{\mathcal{D}(G)})}{\Lambda(T_G)} \hookrightarrow \pi_1(G) = \frac{\Lambda(T_G)}{\Lambda(T_G)} \twoheadrightarrow \pi_1(G^{\text{ab}}) = \Lambda(G^{\text{ab}}), \tag{2.15}$$

where the first term is the torsion subgroup of $\pi_1(G)$ and the last term is the torsion-free quotient of $\pi_1(G)$.

Let us now describe the **center** $\mathcal{Z}(G)$ of G , which is a multiplicative (algebraic) group. By taking the centers of the algebraic groups appearing in the

cross-like diagram (2.1), we obtain the (non functorial) cross-like diagram of multiplicative groups

$$\begin{array}{ccccc}
 & & \mathcal{L}(\mathcal{D}(G)) & & \\
 & & \downarrow & \searrow & \\
 \mathcal{R}(G) = \mathcal{L}(G)^{\circ} & \hookrightarrow & \mathcal{L}(G) & \twoheadrightarrow & \mathcal{L}(G^{ss}) = \pi_0(\mathcal{L}(G)) \\
 & \searrow & \downarrow & & \\
 & & G^{ab} & &
 \end{array} \tag{2.16}$$

where the horizontal and vertical lines are short exact sequences, the upper-right diagonal morphism is an isogeny of finite multiplicative groups and the lower-left diagonal arrow is an isogeny of tori. By passing to the character groups $\Lambda^*(-) = \text{Hom}(-, \mathbb{G}_m)$ of the multiplicative groups appearing in (2.16), we get the (non functorial) diagram of finitely generated Abelian groups

$$\begin{array}{ccccc}
 & & \Lambda^*(\mathcal{L}(\mathcal{D}(G))) = \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{ad}})} & & \\
 & & \uparrow \overline{\Lambda^*_{\mathcal{D}}} & \swarrow & \\
 \Lambda^*(\mathcal{R}(G)) & \longleftarrow & \Lambda^*(\mathcal{L}(G)) = \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{ad}})} & \longrightarrow & \Lambda^*(\mathcal{L}(G^{ss})) = \frac{\Lambda^*(T_{G^{ss}})}{\Lambda^*(T_{G^{ad}})} \\
 & \swarrow & \downarrow \overline{\Lambda^*_{ab}} & & \\
 & & \Lambda^*(G^{ab}), & &
 \end{array} \tag{2.17}$$

where the horizontal line is the canonical decomposition of $\Lambda^*(\mathcal{L}(G))$ into its torsion subgroup and torsion-free quotient, the vertical line is exact, the lower-left diagonal arrow is a finite inclusion of lattices, the upper-right diagonal arrow is an inclusion of finite Abelian groups.

2.2. Integral bilinear (even) symmetric forms on $\Lambda(T_G)$

In this subsection we will prove some results on (\mathcal{W}_G -invariant) integral bilinear (even) symmetric forms on the lattice $\Lambda(T_G)$.

Given a lattice Λ of rank r (i.e., $\Lambda \cong \mathbb{Z}^r$), we denote the lattice of integral bilinear (respectively even) symmetric forms on Λ by

$$\begin{aligned}
 \text{Bil}^s(\Lambda) &:= \{b : \Lambda \otimes \Lambda \rightarrow \mathbb{Z} \text{ such that } b \text{ is symmetric}\}, \\
 \text{Bil}^{s, \text{ev}}(\Lambda) &:= \{b \in \text{Bil}^s(\Lambda) : b(x, x) \text{ is even for any } x \in \Lambda\}.
 \end{aligned} \tag{2.18}$$

Given $b \in \text{Bil}^s(\Lambda)$, we will denote by $b^{\mathbb{Q}} : \Lambda_{\mathbb{Q}} \otimes \Lambda_{\mathbb{Q}} \rightarrow \mathbb{Q}$ the rational extension of b to the \mathbb{Q} -vector space $\Lambda_{\mathbb{Q}} := \Lambda \otimes_{\mathbb{Z}} \mathbb{Q}$.

The above lattices (2.18), which are contravariantly functorial with respect to morphisms of lattices, can be described in terms of the dual lattice $\Lambda^* := \text{Hom}(\Lambda, \mathbb{Z})$ in the following way. Consider the following lattices (of rank $\binom{r+1}{2}$)

$$(\Lambda^* \otimes \Lambda^*)^s \subset \Lambda^* \otimes \Lambda^* \quad \text{and} \quad \text{Sym}^2(\Lambda^*) := \frac{\Lambda^* \otimes \Lambda^*}{\langle \chi \otimes \mu - \mu \otimes \chi \rangle}, \quad (2.19)$$

where $(\Lambda^* \otimes \Lambda^*)^s$ is the subspace of symmetric tensors of $\Lambda^* \otimes \Lambda^*$, *i.e.*, tensors that are invariant under the involution $\chi \otimes \mu \mapsto \mu \otimes \chi$. We will denote the elements of $\text{Sym}^2(\Lambda^*)$ by $\chi \cdot \mu := [\chi \otimes \mu]$.

The lattices in (2.18) are isomorphic to those in (2.19) via the following isomorphisms:

$$\begin{aligned} (\Lambda^* \otimes \Lambda^*)^s &\xrightarrow{\cong} \text{Bil}^s(\Lambda) \\ \chi \otimes \mu &\mapsto (\chi \otimes \mu)(x \otimes y) := \chi(x)\mu(y), \quad \text{and} \\ \text{Sym}^2 \Lambda^* &\xrightarrow{\cong} \text{Bil}^{s,\text{ev}}(\Lambda) \\ \chi \cdot \mu &\mapsto (\chi \cdot \mu)(x \otimes y) := \chi(x)\mu(y) + \mu(x)\chi(y). \end{aligned} \quad (2.20)$$

In terms of the isomorphisms (2.20), the inclusion $\text{Bil}^{s,\text{ev}}(\Lambda) \subset \text{Bil}^s(\Lambda)$ corresponds to the injective morphism

$$\begin{aligned} \psi : \text{Sym}^2(\Lambda^*) &\hookrightarrow (\Lambda^* \otimes \Lambda^*)^s, \\ \chi \cdot \mu &\mapsto \chi \otimes \mu + \mu \otimes \chi. \end{aligned} \quad (2.21)$$

If we fix a basis $\{\chi_i\}_{i=1}^r$ of Λ^* , then

$$\begin{aligned} \{\{\chi_i \otimes \chi_i\} \cup \{\chi_i \otimes \chi_j + \chi_j \otimes \chi_i\}_{i < j}\} &\text{ is a basis of } (\Lambda^* \otimes \Lambda^*)^s, \\ \{\chi_i \cdot \chi_j\}_{i \leq j} &\text{ is a basis of } \text{Sym}^2(\Lambda^*). \end{aligned} \quad (2.22)$$

Using the above basis, it follows that

$$\text{coker}(\psi) = (\mathbb{Z}/2\mathbb{Z})^r. \quad (2.23)$$

We now come back to the setting of Subsection 2.1. Let G be a reductive group with maximal torus $T_G \subset G$ and consider the natural action of the Weyl group \mathscr{W}_G on $\Lambda(T_G)$ and on $\Lambda^*(T_G)$. We now want to describe the lattices

$$\begin{aligned} \text{Bil}^s(\Lambda(T_G))^{\mathscr{W}_G} &\cong ((\Lambda^*(T_G) \otimes \Lambda^*(T_G))^s)^{\mathscr{W}_G} \quad \text{and} \\ \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathscr{W}_G} &\cong \text{Sym}^2(\Lambda^*(T_G))^{\mathscr{W}_G} \end{aligned}$$

of \mathscr{W}_G -invariant integral bilinear (respectively even) symmetric forms on $\Lambda(T_G)$.

Proposition 2.4. *With the above notation, we have an exact sequence of lattices*

$$\begin{aligned} \text{Bil}^{s,(\text{ev})}(\Lambda(G^{\text{ab}})) \xrightarrow{B_{\text{ab}}^*} \text{Bil}^{s,(\text{ev})}(\Lambda(T_G)) \mathscr{W}_G \xrightarrow{\text{res}_{\mathscr{D}}} \text{Bil}^{s,(\text{ev})}(\Lambda(T_{\mathscr{D}(G)}) | \Lambda(T_{G^{\text{ss}}})) \mathscr{W}_G, \\ b(- \otimes -) \mapsto b(\Lambda_{\text{ab}}(-) \otimes \Lambda_{\text{ab}}(-)) \\ b \mapsto b|_{\Lambda(T_{\mathscr{D}(G)}) \otimes \Lambda(T_{\mathscr{D}(G)})} \end{aligned} \tag{2.24}$$

where $\Lambda_{\text{ab}} : \Lambda(T_G) \rightarrow \Lambda(G^{\text{ab}})$ is the homomorphism defined in (2.8) and

$$\begin{aligned} & \text{Bil}^{s,(\text{ev})}(\Lambda(T_{\mathscr{D}(G)}) | \Lambda(T_{G^{\text{ss}}})) \mathscr{W}_G \\ := & \left\{ b \in \text{Bil}^{s,(\text{ev})}(\Lambda(T_{\mathscr{D}(G)})) \mathscr{W}_G : b|_{\Lambda(T_{\mathscr{D}(G)}) \otimes \Lambda(T_{G^{\text{ss}}})} \text{ is integral } \right\}. \end{aligned}$$

Moreover, the exact sequence (2.24) is contravariant with respect to homomorphisms of reductive groups $\phi : H \rightarrow G$ such that $\phi(T_H) \subseteq T_G$.

The above notation $\text{Bil}^{s,(\text{ev})}$ means that the result applies by putting Bil^s everywhere or by putting $\text{Bil}^{s,\text{ev}}$ everywhere.

In the proof of the above proposition, we will use the following

Lemma 2.5. *Let G be a reductive group with maximal torus $T_G \subset G$ and consider the natural action of the Weyl group \mathscr{W}_G on $\Lambda(T_G)$ and on $\Lambda^*(T_G)$. Then we have isomorphisms*

$$\Lambda_{\mathscr{R}} : \Lambda(\mathscr{R}(G)) \xrightarrow{\cong} \Lambda(T_G) \mathscr{W}_G \quad \text{and} \quad \Lambda_{\text{ab}}^* : \Lambda^*(G^{\text{ab}}) \xrightarrow{\cong} \Lambda^*(T_G) \mathscr{W}_G.$$

Proof. The second isomorphism is proved in [14, Lemma 2.1.1]. The proof of the first isomorphism is similar. \square

Proof of Proposition 2.4. Clearly, the morphism B_{ab}^* is injective and $\text{res}_{\mathscr{D}} \circ B_{\text{ab}}^* = 0$.

In order to complete the proof, we will need the following

Claim: If $b \in \text{Bil}^s(\Lambda(T_G)) \mathscr{W}_G$ then $b|_{\Lambda(\mathscr{R}(G)) \otimes \Lambda(T_{\mathscr{D}(G)})} \equiv 0$. In particular, $b|_{\Lambda(T_G)_{\mathbb{Q}}^{\text{ab}} \otimes \Lambda(T_G)_{\mathbb{Q}}^{\text{ss}}} \equiv 0$.

Indeed, for any $x \in \Lambda(\mathscr{R}(G))$, the restriction $b(x \otimes -) : \Lambda(T_G) \rightarrow \mathbb{Z}$ is \mathscr{W}_G -invariant since b is \mathscr{W}_G -invariant and x is fixed by \mathscr{W}_G (because the action of \mathscr{W}_G is trivial on $\Lambda(\mathscr{R}(G))$). Hence, Lemma 2.5 implies that $b(x \otimes -)$ is the pull-back of an integral functional on $\Lambda(G^{\text{ab}})$, or in other words that $b(x \otimes -)|_{\Lambda(T_{\mathscr{D}(G)})} \equiv 0$. Since this is true for any $x \in \Lambda(\mathscr{R}(G))$, we get that $b|_{\Lambda(\mathscr{R}(G)) \otimes \Lambda(T_{\mathscr{D}(G)})} \equiv 0$. The last assertion follows from the fact that $\Lambda(\mathscr{R}(G))_{\mathbb{Q}} = \Lambda(T_G)_{\mathbb{Q}}^{\text{ab}}$ and $\Lambda(T_{\mathscr{D}(G)})_{\mathbb{Q}} = \Lambda(T_G)_{\mathbb{Q}}^{\text{ss}}$.

We now go back to proof of the proposition. Let us first prove that the sequence (2.24) is exact in the middle, *i.e.*, $\ker(\text{res}_{\mathscr{D}}) \subseteq \text{Im}(B_{\text{ab}}^*)$. Consider a form $b \in \text{Bil}^{s,(\text{ev})}(\Lambda(T_G)) \mathscr{W}_G$ such that $\text{res}_{\mathscr{D}}(b) = 0$. This assumption, together with the above claim, implies that $b|_{\Lambda(T_{\mathscr{D}(G)}) \otimes \Lambda(T_G)} \equiv 0$, which implies that b is the pull-back of an integral bilinear (respectively even) symmetric form on $\Lambda(G^{\text{ab}})$.

Let us now prove that the morphism $\text{res}_{\mathcal{D}}$ is well-defined, *i.e.*, that $\text{res}_{\mathcal{D}}(b)$ is integral on $\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_{G^{ss}})$ for every $b \in \text{Bil}^{s,(\text{ev})}(\Lambda(T_G))^{\mathcal{W}_G}$. For any element $y \in \Lambda(T_{\mathcal{D}(G)})$, consider the integral functional $b(y \otimes -) : \Lambda(T_G) \rightarrow \mathbb{Z}$. By the claim, we have that $b(y \otimes -)|_{\Lambda(T_{\mathcal{D}(G)})} \equiv 0$, which implies that $b(y \otimes -)$ is the restriction of an integral functional on $\Lambda(T_{G^{ss}})$. Since this is true for any $y \in \Lambda(T_{\mathcal{D}(G)})$, we deduce that $\text{res}_{\mathcal{D}}(b)$ is integral on $\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_{G^{ss}})$.

In order to show that the sequence (2.24) is exact, it remains to prove that $\text{res}_{\mathcal{D}}$ is surjective. Let $\tilde{b} \in \text{Bil}^{s,(\text{ev})}(\Lambda(T_{\mathcal{D}(G)})|_{\Lambda(T_{G^{ss}})})^{\mathcal{W}_G}$. Since $\tilde{b}^{\mathbb{Q}}$ is integral on $\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_{G^{ss}})$, by composing $\tilde{b}^{\mathbb{Q}}$ with the surjection $\Lambda(T_G) \rightarrow \Lambda(T_{G^{ss}})$ we get a \mathcal{W}_G -invariant (respectively even) symmetric integral form

$$\widehat{b} : \Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_G) + \Lambda(T_G) \otimes \Lambda(T_{\mathcal{D}(G)}) \rightarrow \mathbb{Z}. \tag{2.25}$$

Now consider the \mathcal{W}_G -equivariant short exact sequence of lattices

$$\begin{aligned} \text{Bil}^{s,(\text{ev})}(\Lambda(G^{\text{ab}})) \hookrightarrow \text{Bil}^{s,(\text{ev})}(\Lambda(T_G)) &\rightarrow \text{Hom}^{s,(\text{ev})}(\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_G) + \Lambda(T_G) \otimes \Lambda(T_{\mathcal{D}(G)}), \mathbb{Z}) \\ b &\mapsto b|_{\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_G) + \Lambda(T_G) \otimes \Lambda(T_{\mathcal{D}(G)})} \end{aligned} \tag{2.26}$$

where $\text{Hom}^{s,(\text{ev})}(\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_G) + \Lambda(T_G) \otimes \Lambda(T_{\mathcal{D}(G)}), \mathbb{Z})$ is the lattice of even symmetric integral forms on $\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_G) + \Lambda(T_G) \otimes \Lambda(T_{\mathcal{D}(G)}) \subseteq \Lambda(T_G) \otimes \Lambda(T_G)$. Since the action of \mathcal{W}_G is trivial, we have that

$$H^1(\mathcal{W}_G, \text{Bil}^{s,(\text{ev})}(\Lambda(G^{\text{ab}}))) = \text{Hom}(\mathcal{W}_G, \text{Bil}^{s,(\text{ev})}(\Lambda(G^{\text{ab}}))), \tag{2.27}$$

and the last group is zero since \mathcal{W}_G is a finite group and $\text{Bil}^{s,(\text{ev})}(\Lambda(G^{\text{ab}}))$ is torsion-free. By taking the long exact sequence in \mathcal{W}_G -cohomology associated to the exact sequence (2.26) and using the vanishing $H^1(\mathcal{W}_G, \text{Bil}^{s,(\text{ev})}(\Lambda(G^{\text{ab}}))) = 0$, we get a surjection

$$\text{Bil}^{s,(\text{ev})}(\Lambda(T_G))^{\mathcal{W}_G} \twoheadrightarrow \text{Hom}^{s,(\text{ev})}(\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_G) + \Lambda(T_G) \otimes \Lambda(T_{\mathcal{D}(G)}), \mathbb{Z})^{\mathcal{W}_G}.$$

Hence, the form \widehat{b} of (2.25) is the restriction of a form $b \in \text{Bil}^{s,(\text{ev})}(\Lambda(T_G))^{\mathcal{W}_G}$. By construction we have that $\text{res}_{\mathcal{D}}(b) = \tilde{b}$, which concludes the proof of the surjectivity of $\text{res}_{\mathcal{D}}$.

Finally, the (contravariant) functoriality of the exact sequence (2.24) follows from the fact that $\text{Bil}^{s,(\text{ev})}(\Lambda(T_G))^{\mathcal{W}_G}$ is functorial by [3, Lemma 4.3.1] (and the discussion that follows) while $\text{Bil}^{s,(\text{ev})}(\Lambda(G^{\text{ab}}))$ and the morphism B_{ab}^* are functorial by Remark 2.3. □

By combining the two exact sequences of Proposition 2.4, we get a new exact sequence:

Corollary 2.6. *With the above notation, we have an exact sequence of lattices*

$$\text{Bil}^s(\Lambda(G^{\text{ab}})) \xleftarrow{B_{\text{ab}}^*} \text{Bil}^{s, \mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathcal{W}_G} \xrightarrow{\text{res}_{\mathcal{D}}} \text{Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)})|_{\Lambda(T_{G^{ss}})})^{\mathcal{W}_G}, \tag{2.28}$$

where

$$\text{Bil}^{s, \mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathscr{W}_G} := \{b \in \text{Bil}^s(\Lambda(T_G))^{\mathscr{W}_G} : \text{res}_{\mathcal{D}}(b) \text{ is even} \}.$$

Moreover, the exact sequence (2.28) is contravariant with respect to homomorphisms of reductive groups $\phi : H \rightarrow G$ such that $\phi(T_H) \subseteq T_G$.

Proof. Consider the commutative diagram of Abelian groups

$$\begin{array}{ccccc} \text{Bil}^s(\Lambda(G^{\text{ab}})) \hookrightarrow & \text{res}_{\mathcal{D}}^{-1}(\text{Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G}) & \twoheadrightarrow & \text{Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} \\ \parallel & \downarrow & & \downarrow \\ \text{Bil}^s(\Lambda(G^{\text{ab}})) \hookrightarrow & \text{Bil}^s(\Lambda(T_G))^{\mathscr{W}_G} & \xrightarrow{\text{res}_{\mathcal{D}}} & \text{Bil}^s(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} \end{array}$$

where the rows are exact and the columns are the obvious inclusions. The bottom row is the non-even version the exact sequence (2.24). By definition, we have that

$$\text{res}_{\mathcal{D}}^{-1}(\text{Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G}) = \text{Bil}^{s, \mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathscr{W}_G},$$

and hence the top row is the required sequence (2.28). □

Corollary 2.7. *The ranks of $\text{Bil}^{s, \text{ev}}(\Lambda(T_G))^{\mathscr{W}_G}$ and of $\text{Bil}^{s, \mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathscr{W}_G}$ are equal to*

$$\binom{\dim G^{\text{ab}} + 1}{2} + |\{\text{simple factors of } \mathfrak{g}^{\text{ss}}\}|.$$

Proof. Since $\text{Bil}^{s, \text{ev}}(\Lambda(T_G))^{\mathscr{W}_G} \subseteq \text{Bil}^{s, \mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathscr{W}_G} \subseteq \text{Bil}^s(\Lambda(T_G))^{\mathscr{W}_G}$ are finite index inclusions, it is enough to prove the result for $\text{Bil}^{s, \text{ev}}(\Lambda(T_G))^{\mathscr{W}_G}$. Proposition 2.4 implies that

$$\begin{aligned} \text{rk Bil}^{s, \text{ev}}(\Lambda(T_G))^{\mathscr{W}_G} &= \text{rk Bil}^{s, \text{ev}}(\Lambda(G^{\text{ab}})) + \text{rk Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} \\ &= \binom{\dim G^{\text{ab}} + 1}{2} + \text{rk Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G}. \end{aligned}$$

We conclude observing that we have finite index inclusions

$$\text{Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} \subseteq \text{Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)}))^{\mathscr{W}_G} \subseteq \text{Bil}^{s, \text{ev}}(\Lambda(T_G))^{\mathscr{W}_G}$$

and that the last lattice has rank equal to the number of simple factors of \mathfrak{g}^{ss} (see, e.g., [14, Lemma 2.2.1]). □

We now define evaluation homomorphisms from the exact sequence (2.28) onto the vertical exact sequence in (2.17).

Definition/Lemma 2.8. Fix the same notation as above. Let $\delta \in \pi_1(G)$ and set $\delta^{ss} := \pi_1(ss)(\delta) \in \pi_1(G^{ss})$ and $\delta^{ab} := \pi_1(ab)(\delta) \in \pi_1(G^{ab})$. There is a (non functorial) commutative diagram with exact rows

$$\begin{array}{ccccc}
 \text{Bil}^s(\Lambda(G^{ab})) & \xrightarrow{B_{ab}^*} & \text{Bil}^{s, \mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathcal{W}_G} & \xrightarrow{\text{res}_{\mathcal{D}}} & \text{Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{ss}}))^{\mathcal{W}_G} \\
 \downarrow \text{ev}_{G^{ab}}^{\delta^{ab}} & & \downarrow \text{ev}_G^\delta & & \downarrow \text{ev}_{\mathcal{D}(G)}^\delta \\
 \Lambda^*(G^{ab}) & \xrightarrow{\Lambda_{ab}^*} & \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{ad}})} & \xrightarrow{\Lambda_{\mathcal{D}}^*} & \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{ad}})}
 \end{array}$$

where the vertical arrows, called evaluation homomorphisms, are defined as follows:

- (i) $\text{ev}_{G^{ab}}^{\delta^{ab}}(b) = b(\delta^{ab} \otimes -)$;
- (ii) $\text{ev}_G^\delta(b) = b(\delta \otimes -) := [b(d \otimes -)]$, for some lifting $d \in \Lambda(T_G)$ of $\delta \in \pi_1(G)$;
- (iii) $\text{ev}_{\mathcal{D}(G)}^\delta(b) = b(\delta^{ss} \otimes -) := [b^{\mathbb{Q}}(d^{ss} \otimes -) |_{\Lambda(T_{\mathcal{D}(G)})}]$, for some lifting $d^{ss} \in \Lambda(T_{G^{ss}})$ of $\delta^{ss} \in \pi_1(G^{ss})$.

Note that the notation in the above Definition/Lemma is coherent since if G is a torus then $\text{ev}_G^\delta = \text{ev}_{G^{ab}}^{\delta^{ab}}$ and if G is semisimple then $\text{ev}_G^\delta = \text{ev}_{\mathcal{D}(G)}^\delta$.

In order to prove that the last two evaluation homomorphisms are well-defined, we will need the following:

Lemma 2.9. If $b \in \text{Bil}^{s, \text{ev}}(\Lambda(T_G))^{\mathcal{W}_G}$, then its rational extension $b^{\mathbb{Q}}$ is integral on $\Lambda(T_G) \otimes \Lambda(T_{G^{ad}}) + \Lambda(T_{G^{ad}}) \otimes \Lambda(T_G) \subseteq \Lambda(T_G)_{\mathbb{Q}} \otimes \Lambda(T_G)_{\mathbb{Q}}$.

Proof. See [3, Lemma 4.3.4]. □

Proof of Definition/Lemma 2.8. The fact that the evaluation homomorphism ev_G^δ (respectively $\text{ev}_{\mathcal{D}(G)}^\delta$) is well-defined follows from the fact that any two lifts of δ (respectively of δ^{ss}) differ by an element $e \in \Lambda(T_G)$, together with Lemma 2.9 which implies that $b^{\mathbb{Q}}(e \otimes -)$ is integral on $\Lambda(T_{G^{ad}})$.

The commutativity of the left square follows from the fact that if $b \in \text{Bil}^s(\Lambda(G^{ab}))$ then

$$\begin{aligned}
 \text{ev}_G^\delta(B_{ab}^*(b)) &= [B_{ab}^*(b)(d \otimes -)] = [b(\delta^{ab} \otimes \Lambda_{ab}(-))] \\
 &= \overline{\Lambda_{ab}^*}(b(\delta^{ab} \otimes -)) = \overline{\Lambda_{ab}^*}(\text{ev}_{G^{ab}}^{\delta^{ab}}(b)),
 \end{aligned}$$

where we have used that any lift $d \in \Lambda(T_G)$ of δ satisfies $\Lambda_{ab}(d) = \delta^{ab}$.

Next, observe that, by (2.10), any lift $d \in \Lambda(T_G)$ of $\delta \in \pi_1(G)$ decomposes as $d = \delta^{ab} + d^{ss}$, where $d^{ss} := p_2(d) \in \Lambda(T_{G^{ss}})$ has the property that its class in $\pi_1(G^{ss})$ coincides with δ^{ss} . Therefore, the commutativity of the right square follows since for any $b \in \text{Bil}^{s, \mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathcal{W}G}$ we have that

$$\begin{aligned} \overline{\Lambda^*_{\mathcal{D}}}\left(\text{ev}_G^\delta(b)\right) &= [b(d \otimes -)|_{\Lambda(T_{\mathcal{D}(G)})}] \\ &= [b^{\mathbb{Q}}(d^{ss} \otimes -)|_{\Lambda(T_{\mathcal{D}(G)})}] = \text{ev}_{\mathcal{D}(G)}^\delta(\text{res}_{\mathcal{D}}(b)), \end{aligned}$$

where we have used that $b^{\mathbb{Q}}(\delta^{ab} \otimes -)|_{\Lambda(T_{\mathcal{D}(G)})} = 0$ by the claim in the proof of Proposition 2.4. □

Remark 2.10. The last evaluation homomorphism $\text{ev}_{\mathcal{D}(G)}^\delta$ can be compared with the evaluation homomorphisms of the semisimple groups $\mathcal{D}(G)$ and G^{ss} in the following way. First all, note that we have injective restriction homomorphisms

$$\text{Bil}^{s, (\text{ev})}(\Lambda(T_{G^{ss}}))^{\mathcal{W}G} \hookrightarrow \text{Bil}^{s, (\text{ev})}(\Lambda(T_{\mathcal{D}(G)})|_{\Lambda(T_{G^{ss}})})^{\mathcal{W}G} \hookrightarrow \text{Bil}^{s, (\text{ev})}(\Lambda(T_{\mathcal{D}(G)}))^{\mathcal{W}G}.$$

Then we have that:

- (1) For any $\delta \in \pi_1(G)$, the following diagram is commutative:

$$\begin{array}{ccc} \text{Bil}^{s, \text{ev}}(\Lambda(T_{G^{ss}}))^{\mathcal{W}G} & \hookrightarrow & \text{Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)})|_{\Lambda(T_{G^{ss}})})^{\mathcal{W}G} \\ \downarrow \text{ev}_G^{\delta^{ss}} & & \downarrow \text{ev}_{\mathcal{D}(G)}^\delta \\ \frac{\Lambda^*(T_{G^{ss}})}{\Lambda^*(T_{G^{\text{ad}}})} & \hookrightarrow & \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}})} \end{array}$$

where δ^{ss} is the image of δ in $\pi_1(G^{ss})$;

- (2) For any $\delta \in \pi_1(G)$ which is the image of a (necessarily unique) element $\delta^{\mathcal{D}} \in \pi_1(\mathcal{D}(G))$ (which happens precisely when δ is a torsion element of $\pi_1(G)$, see (2.15)), then we have the following commutative diagram

$$\begin{array}{ccc} \text{Bil}^{s, \text{ev}}(\Lambda(T_{\mathcal{D}(G)})|_{\Lambda(T_{G^{ss}})})^{\mathcal{W}G} & \hookrightarrow & \text{Bil}^{s, \text{ev}}(\Lambda(T_{G^{ss}}))^{\mathcal{W}G} \\ \downarrow \text{ev}_{\mathcal{D}(G)}^\delta & & \downarrow \text{ev}_{\mathcal{D}(G)}^{\delta^{\mathcal{D}}} \\ \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}})} & \xlongequal{\quad\quad\quad} & \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}})}. \end{array}$$

The homomorphism $\text{ev}_{\mathcal{D}(G)}^\delta$ of Definition/Lemma 2.8 can be extended to a slightly larger lattice, as we now show.

Definition/Lemma 2.11. *Fix the same notation as above. Consider the lattice*

$$\begin{aligned} & \text{Bil}^{s,-\text{ev}} \left(\Lambda(T_{\mathcal{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}}) \right)^{\mathcal{W}G} \\ & := \left\{ b \in \text{Bil}^s \left(\Lambda(T_{\mathcal{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}}) \right)^{\mathcal{W}G} : b_{\Lambda(T_G) \otimes \Lambda(T_G)} \text{ is even} \right\}. \end{aligned}$$

(i) *The natural inclusion*

$$r_G : \text{Bil}^{s,\text{ev}} \left(\Lambda(T_{\mathcal{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}}) \right)^{\mathcal{W}G} \hookrightarrow \text{Bil}^{s,-\text{ev}} \left(\Lambda(T_{\mathcal{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}}) \right)^{\mathcal{W}G}$$

has an elementary 2-Abelian cokernel;

(ii) *For any $\delta \in \pi_1(G)$, the evaluation homomorphism*

$$\text{ev}_{\mathcal{D}(G)}^\delta : \text{Bil}^{s,\text{ev}} \left(\Lambda(T_{\mathcal{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}}) \right)^{\mathcal{W}G} \rightarrow \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}})}$$

of Definition/Lemma 2.8 can be extended to a homomorphism

$$\begin{aligned} \tilde{\text{ev}}_{\mathcal{D}(G)}^\delta : \text{Bil}^{s,-\text{ev}} \left(\Lambda(T_{\mathcal{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}}) \right)^{\mathcal{W}G} & \longrightarrow \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}})}, \\ b \mapsto b(\delta^{\text{ss}} \otimes -) & := [b^{\mathbb{Q}}(d^{\text{ss}} \otimes -)_{\mid \Lambda(T_{\mathcal{D}(G)})}], \end{aligned}$$

where $d^{\text{ss}} \in \Lambda(T_{G^{\text{ss}}})$ is any lifting of $\delta^{\text{ss}} \in \pi_1(G^{\text{ss}})$.

Proof. Part (i) follows from the inclusions

$$\begin{aligned} \text{Bil}^{s,\text{ev}} \left(\Lambda(T_{\mathcal{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}}) \right)^{\mathcal{W}G} & \xrightarrow{r_G} \text{Bil}^{s,-\text{ev}} \left(\Lambda(T_{\mathcal{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}}) \right)^{\mathcal{W}G} \\ & \subseteq \text{Bil}^s \left(\Lambda(T_{\mathcal{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}}) \right)^{\mathcal{W}G} \end{aligned}$$

together with (2.23).

Part (ii): the fact that $\tilde{\text{ev}}_G^\delta$ is well-defined follows from the fact that any two lifts of δ^{ss} differ by an element $e \in \Lambda(T_G)$, together with Lemma 2.9 which implies that $b^{\mathbb{Q}}(e \otimes -)$ is integral on $\Lambda(T_{G^{\text{ad}}})$. The fact that $\tilde{\text{ev}}_{\mathcal{D}(G)}^\delta \circ r_G = \text{ev}_{\mathcal{D}(G)}^\delta$ is obvious from the definitions. □

3. The universal moduli stack $\text{Bun}_{G,g,n}$ and its Picard group

Let G be a reductive group over $k = \bar{k}$. We denote by $\text{Bun}_{G,g,n}$ the universal moduli stack of G -bundles over n -marked curves of genus g . More precisely, for any scheme S , $\text{Bun}_{G,g,n}(S)$ is the groupoid of triples $(\mathcal{C} \rightarrow S, \underline{\sigma}, E)$, where $(\pi : \mathcal{C} \rightarrow S, \underline{\sigma} = \{\sigma_1, \dots, \sigma_n\})$ is a family of n -pointed curves of genus g over S and E is a G -bundle on \mathcal{C} . We will denote by $(\pi : \mathcal{C}_{G,g,n} \rightarrow \text{Bun}_{G,g,n}, \underline{\sigma}, \mathcal{E})$ the universal family of G -bundles.

By definition, we have a forgetful surjective morphism

$$\begin{aligned} \Phi_G (= \Phi_{G,g,n}) : \text{Bun}_{G,g,n} &\longrightarrow \mathcal{M}_{g,n} \\ (\mathcal{C} \rightarrow S, \underline{\sigma}, E) &\mapsto (\mathcal{C} \rightarrow S, \underline{\sigma}) \end{aligned} \tag{3.1}$$

onto the moduli stack $\mathcal{M}_{g,n}$ of n -marked curves of genus g . Note that the universal n -marked curve $(\mathcal{C}_{G,g,n} \rightarrow \text{Bun}_{G,g,n}, \underline{\sigma})$ over $\text{Bun}_{G,g,n}$ is the pull-back of the universal n -marked curve $(\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}, \underline{\sigma})$ over $\mathcal{M}_{g,n}$.

Any morphism of reductive groups $\phi : G \rightarrow H$ determines a morphism of stacks over $\mathcal{M}_{g,n}$

$$\begin{aligned} \phi_{\#} (= \phi_{\#,g,n}) : \text{Bun}_{G,g,n} &\longrightarrow \text{Bun}_{H,g,n} \\ (\mathcal{C} \rightarrow S, \underline{\sigma}, E) &\mapsto (\mathcal{C} \rightarrow S, \underline{\sigma}, (E \times H)/G) \end{aligned} \tag{3.2}$$

where the (right) action of G on $E \times H$ is $(p, h).g := (p.g, \phi(g)^{-1}h)$.

The fiber of $\Phi_{G,g,n}$ over a n -pointed curve $(C, p_1, \dots, p_n) \in \mathcal{M}_{g,n}(k)$ is equal to the k -stack $\text{Bun}_G(C)$ of G -bundles on C , i.e., the stack over k whose S -points $\text{Bun}_G(C)(S)$ is the groupoid of G -bundles on $C_S := C \times_k S$ for any k -scheme S . For any morphism of reductive groups $\phi : G \rightarrow H$, the restriction of the morphism $\phi_{\#,g,n}$ to the fiber over $(C, p_1, \dots, p_n) \in \mathcal{M}_{g,n}(k)$ gives rise to a morphism

$$\phi_{\#}(C) : \text{Bun}_G(C) \rightarrow \text{Bun}_H(C).$$

We collect in the following theorem the geometric properties of $\text{Bun}_{G,g,n}$ and of the forgetful morphism $\Phi_{G,g,n}$.

Theorem 3.1. *Let G be a reductive group.*

- (1) *The morphism $\Phi_{G,g,n}$ is locally of finite presentation, smooth, with affine and finitely presented relative diagonal;*
- (2) *There is a functorial decomposition into connected components*

$$\Phi_{G,g,n} : \coprod_{\delta \in \pi_1(G)} \text{Bun}_{G,g,n}^{\delta} \xrightarrow{\Phi_{G,g,n}^{\delta}} \mathcal{M}_{g,n}. \tag{3.3}$$

Similarly, the fiber $\text{Bun}_G(C)$ of $\Phi_{G,g,n}$ over $(C, p_1, \dots, p_n) \in \mathcal{M}_{g,n}(k)$ admits a functorial decomposition into connected components

$$\text{Bun}_G(C) = \coprod_{\delta \in \pi_1(G)} \text{Bun}_G^{\delta}(C);$$

- (3) *For each $\delta \in \pi_1(G)$, the stack $\text{Bun}_{G,g,n}^{\delta}$ is smooth and integral of relative dimension over $\mathcal{M}_{g,n}$ equal to $(g - 1) \dim G$;*
- (4) *$\Phi_G^{\delta} : \text{Bun}_{G,g,n}^{\delta} \rightarrow \mathcal{M}_{g,n}$ is of finite type (or equivalently quasi-compact) for any (or equivalently for some) $\delta \in \pi_1(G)$ if and only if G is a torus;*

- (5) For any $\delta \in \pi_1(G)$ the morphism $\Phi_G^\delta : \text{Bun}_{G,g,n}^\delta \rightarrow \mathcal{M}_{g,n}$ is fpqc (i.e., faithfully flat and locally quasi-compact) and cohomologically flat in degree zero (i.e., the natural morphism $(\Phi_G^\delta)^\# : \mathcal{O}_{\mathcal{M}_{g,n}} \rightarrow (\Phi_G^\delta)_*(\mathcal{O}_{\text{Bun}_{G,g,n}^\delta})$ is a universal isomorphism).

Proof. See [14, Section 3]. □

3.1. The Picard group of $\text{Bun}_{G,g,n}^\delta$

The aim of this subsection is to recall the results on the Picard group of $\text{Bun}_{G,g,n}^\delta$ obtained in [14]. We will focus on the case $g \geq 1$; the case $g = 0$ is easier to deal with and is completely described in [14, Theorem D].

Note that the Picard group of $\mathcal{M}_{g,n}$ is well-known up to torsion (and completely known if $\text{char}(k) \neq 2$ by [13]) and the pull-back morphism

$$(\Phi_G^\delta)^* : \text{Pic}(\mathcal{M}_{g,n}) \rightarrow \text{Pic}(\text{Bun}_{G,g,n}^\delta)$$

is injective since Φ_G^δ is fpqc and cohomologically flat in degree zero by Theorem 3.1(5). Therefore, we can focus our attention onto the relative Picard group

$$\text{RPic}(\text{Bun}_{G,g,n}^\delta) := \text{Pic}(\text{Bun}_{G,g,n}^\delta) / \left((\Phi_G^\delta)^*(\text{Pic}(\mathcal{M}_{g,n})) \right). \tag{3.4}$$

A first source of line bundles on $\text{Bun}_{G,g,n}$ comes from the determinant of cohomology $d_\pi(-)$ and the Deligne pairing $\langle -, - \rangle_\pi$ of line bundles on the universal curve $\pi : \mathcal{C}_{G,g,n} \rightarrow \text{Bun}_{G,g,n}$ (see [1, Chapter XIII, Sections 4, 5] for the definition and main properties of $d_\pi(-)$ and $\langle -, - \rangle_\pi$). To be more precise, any character $\chi : G \rightarrow \mathbb{G}_m \in \Lambda^*(G) := \text{Hom}(G, \mathbb{G}_m)$ gives rise to a morphism of stacks

$$\chi_\# : \text{Bun}_{G,g,n} \rightarrow \text{Bun}_{\mathbb{G}_m,g,n}$$

and, by pulling back via $\chi_\#$ the universal \mathbb{G}_m -bundle (i.e., line bundle) on the universal curve over $\text{Bun}_{\mathbb{G}_m,g,n}$, we get a line bundle \mathcal{L}_χ on $\mathcal{C}_{G,g,n}$. Then, using these line bundles \mathcal{L}_χ and the sections $\sigma_1, \dots, \sigma_n$ of π , we define the following two types of line bundles, that we call *tautological line bundles*, on $\text{Bun}_{G,g,n}$ (and hence, by restriction, also on $\text{Bun}_{G,g,n}^\delta$):

- $\mathcal{L}(\chi, \zeta) := d_\pi(\mathcal{L}_\chi(\zeta_1 \cdot \sigma_1 + \dots + \zeta_n \cdot \sigma_n))$;
- $\langle (\chi, \zeta), (\chi', \zeta') \rangle := \langle \mathcal{L}_\chi(\zeta_1 \cdot \sigma_1 + \dots + \zeta_n \cdot \sigma_n), \mathcal{L}_{\chi'}(\zeta'_1 \cdot \sigma_1 + \dots + \zeta'_n \cdot \sigma_n) \rangle_\pi$;

for $\chi, \chi' \in \text{Hom}(G, \mathbb{G}_m)$ and $\zeta = (\zeta_1, \dots, \zeta_n), \zeta' = (\zeta'_1, \dots, \zeta'_n) \in \mathbb{Z}^n$. From the standard relations between the Deligne pairing and the determinant of cohomology, we deduce that

$$\langle -, - \rangle \text{ is bilinear on } \text{Hom}(G, \mathbb{G}_m) \times \mathbb{Z}^n, \tag{3.5}$$

$$\langle (\chi, \zeta), (\chi', \zeta') \rangle = \mathcal{L}(\chi + \chi', \zeta + \zeta') \otimes \mathcal{L}(\chi, \zeta)^{-1} \otimes \mathcal{L}(\chi', \zeta')^{-1} \otimes \mathcal{L}(0, 0).$$

See [14, Section 3.5] for more details.

In the case of a torus $G = T$, the relative Picard group $\text{RPic}(\text{Bun}_{T,g,n}^\delta)$ is generated by tautological line bundles and the following theorem also clarifies the dependence relations among the tautological line bundles.

Theorem 3.2 ([14, Theorem B]). *Assume that $g \geq 1$. Let T be an algebraic torus and let $d \in \pi_1(T)$. The relative Picard group $\text{RPic}(\text{Bun}_{T,g,n}^d)$ is a free Abelian group of finite rank generated by the tautological line bundles and sitting in the exact sequences*

$$\text{Sym}^2(\Lambda^*(T)) \oplus (\Lambda^*(T) \otimes \mathbb{Z}^n) \xrightarrow{\tau_T^d + \sigma_T^d} \text{RPic}(\text{Bun}_{T,g,n}^d) \xrightarrow{\rho_T^d} \Lambda^*(T) \quad \text{if } g \geq 2, \tag{3.6}$$

$$\text{Sym}^2(\Lambda^*(T)) \oplus (\Lambda^*(T) \otimes \mathbb{Z}^n) \xrightarrow{\tau_T^d + \sigma_T^d} \text{RPic}(\text{Bun}_{T,1,n}^d) \xrightarrow{\rho_T^d} \frac{\Lambda^*(T)}{2\Lambda^*(T)} \quad \text{if } g = 1, \tag{3.7}$$

where τ_T^d (called transgression map) and σ_T^d are defined by

$$\begin{aligned} \tau_T^d(\chi \cdot \chi') &= \langle (\chi, 0), (\chi', 0) \rangle, \quad \text{for any } \chi, \chi' \in \Lambda^*(T), \\ \sigma_T^d(\chi \otimes \zeta) &= \langle (\chi, 0), (0, \zeta) \rangle, \quad \text{for any } \chi \in \Lambda^*(T) \text{ and } \zeta \in \mathbb{Z}^n, \end{aligned}$$

and ρ_T^d is the unique homomorphism such that

$$\rho_T^d(\mathcal{L}(\chi, \zeta)) = \begin{cases} \chi \in \Lambda^*(T) & \text{if } g \geq 2 \\ [\chi] \in \frac{\Lambda^*(T)}{2\Lambda^*(T)} & \text{if } g = 1 \end{cases} \quad \text{for any } \chi \in \Lambda^*(T) \text{ and } \zeta \in \mathbb{Z}^n.$$

Furthermore, the exact sequences (3.6) and (3.7) are contravariant with respect to homomorphisms of tori.

Now consider the case of an arbitrary reductive group G . Note that any character of G factors through its maximal Abelian quotient $\text{ab} : G \rightarrow G^{\text{ab}}$, i.e., the quotient of G by its derived subgroup. Hence, the tautological line bundles on $\text{Bun}_{G,g,n}^\delta$ are all pull-backs of line bundles via the morphism (induced by ab)

$$\text{ab}_\# : \text{Bun}_{G,g,n}^\delta \rightarrow \text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}$$

where $\delta^{\text{ab}} := \pi_1(\text{ab})(\delta) \in \pi_1(G^{\text{ab}})$. Moreover, Theorem 3.2 implies that the subgroup of $\text{RPic}(\text{Bun}_{G,g,n}^\delta)$ generated by the tautological line bundles coincides with the pull-back of $\text{RPic}(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}})$ via $\text{ab}_\#$.

The next result says that, for an arbitrary reductive group G , the relative Picard group of $\text{Bun}_{G,g,n}^\delta$ is generated by the image of the pull-back $\text{ab}_\#^*$ together with the image of a functorial transgression map τ_G^δ (which coincides with the transgression map τ_T^d in Theorem 3.2 if $G = T$ is a torus).

Theorem 3.3 ([14, Theorem C]). *Assume that $g \geq 1$. Let G be a reductive group and let $\text{ab} : G \rightarrow G^{\text{ab}}$ be its maximal Abelian quotient. Choose a maximal torus $\iota : T_G \hookrightarrow G$ and let \mathscr{W}_G be the Weyl group of G . Fix $\delta \in \pi_1(G)$ and denote by δ^{ab} its image in $\pi_1(G^{\text{ab}})$.*

- (1) *There exists a unique injective homomorphism (called transgression map for G)¹*

$$\tau_G^\delta : \text{Sym}^2(\Lambda^*(T_G))^{\mathscr{W}_G} \hookrightarrow \text{RPic}\left(\text{Bun}_{G,g,n}^\delta\right), \tag{3.8}$$

such that, for any lift $d \in \pi_1(T_G)$ of $\delta \in \pi_1(G)$, the composition of τ_G^δ with

$$\iota_\#^* : \text{RPic}\left(\text{Bun}_{G,g,n}^\delta\right) \rightarrow \text{RPic}\left(\text{Bun}_{T_G,g,n}^d\right)$$

is equal to the \mathscr{W}_G -invariant part of the transgression homomorphism $\tau_{T_G}^d$ defined in Theorem 3.2;

- (2) *There is a push-out diagram of injective homomorphisms of Abelian groups*

$$\begin{array}{ccc} \text{Sym}^2(\Lambda^*(G^{\text{ab}})) & \xrightarrow{\text{Sym}^2 \Lambda_{\text{ab}}^*} & \text{Sym}^2(\Lambda^*(T_G))^{\mathscr{W}_G} \\ \downarrow \tau_{G^{\text{ab}}}^{\delta^{\text{ab}}} & & \downarrow \tau_G^\delta \\ \text{RPic}\left(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}\right) & \xrightarrow{\text{ab}_\#^*} & \text{RPic}\left(\text{Bun}_{G,g,n}^\delta\right) \end{array} \tag{3.9}$$

where $\text{Sym}^2 \Lambda_{\text{ab}}^$ is the homomorphism induced by the morphism of tori $T_G \xrightarrow{\iota} G \xrightarrow{\text{ab}} G^{\text{ab}}$.*

Furthermore, the transgression homomorphism (3.8) and the diagram (3.9) are contravariant with respect to homomorphisms of reductive groups $\phi : H \rightarrow G$ such that $\phi(T_H) \subseteq T_G$.

Corollary 3.4. *With the notation of Theorem 3.3, there is an exact sequence of lattices*

$$\text{RPic}\left(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}\right) \xrightarrow{\text{ab}_\#^*} \text{RPic}\left(\text{Bun}_{G,g,n}^\delta\right) \xrightarrow{\theta_G^\delta} \text{Bil}^{s,\text{ev}}\left(\Lambda(T_{\mathscr{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}})\right)^{\mathscr{W}_G}, \tag{3.10}$$

such that $\theta_G^\delta \circ \tau_G^\delta$ is equal to the restriction homomorphism

$$\text{res}_{\mathscr{D}} : \text{Sym}^2 \Lambda^*(T_G)^{\mathscr{W}_G} \cong \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathscr{W}_G} \twoheadrightarrow \text{Bil}^{s,\text{ev}}\left(\Lambda(T_{\mathscr{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}})\right)^{\mathscr{W}_G}.$$

Furthermore, the above exact sequence (3.10) is contravariant with respect to homomorphisms of reductive groups $\phi : H \rightarrow G$ such that $\phi(T_H) \subseteq T_G$.

¹ This is the algebraic analogue of the topological transgression map $H^4(BG, \mathbb{Z}) \rightarrow H^2(\text{Bun}_{G,g,n}^\delta, \mathbb{Z})$, see [27, Section 1].

Proof. This follows from the push-out diagram (3.9) together with Proposition 2.4. \square

For an analogue of the above exact sequence for $g = 0$, see (4.16).

3.2. Two alternative presentations of $\mathrm{RPic}(\mathrm{Bun}_{G,g,n}^\delta)$

The aim of this subsection is to give two alternative presentations of $\mathrm{RPic}(\mathrm{Bun}_{G,g,n}^\delta)$, for G a reductive group and $g \geq 1$. We will freely use the notation from Subsections 2.1 and 2.2 with respect to a fixed maximal torus $\iota : T_G \hookrightarrow G$.

The first presentation of $\mathrm{RPic}(\mathrm{Bun}_{G,g,n}^\delta)$ is based on the following homomorphism.

Definition/Lemma 3.5. *Assume that $g \geq 1$. Let G be a reductive group with maximal torus T_G and Weyl group \mathscr{W}_G , and fix $\delta \in \pi_1(G)$. There exists a well-defined homomorphism*

$$\gamma_G^\delta : \mathrm{RPic}(\mathrm{Bun}_{G,g,n}^\delta) \rightarrow \mathrm{Bil}^{s, \mathscr{D}\text{-ev}}(\Lambda(T_G))^{\mathscr{W}_G}$$

uniquely determined by the following properties:

- (i) *The composition $\gamma_G^\delta \circ \tau_G^\delta$ is equal to*

$$\alpha : \mathrm{Sym}^2 \Lambda^*(T_G)^{\mathscr{W}_G} \cong \mathrm{Bil}^{s, \mathrm{ev}}(\Lambda(T_G))^{\mathscr{W}_G} \hookrightarrow \mathrm{Bil}^{s, \mathscr{D}\text{-ev}}(\Lambda(T_G))^{\mathscr{W}_G},$$

where the first isomorphism follows from (2.20), and the second injective homomorphism is the obvious inclusion;

- (ii) *The composition $\gamma_G^\delta \circ \mathrm{ab}_\#^*$ is equal to the composition*

$$\begin{aligned} \mathrm{RPic}(\mathrm{Bun}_{G^{\mathrm{ab}},g,n}^{\delta^{\mathrm{ab}}}) &\xrightarrow{\gamma_{G^{\mathrm{ab}}}^{\delta^{\mathrm{ab}}}} (\Lambda^*(G^{\mathrm{ab}}) \otimes \Lambda^*(G^{\mathrm{ab}}))^s \cong \mathrm{Bil}^s(\Lambda(G^{\mathrm{ab}})) \\ &\xrightarrow{B_{\mathrm{ab}}^*} \mathrm{Bil}^{s, \mathscr{D}\text{-ev}}(\Lambda(T_G))^{\mathscr{W}_G}, \end{aligned}$$

where $\gamma_{G^{\mathrm{ab}}}^{\delta^{\mathrm{ab}}}$ is the unique homomorphism such that

$$\begin{aligned} \gamma_{G^{\mathrm{ab}}}^{\delta^{\mathrm{ab}}}(\mathscr{L}(\chi, \zeta)) &= \chi \otimes \chi, \\ \gamma_{G^{\mathrm{ab}}}^{\delta^{\mathrm{ab}}}(\langle (\chi, \zeta), (\chi', \zeta') \rangle) &= \chi \otimes \chi' + \chi' \otimes \chi, \end{aligned}$$

the second isomorphism follows from (2.20) and B_{ab}^* is the homomorphism defined in Corollary 2.6.

Moreover, the homomorphism γ_G^δ is contravariant with respect to homomorphisms of reductive groups $\phi : H \rightarrow G$ such that $\phi(T_H) \subseteq T_G$.

Proof. The fact that $\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}$ is a well-defined homomorphism has been shown in [14, Proposition 4.1.2, Equation (4.1.3)].

In order to show that there exists a unique homomorphism satisfying (i) and (ii), using that $\text{RPic}(\text{Bun}_{G,g,n}^{\delta})$ is the pushout (3.9), it is enough to show that

$$B_{\text{ab}}^* \circ \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}} \circ \tau_{G^{\text{ab}}}^{\delta^{\text{ab}}} = \alpha \circ \text{Sym}^2 \Lambda_{\text{ab}}^* : \text{Sym}^2 \Lambda^*(G^{\text{ab}}) \rightarrow \text{Bil}^{s, \mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathcal{W}_G}. \quad (3.11)$$

Given $\chi, \chi' \in \Lambda^*(G^{\text{ab}})$ and $x, y \in \Lambda(T_G)$, we compute using the isomorphisms (2.20):

$$\begin{aligned} & \left(B_{\text{ab}}^* \circ \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}} \circ \tau_{G^{\text{ab}}}^{\delta^{\text{ab}}} \right) (\chi \cdot \chi')(x \otimes y) \\ &= \left(B_{\text{ab}}^* \circ \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}} \right) ((\chi, 0), (\chi', 0))(x \otimes y) \\ &= B_{\text{ab}}^*(\chi \otimes \chi' + \chi' \otimes \chi)(x \otimes y) \\ &= (\Lambda_{\text{ab}}^*(\chi) \otimes \Lambda_{\text{ab}}^*(\chi') + \Lambda_{\text{ab}}^*(\chi') \otimes \Lambda_{\text{ab}}^*(\chi))(x \otimes y) \\ &= \Lambda_{\text{ab}}^*(\chi)(x)\Lambda_{\text{ab}}^*(\chi')(y) + \Lambda_{\text{ab}}^*(\chi')(x)\Lambda_{\text{ab}}^*(\chi)(y). \end{aligned} \quad (3.12)$$

On the other hand,

$$\begin{aligned} & (\alpha \circ \text{Sym}^2 \Lambda_{\text{ab}}^*) (\chi \cdot \chi')(x \otimes y) \\ &= \alpha (\Lambda_{\text{ab}}^*(\chi) \cdot \Lambda_{\text{ab}}^*(\chi')) (x \otimes y) \\ &= \Lambda_{\text{ab}}^*(\chi)(x)\Lambda_{\text{ab}}^*(\chi')(y) + \Lambda_{\text{ab}}^*(\chi')(x)\Lambda_{\text{ab}}^*(\chi)(y). \end{aligned} \quad (3.13)$$

Hence, we conclude that the equality (3.11) holds and we are done.

Finally, the (contravariant) functoriality of γ_G^{δ} follows from the functoriality of B_{ab}^* (see Corollary 2.6) and of α and $\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}$ (which are obvious). \square

Using the homomorphism γ_G^{δ} , we get the required new presentation of $\text{RPic}(\text{Bun}_{G,g,n}^{\delta})$.

Theorem 3.6. *Assume that $g \geq 1$. Let G be a reductive group with maximal torus T_G and Weyl group \mathcal{W}_G , and fix $\delta \in \pi_1(G)$. Consider the following group:*

$$\widehat{H}_{g,n} := \begin{cases} \mathbb{Z} \oplus \mathbb{Z}^n & \text{if } g \geq 2 \\ \mathbb{Z}^n & \text{if } g = 1. \end{cases}$$

There is an exact sequence

$$0 \rightarrow \Lambda^*(G^{\text{ab}}) \otimes \widehat{H}_{g,n} \xrightarrow{i_G^{\delta}} \text{RPic}(\text{Bun}_{G,g,n}^{\delta}) \xrightarrow{\gamma_G^{\delta}} \text{Bil}^{s, \mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathcal{W}_G} \rightarrow 0, \quad (3.14)$$

where the morphism i_G^δ is defined as

$$\begin{aligned}
 i_G^\delta(\chi \otimes (m, \zeta)) &= \text{ab}_\#^* \left(\left\langle \mathcal{L}_\chi, \omega_\pi^m \left(\sum_{i=1}^n \zeta_i \sigma_i \right) \right\rangle \right) \quad \text{if } g \geq 2, \\
 i_G^\delta(\chi \otimes \zeta) &= \text{ab}_\#^* \left(\left\langle \mathcal{L}_\chi, \mathcal{O} \left(\sum_{i=1}^n \zeta_i \sigma_i \right) \right\rangle \right) \quad \text{if } g = 1.
 \end{aligned}
 \tag{3.15}$$

Moreover, the exact sequence (3.14) is contravariant with respect to homomorphisms of reductive groups $\phi : H \rightarrow G$ such that $\phi(T_H) \subseteq T_G$.

Proof. Consider the following diagram:

$$\begin{array}{ccccc}
 \text{RPic}(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}) & \xrightarrow{\text{ab}_\#^*} & \text{RPic}(\text{Bun}_{G,g,n}^\delta) & \xrightarrow{\theta_G^\delta} & \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}G} \\
 \downarrow \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}} & & \downarrow \gamma_G^\delta & & \parallel \\
 \text{Bil}^s(\Lambda(G^{\text{ab}})) & \xrightarrow{B_{\text{ab}}^*} & \text{Bil}^{s,\mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathcal{W}G} & \xrightarrow{\text{res}_{\mathcal{D}}} & \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}G}.
 \end{array}
 \tag{3.16}$$

Claim: The diagram (3.16) is commutative with exact rows.

Indeed, the first row is exact by Corollary 3.4 while the second row is exact by Corollary 2.6. The commutativity of the left square follows from Definition/Lemma 3.5(ii). In order to prove the commutativity of the right square, using that $\text{RPic}(\text{Bun}_{G,g,n}^\delta)$ is the pushout (3.9), it is enough to show that

$$\theta_G^\delta \circ \text{ab}_\#^* = \text{res}_{\mathcal{D}} \circ \gamma_G^\delta \circ \text{ab}_\#^*, \tag{3.17}$$

$$\theta_G^\delta \circ \tau_G^\delta = \text{res}_{\mathcal{D}} \circ \gamma_G^\delta \circ \tau_G^\delta. \tag{3.18}$$

Equality (3.17) holds since, by what observed above, we have that

$$\theta_G^\delta \circ \text{ab}_\#^* = 0 \quad \text{and} \quad \text{res}_{\mathcal{D}} \circ \gamma_G^\delta \circ \text{ab}_\#^* = \text{res}_{\mathcal{D}} \circ B_{\text{ab}}^* \circ \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}} = 0.$$

Equality (3.18) holds since, by Corollary 3.4, $\theta_G^\delta \circ \tau_G^\delta$ is equal to the restriction homomorphism

$$\text{Sym}^2 \Lambda^*(T_G)^{\mathcal{W}G} \cong \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathcal{W}G} \rightarrow \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}G},$$

which, by Corollary 2.6 and Definition/Lemma 3.5(i), is equal to $\text{res}_{\mathcal{D}} \circ \gamma_G^\delta \circ \tau_G^\delta$.

By the above claim, we can apply the snake lemma to (3.16) and we obtain the two isomorphisms

$$\begin{aligned}
 \text{ab}_\#^* : \ker(\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}) &\xrightarrow{\cong} \ker(\gamma_G^\delta), \\
 B_{\text{ab}}^* : \text{coker}(\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}) &\xrightarrow{\cong} \text{coker}(\gamma_G^\delta).
 \end{aligned}
 \tag{3.19}$$

From the definition of $\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}$ (see Definition/Lemma 3.5(ii)) together with (2.22), it follows that $\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}$ is surjective. Therefore, the second isomorphism in (3.19) implies that also γ_G^δ is surjective.

It remains to prove that the kernel morphism of γ_G^δ is equal to i_G^δ . Using the first isomorphism in (3.19) and the fact that $i_G^\delta = \text{ab}_\#^* \circ i_{G^{\text{ab}}}^{\delta^{\text{ab}}}$ by definition, it is enough to prove that

$$\text{the kernel morphism of } \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}} \text{ is equal to } i_{G^{\text{ab}}}^{\delta^{\text{ab}}}. \tag{3.20}$$

With the aim of proving (3.20), let us recall some results from [14]. Fix an isomorphism $G^{\text{ab}} \cong \mathbb{G}_m^r$ which induces an isomorphism $\Lambda^*(G^{\text{ab}}) \cong \Lambda^*(\mathbb{G}_m^r) = \mathbb{Z}^r$. Denote by $\{e_i\}_{i=1}^r$ the canonical basis of \mathbb{Z}^r and by $\{f_j\}_{j=1}^n$ the canonical basis of \mathbb{Z}^n . By [14, Theorem 4.0.1(2)], the relative Picard group of $\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}$ is freely generated by

$$\begin{aligned} &\langle (e_i, 0), (0, f_j) \rangle && \text{for } i = 1, \dots, r \text{ and } j = 1, \dots, n \\ &\langle (e_i, 0), (e_k, 0) \rangle = \langle \mathcal{L}_{e_i}, \mathcal{L}_{e_k} \rangle && \text{for } \begin{cases} 1 \leq i \leq k \leq r & \text{if } g \geq 2 \\ 1 \leq i < k \leq r & \text{if } g = 1 \end{cases} \\ &\mathcal{L}(e_i, 0) = d_\pi(\mathcal{L}_{e_i}) && \text{for } i = 1, \dots, r. \end{aligned}$$

Take now an element $\mathcal{M} \in \text{RPic}(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}})$ and write it as

$$\mathcal{M} = \sum_{1 \leq i \leq j \leq r} a_{ij} \langle (e_i, 0), (e_j, 0) \rangle + \sum_{1 \leq k \leq r} \langle (e_k, 0), (0, \zeta^k) \rangle + \sum_{1 \leq l \leq r} b_l \mathcal{L}(e_l, 0),$$

for some unique $a_{ij}, b_l \in \mathbb{Z}$, $\zeta^k = (\zeta_1^k, \dots, \zeta_n^k) \in \mathbb{Z}^n$, with the property that $a_{ii} = 0$ if $g = 1$. From the definition of $\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}$ (see Definition/Lemma 3.5(ii)), we compute

$$\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}(\mathcal{M}) = \sum_{1 \leq i \leq j \leq r} a_{ij} (e_i \otimes e_j + e_j \otimes e_i) + \sum_{1 \leq l \leq r} b_l e_l \otimes e_l.$$

Hence, we have that

$$\mathcal{M} \in \ker \left(\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}} \right) \Leftrightarrow \begin{cases} a_{ij} = 0 \text{ for } i < j \\ 2a_{ii} + b_i = 0 \end{cases} \Leftrightarrow \begin{cases} a_{ij} = 0 \text{ for } i < j \\ b_i = -2a_{ii} \\ (a_{ii}, \zeta^i) \in \widehat{H}_{g,n}. \end{cases}$$

In other words, \mathcal{M} belongs to the kernel of $\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}$ if and only if \mathcal{M} has the following form

$$\begin{aligned} \mathcal{M} &= \sum_{\substack{1 \leq i \leq r \\ (a_{ii}, \zeta^i) \in \widehat{H}_{g,n}}} \left[a_{ii} \langle \mathcal{L}_{e_i}, \mathcal{L}_{e_i} \rangle - 2a_{ii} d_\pi(\mathcal{L}_{e_i}) + \left\langle \mathcal{L}_{e_i}, \mathcal{O} \left(\sum_k \zeta_k^i \sigma_k \right) \right\rangle \right] \\ &= \sum_{\substack{1 \leq i \leq r \\ (a_{ii}, \zeta^i) \in \widehat{H}_{g,n}}} \left\langle \mathcal{L}_{e_i}, \omega_\pi^{a_{ii}} \left(\sum_k \zeta_k^i \sigma_k \right) \right\rangle, \end{aligned}$$

where the second equality follows from [14, Remark 3.5.1]. This shows that $i_{G^{\text{ab}}}^{\delta^{\text{ab}}}$ is an injective homomorphism whose image is equal to the kernel of $\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}$, which proves (3.20). \square

We now want to get a second presentation of $\text{RPic}(\text{Bun}_{G,g,n}^{\delta})$. With this aim, we introduce the following homomorphism.

Definition/Lemma 3.7. *Assume that $g \geq 1$. Let G be a reductive group with maximal torus T_G and Weyl group \mathscr{W}_G , and fix $\delta \in \pi_1(G)$. There exists a well-defined (non functorial) homomorphism*

$$\omega_G^{\delta} : \text{RPic}(\text{Bun}_{G,g,n}^{\delta}) \rightarrow \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})}$$

uniquely determined by the following properties:

- (i) *The composition $\omega_G^{\delta} \circ \tau_G^{\delta}$ is equal to the composition*

$$\begin{aligned} \text{Sym}^2 \Lambda^*(T_G)^{\mathscr{W}_G} &\xrightarrow{\cong} \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathscr{W}_G} \subseteq \text{Bil}^{s,\mathscr{Q}-\text{ev}}(\Lambda(T_G))^{\mathscr{W}_G} \\ &\xrightarrow{\text{ev}_G^{\delta}} \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} \end{aligned}$$

where the first isomorphism is induced by (2.20) and ev_G^{δ} is the homomorphism in Definition/Lemma 2.8;

- (ii) *The composition $\omega_G^{\delta} \circ \text{ab}_{\#}^{\delta}$ is equal to the composition*

$$\begin{aligned} \text{RPic}(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}) &\xrightarrow{\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}}} \Lambda^*(G^{\text{ab}}) \xrightarrow{\overline{\Lambda_{\text{ab}}^*}} \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})}, \\ \mathcal{L}(\chi, \zeta) &\mapsto [\chi(\delta^{\text{ab}}) + |\zeta| + 1 - g] \chi, \\ \langle (\chi, \zeta), (\chi', \zeta') \rangle &\mapsto [\chi'(\delta^{\text{ab}}) + |\zeta'|] \chi + [\chi(\delta^{\text{ab}}) + |\zeta|] \chi', \end{aligned}$$

where $|\zeta| = \sum_i \zeta_i \in \mathbb{Z}$ and similarly for $|\zeta'|$, and $\overline{\Lambda_{\text{ab}}^*}$ is the homomorphism in (2.17).

Proof. The fact that $\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}}$ is well-defined has been proved in [14, Proposition 4.1.2(i)].

In order to show that there exists a unique homomorphism ω_G^{δ} satisfying properties (i) and (ii), using that $\text{RPic}(\text{Bun}_{G,g,n}^{\delta})$ is the pushout (3.9), it is enough to show that

$$\overline{\Lambda_{\text{ab}}^*} \circ \omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} \circ \tau_{G^{\text{ab}}}^{\delta^{\text{ab}}} = \text{ev}_G^{\delta} \circ \text{Sym}^2 \Lambda_{\text{ab}}^* : \text{Sym}^2 \Lambda^*(G^{\text{ab}}) \rightarrow \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})}. \quad (3.21)$$

Given $\chi, \chi' \in \Lambda^*(G^{\text{ab}})$, we compute

$$\begin{aligned} \left(\overline{\Lambda_{\text{ab}}^*} \circ \omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} \circ \tau_{G^{\text{ab}}}^{\delta^{\text{ab}}}\right) (\chi \cdot \chi') &= \left(\overline{\Lambda_{\text{ab}}^*} \circ \omega_{G^{\text{ab}}}^{\delta^{\text{ab}}}\right) ((\chi, 0), (\chi', 0)) \\ &= \overline{\Lambda_{\text{ab}}^*} (\chi(\delta^{\text{ab}})\chi' + \chi'(\delta^{\text{ab}})\chi) \\ &= \chi(\delta^{\text{ab}})\Lambda_{\text{ab}}^*(\chi') + \chi'(\delta^{\text{ab}})\Lambda_{\text{ab}}^*(\chi), \end{aligned} \tag{3.22}$$

$$\begin{aligned} \left(\text{ev}_G^\delta \circ \text{Sym}^2 \Lambda_{\text{ab}}^*\right) (\chi \cdot \chi') &= \text{ev}_G^\delta (\Lambda_{\text{ab}}^*(\chi) \cdot \Lambda_{\text{ab}}^*(\chi')) \\ &= \Lambda_{\text{ab}}^*(\chi)(d)\Lambda_{\text{ab}}^*(\chi') + \Lambda_{\text{ab}}^*(\chi')(d)\Lambda_{\text{ab}}^*(\chi) \\ &= \chi(\Lambda_{\text{ab}}(d))\Lambda_{\text{ab}}^*(\chi') + \chi'(\Lambda_{\text{ab}}(d))\Lambda_{\text{ab}}^*(\chi). \end{aligned} \tag{3.23}$$

The expressions (3.22) and (3.23) coincide since $\Lambda_{\text{ab}}(d) = \pi_1(\text{ab})(\delta) = \delta^{\text{ab}}$ for any lift $d \in \Lambda(T_G)$ of $\delta \in \pi_1(G)$; hence, the equality (3.21) holds and we are done. \square

Remark 3.8. We remark that the homomorphism ω_G^δ is not equal to the composition

$$\text{RPic} \left(\text{Bun}_{G,g,n}^\delta \right) \xrightarrow{\gamma_G^\delta} \text{Bil}^{s, \mathcal{D}-\text{ev}} (\Lambda(T_G))^{\mathscr{W}_G} \xrightarrow{\text{ev}_G^\delta} \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})},$$

where γ_G^δ is the homomorphism in Definition/Lemma 3.5 and τ_G^δ is the homomorphism in Definition/Lemma 2.8. More precisely, their compositions with the pull-back $\text{ab}_\#^*$ are different.

By putting together the homomorphisms of Definition/Lemmas 3.5 and 3.7, we get the following homomorphism:

$$\omega_G^\delta \oplus \gamma_G^\delta : \text{RPic} \left(\text{Bun}_{G,g,n}^\delta \right) \rightarrow \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} \oplus \text{Bil}^{s, \mathcal{D}-\text{ev}} (\Lambda(T_G))^{\mathscr{W}_G}.$$

With the aim of describing its image, we give the following:

Definition 3.9. Let G be a reductive group with maximal torus T_G and Weyl group \mathscr{W}_G , and fix $\delta \in \pi_1(G)$. Denote by

$$\text{NS} \left(\text{Bun}_{G,g,n}^\delta \right) \subseteq \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} \oplus \text{Bil}^{s, \mathcal{D}-\text{ev}} (\Lambda(T_G))^{\mathscr{W}_G}$$

the subgroup consisting of all the elements $([\chi], b)$ such that

$$\begin{aligned} [\chi|_{\Lambda(T_{\mathcal{D}(G)})}] &= \overline{\Lambda_{\mathcal{D}}^*}([\chi]) \text{ is equal to } \left(\overline{\Lambda_{\mathcal{D}}^*} \circ \text{ev}_G^\delta\right) (b) \\ &= b(\delta \otimes -)|_{\Lambda(T_{\mathcal{D}(G)})} := [b(d \otimes -)|_{\Lambda(T_{\mathcal{D}(G)})}] \end{aligned} \tag{3.24}$$

as elements in $\frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}})}$, where $d \in \Lambda(T_G)$ is any lift of δ , and $\overline{\Lambda_{\mathcal{D}}^*}$ and ev_G^δ are the homomorphisms of Definition/Lemma 2.8.

The group $\text{NS}(\text{Bun}_{G,g,n}^\delta)$ is contravariant with respect to homomorphisms of reductive groups $\phi : H \rightarrow G$ such that $\phi(T_H) \subseteq T_G$.

Definition/Lemma 3.10. *Let $\phi : H \rightarrow G$ be a homomorphism of reductive groups, and choose maximal tori $T_G \subseteq G$ and $T_H \subseteq H$ in such a way that $\phi(T_H) \subseteq T_G$. Let $\epsilon \in \pi_1(H)$ and set $\delta := \pi_1(\phi)(\epsilon) \in \pi_1(G)$. Pick a lift $e \in \Lambda(T_H)$ of $\epsilon \in \pi_1(H)$. Then there exists a well-defined homomorphism*

$$\begin{aligned} \phi^{*,\text{NS}} : \text{NS}(\text{Bun}_{G,g,n}^\delta) &\longrightarrow \text{NS}(\text{Bun}_{H,g,n}^\epsilon), \\ ([\chi], b) &\mapsto \left(\left[\Lambda_\phi^*(\chi^{b(\Lambda_\phi(e) \otimes -)}) \right], B_\phi^*(b) \right), \end{aligned} \tag{3.25}$$

where $\Lambda_\phi : \Lambda(T_H) \rightarrow \Lambda(T_G)$, $\Lambda_\phi^* : \Lambda^*(T_G) \rightarrow \Lambda^*(T_H)$ and $B_\phi^* : \text{Bil}^s(\Lambda(T_G)) \rightarrow \text{Bil}^s(\Lambda(T_H))$ are the natural morphisms induced by $\phi : T_H \rightarrow T_G$, and $\chi^{b(\Lambda_\phi(e) \otimes -)} \in \Lambda^*(T_G)$ is the unique lift of $[\chi] \in \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G\text{ad}})}$ such that

$$\left(\chi^{b(\Lambda_\phi(e) \otimes -)} \right)_{|\Lambda(T_{\mathcal{D}(G)})} = b(\Lambda_\phi(e) \otimes -)_{|\Lambda(T_{\mathcal{D}(G)})}.$$

Moreover, if $\psi : L \rightarrow H$ is another homomorphism of reductive groups and we choose a maximal torus $T_L \subseteq L$ in such a way that $\psi(T_L) \subseteq T_H$, then $(\phi \circ \psi)^* = \psi^* \circ \phi^*$.

Proof. Let us first consider the following two special cases:

Special case I: $\phi : T' \rightarrow T$ is a morphism of tori.

Choose $d' \in \Lambda(T')$ and set $d := \Lambda_\phi(d') \in \Lambda(T)$. The definition (3.25) reduces in this special case to

$$\begin{aligned} \phi^{*,\text{NS}} : \text{NS}(\text{Bun}_{T,g,n}^d) &\longrightarrow \text{NS}(\text{Bun}_{T',g,n}^{d'}), \\ (\chi, b) &\mapsto (\Lambda_\phi^*(\chi), B_\phi^*(b)), \end{aligned} \tag{3.26}$$

which is clearly a well-defined homomorphism. Moreover, the association $\phi \mapsto \phi^{*,\text{NS}}$ is compatible with the composition of morphisms of tori.

Special case II: $\phi = \iota : T_G \rightarrow G$ is the inclusion of a maximal torus inside a reductive group G .

Choose a lift $d \in \Lambda(T_G)$ of $\delta \in \pi_1(G)$. Pick an element $([\chi], b) \in \text{NS}(\text{Bun}_{G,g,n}^\delta)$. Consider the following commutative diagram with surjective arrows:

$$\begin{array}{ccc} \Lambda^*(T_G) & \xrightarrow{\Lambda_{\mathcal{D}}} & \Lambda^*(T_{\mathcal{D}(G)}) \\ \downarrow & & \downarrow \\ \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G\text{ad}})} & \xrightarrow{\overline{\Lambda_{\mathcal{D}}}} & \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G\text{ad}})}. \end{array} \tag{3.27}$$

The diagram (3.27) is a pull-back diagram since the kernels of the vertical surjections are both equal to $\Lambda^*(T_{G^{ad}})$ while the kernels of the horizontal surjections are both equal to $\Lambda^*(G^{ab})$ and $\Lambda^*(T_{G^{ad}}) \cap \Lambda^*(G^{ab}) = \{0\}$. From this and condition (3.24), it follows that there exists a unique lift of $[\chi] \in \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{ad}})}$, that we denote by $\chi^{b(d \otimes -)} \in \Lambda^*(T_G)$, with the property that

$$\left(\chi^{b(d \otimes -)}\right)_{|\Lambda(T_{\mathcal{D}(G)})} = b(d \otimes -)_{|\Lambda(T_{\mathcal{D}(G)})}. \tag{3.28}$$

Definition (3.25) reduces in this special case to

$$\begin{aligned} \iota^{*,NS} : \text{NS}\left(\text{Bun}_{G,g,n}^\delta\right) &\longrightarrow \text{NS}\left(\text{Bun}_{T_G,g,n}^d\right), \\ ([\chi], b) &\mapsto \left(\chi^{b(d \otimes -)}, b\right), \end{aligned} \tag{3.29}$$

which is a well-defined and injective homomorphism whose image is equal to

$$\begin{aligned} \text{Im}(\iota^{*,NS}) &= \{(\chi, b) : \chi_{|\Lambda(T_{\mathcal{D}(G)})} \\ &= b(d \otimes -)_{|\Lambda(T_{\mathcal{D}(G)})} \text{ and } b \in \text{Bil}^{s,\mathcal{D}\text{-ev}}(\Lambda(T_G))^{\#G}\}. \end{aligned} \tag{3.30}$$

We now go back to the general case. Denote the inclusions of the maximal tori in G and H by, respectively, $\iota_G : T_G \hookrightarrow G$ and $\iota_H : T_H \hookrightarrow H$ and set $\phi_T := \phi_{|T_H} : T_H \rightarrow T_G$. Set also $d := \Lambda_\phi(e) \in \Lambda(T_G)$, which is a lift of $\delta \in \pi_1(G)$. Consider the composition

$$\begin{aligned} \phi_T^{*,NS} \circ \iota_G^{*,NS} : \text{NS}\left(\text{Bun}_{G,g,n}^\delta\right) &\longrightarrow \text{NS}\left(\text{Bun}_{T_H,g,n}^e\right) \\ ([\chi], b) &\mapsto \left(\Lambda_\phi^*(\chi^{b(d \otimes -)}), B_\phi^*(b)\right) \end{aligned} \tag{3.31}$$

which is a well-defined homomorphism by the special cases already treated. Moreover, for any $([\chi], b) \in \text{NS}(\text{Bun}_{G,g,n}^\delta)$ and for every $x \in \Lambda(T_{\mathcal{D}(H)})$, we have that

$$\begin{aligned} \Lambda_\phi^*\left(\chi^{b(d \otimes -)}\right)(x) &= \chi^{b(d \otimes -)}(\Lambda_\phi(x)) = b(d \otimes \Lambda_\phi(x)) \\ &= b(\Lambda_\phi(e) \otimes \Lambda_{\phi_T}(x)) = \left(B_\phi^*(b)\right)(e \otimes x), \end{aligned}$$

where in the second equality we have used (3.28). This computation, together with (3.30), implies that the image of $\phi_T^{*,NS} \circ \iota_G^{*,NS}$ is contained in the image of $\iota_H^{*,NS} : \text{NS}(\text{Bun}_{H,g,n}^e) \rightarrow \text{NS}(\text{Bun}_{T_H,g,n}^e)$. Hence, we get a factorization

$$\phi_T^{*,NS} \circ \iota_G^{*,NS} = \iota_H^{*,NS} \circ \phi^{*,NS} \tag{3.32}$$

for some (unique) homomorphism $\phi^{*,NS} : \text{NS}(\text{Bun}_{G,g,n}^\delta) \rightarrow \text{NS}(\text{Bun}_{H,g,n}^e)$.

From the expression (3.31), we conclude that $\phi^{*,NS}$ is given by the formula (3.25).

Finally, the compatibility of the association $\phi \mapsto \phi^{*,NS}$ with the composition of morphisms is due to the factorization (3.32) together with the Special Case I. \square

The group $\text{NS}(\text{Bun}_{G,g,n}^\delta)$ admits a functorial two-step filtration, that we describe in the following:

Proposition 3.11. *Let G be a reductive group with maximal torus T_G and Weyl group \mathcal{W}_G , and fix $\delta \in \pi_1(G)$. We have the commutative diagram, with exact rows and columns*

$$\begin{array}{ccccc}
 \Lambda^*(G^{\text{ab}}) & \xrightarrow{\overline{\Lambda_{\text{ab}}^* \oplus 0}} & \text{NS}(\text{Bun}_{G,g,n}^\delta) & \xrightarrow{\text{res}_G^{\text{NS}}} & \text{Bil}^s(\Lambda(T_G))^{\mathcal{W}_G} \\
 \downarrow i_1 & & \parallel & & \downarrow \text{res}_{\mathcal{D}} \\
 \Lambda^*(G^{\text{ab}}) \oplus \text{Bil}^s(\Lambda(G^{\text{ab}})) & \xrightarrow{\text{ab}^{*,\text{NS}}} & \text{NS}(\text{Bun}_{G,g,n}^\delta) & \xrightarrow{\text{res}_G^{\text{NS}}} & \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_G^{\text{a}}))^{\mathcal{W}_G} \\
 \downarrow p_2 & & & & \downarrow \text{res}_{\mathcal{D}} \\
 \text{Bil}^s(\Lambda(G^{\text{ab}})) & & & &
 \end{array} \tag{3.33}$$

where the identification $\text{NS}(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}) = \Lambda^*(G^{\text{ab}}) \oplus \text{Bil}^s(\Lambda(G^{\text{ab}}))$ follows from Definition 3.9, i_1 is the inclusion of the first factor and p_2 is the projection onto the second factor, the right vertical column is (2.28), res_G^{NS} is the projection onto the second factor and $\text{res}_{\mathcal{D}}^{\text{NS}} := \text{res}_{\mathcal{D}} \circ \text{res}_G^{\text{NS}}$.

Moreover, the diagram (3.33) is contravariant with respect to homomorphisms of reductive groups $\phi : H \rightarrow G$ such that $\phi(T_H) \subseteq T_G$.

Proof. The commutativity of the right square of the diagram is clear, while the commutativity of the left square follows from the fact $\text{ab}^{*,\text{NS}} = \overline{\Lambda_{\text{ab}}^*} \oplus B_{\text{ab}}^*$, as it is easily deduced from Definition/Lemma 3.10.

The exactness of the left and central columns is clear, while the exactness of the right column follows from Corollary 2.6.

The exactness of the upper row follows from the definition of $\text{NS}(\text{Bun}_{G,g,n}^\delta)$ together with the exactness of the column in (2.17).

It remains to prove the exactness of the lower row. The injectivity of $\text{ab}^{*,\text{NS}} = \overline{\Lambda_{\text{ab}}^*} \oplus B_{\text{ab}}^*$ and the fact that $\text{res}_{\mathcal{D}}^{\text{NS}} \circ \text{ab}^{*,\text{NS}} = 0$ are obvious. The surjectivity of $\text{res}_{\mathcal{D}}^{\text{NS}}$ follows from the surjectivity of res_G^{NS} and of $\text{res}_{\mathcal{D}}$. Let us now prove that $\ker(\text{res}_{\mathcal{D}}^{\text{NS}}) \subseteq \text{Im}(\overline{\Lambda_{\text{ab}}^*} \oplus B_{\text{ab}}^*)$. Pick an element $([\chi], b) \in \text{NS}(\text{Bun}_{G,g,n}^\delta)$ such that $0 = \text{res}_{\mathcal{D}}^{\text{NS}}([\chi], b) = \text{res}_{\mathcal{D}}(b)$. By Corollary 2.6, there exists $b^{\text{ab}} \in \text{Bil}^s(\Lambda(G^{\text{ab}}))$ such that $b = B_{\text{ab}}^*(b^{\text{ab}})$. Moreover, from (3.24) it follows that $[\chi]_{\Lambda(T_{\mathcal{D}(G)})} = 0 \in \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G,\text{ad}})}$. Hence, by (2.17), there exists $\chi^{\text{ab}} \in \Lambda^*(G^{\text{ab}})$ such that $\overline{\Lambda_{\text{ab}}^*}(\chi^{\text{ab}}) = [\chi]$. Therefore, $(\chi^{\text{ab}}, b^{\text{ab}}) \in \text{NS}(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}})$ and $\overline{\Lambda_{\text{ab}}^*} \oplus B_{\text{ab}}^*((\chi^{\text{ab}}, b^{\text{ab}})) = ([\chi], b)$ and we are done.

Finally, the functoriality of the diagram (3.33) follows straightforwardly from the definition of the pull-back morphism (3.25). \square

Using the above homomorphisms ω_G^δ and γ_G^δ , we can now give the following new presentation of $\mathrm{RPic}(\mathrm{Bun}_{G,g,n}^\delta)$.

Theorem 3.12. *Assume that $g \geq 1$. Let G be a reductive group with maximal torus T_G and Weyl group \mathcal{W}_G , and fix $\delta \in \pi_1(G)$. Consider the following group:*

$$H_{g,n} := \begin{cases} \{(m, \zeta) \in \mathbb{Z} \oplus \mathbb{Z}^n : (2g - 2)m + |\zeta| = 0\} & \text{if } g \geq 2 \\ \{\zeta \in \mathbb{Z}^n : |\zeta| = 0\} & \text{if } g = 1. \end{cases}$$

(1) *There is an exact sequence*

$$0 \rightarrow \Lambda^*(G^{\mathrm{ab}}) \otimes H_{g,n} \xrightarrow{j_G^\delta} \mathrm{RPic}(\mathrm{Bun}_{G,g,n}^\delta) \xrightarrow{\omega_G^\delta \oplus \gamma_G^\delta} \mathrm{NS}(\mathrm{Bun}_{G,g,n}^\delta), \quad (3.34)$$

where the morphism j_G^δ is defined as

$$j_G^\delta(\chi \otimes (m, \zeta)) = \mathrm{ab}_\#^* \left(\left\langle \mathcal{L}_\chi, \omega_\pi^m \left(\sum_{i=1}^n \zeta_i \sigma_i \right) \right\rangle \right) \quad \text{if } g \geq 2,$$

$$j_G^\delta(\chi \otimes \zeta) = \mathrm{ab}_\#^* \left(\left\langle \mathcal{L}_\chi, \mathcal{O} \left(\sum_{i=1}^n \zeta_i \sigma_i \right) \right\rangle \right) \quad \text{if } g = 1.$$

Moreover, the exact sequence (3.34) is contravariant with respect to homomorphisms of reductive groups $\phi : H \rightarrow G$ such that $\phi(T_H) \subseteq T_G$;

(2) *The image of $\omega_G^\delta \oplus \gamma_G^\delta$ is equal to*

$$\begin{aligned} & \mathrm{Im}(\omega_G^\delta \oplus \gamma_G^\delta) \tag{3.35} \\ &= \begin{cases} \mathrm{NS}(\mathrm{Bun}_{G,g,n}^\delta) & \text{if } n \geq 1 \\ \left\{ ([\chi], b) \in \mathrm{NS}(\mathrm{Bun}_{G,g,n}^\delta) : \begin{array}{l} [\chi(x) - b(\delta \otimes x)] + (g - 1)b(x \otimes x) \\ \text{is a multiple of } 2g - 2, \text{ for any } x \in \Lambda(T_G) \end{array} \right\} & \text{if } n = 0. \end{cases} \end{aligned}$$

Moreover, if $n = 0$ then

$$\frac{\mathrm{NS}(\mathrm{Bun}_{G,g,n}^\delta)}{\mathrm{Im}(\omega_G^\delta \oplus \gamma_G^\delta)} \cong \left(\frac{\mathbb{Z}}{(2g - 2)\mathbb{Z}} \right)^{\dim G^{\mathrm{ab}}}.$$

Remark 3.13.

- (i) For $n = 0$, the subgroup on the right-hand side of (3.35) is well-defined. Indeed, by (3.24), $[\chi(x) - b(\delta \otimes x)]$ is well-defined for any $x \in \Lambda(T_G)$ and it is equal to $[\chi(x^{\mathrm{ab}}) - b(\delta \otimes x^{\mathrm{ab}})]$, where x^{ab} is the image of x in $\Lambda(G^{\mathrm{ab}})$;
- (ii) If either $n = 0$ or $g = n = 1$ then $H_{g,n} = 0$, which implies that the map $\omega_G^\delta \oplus \gamma_G^\delta$ is injective.

Proof. The theorem has been proved for a torus in [14, Proposition 4.3.1], and we are going to apply this result for G^{ab} in order to prove the case of a general reductive group G .

Let us first prove (1) by dividing the proof in two steps.

Step I: The image of

$$\omega_G^\delta \oplus \gamma_G^\delta : \text{RPic} \left(\text{Bun}_{G,g,n}^\delta \right) \rightarrow \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} \oplus \text{Bil}^s(\Lambda(T_G))^{\mathscr{W}G}$$

is contained in $\text{NS}(\text{Bun}_{G,g,n}^\delta)$.

In order to prove this, it is enough to show, using that $\text{RPic}(\text{Bun}_{G,g,n}^\delta)$ is the pushout (3.9), that

$$\begin{aligned} \left((\omega_G^\delta \oplus \gamma_G^\delta) \circ \text{ab}_\#^* \right) \left(\text{RPic} \left(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}} \right) \right) &\subseteq \text{NS} \left(\text{Bun}_{G,g,n}^\delta \right), \\ \left((\omega_G^\delta \oplus \gamma_G^\delta) \circ \tau_G^\delta \right) \left(\text{Sym}^2 \Lambda^*(T_G)^{\mathscr{W}G} \right) &\subseteq \text{NS} \left(\text{Bun}_{G,g,n}^\delta \right). \end{aligned} \tag{3.36}$$

The first containment in (3.36) follows since Definition/Lemma 3.7(ii) and Definition/Lemma 3.5(ii) imply that $(\omega_G^\delta \oplus \gamma_G^\delta) \circ \text{ab}_\#^*$ factorizes as the composition

$$\text{RPic} \left(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}} \right) \xrightarrow{\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} \oplus \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}} \Lambda^*(G^{\text{ab}}) \oplus \text{Bil}^s(\Lambda(G^{\text{ab}})) \xrightarrow{\overline{\Lambda_{\text{ab}}^*} \oplus B_{\text{ab}}^*} \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} \oplus \text{Bil}^s(\Lambda(T_G))^{\mathscr{W}G}, \tag{3.37}$$

and $\text{Im}(\overline{\Lambda_{\text{ab}}^*} \oplus B_{\text{ab}}^*) \subseteq \text{NS}(\text{Bun}_{G,g,n}^\delta)$ as observed in Proposition 3.11.

Take now an element $b \in \text{Sym}^2 \Lambda^*(T_G)^{\mathscr{W}G} \cong \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathscr{W}G}$. Using Definition/Lemma 3.7(i) and Definition/Lemma 3.5(i), we compute

$$\begin{aligned} \left((\omega_G^\delta \oplus \gamma_G^\delta) \circ \tau_G^\delta \right) (b) &= \left(\text{ev}_G^\delta(b), b \right) = (b(\delta \otimes -), b) \\ &\in \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} \oplus \text{Bil}^s(\Lambda(T_G))^{\mathscr{W}G}. \end{aligned} \tag{3.38}$$

From Definition 3.9, it follows easily that the element $(b(\delta \otimes -), b)$ belongs to $\text{NS}(\text{Bun}_{G,g,n}^\delta)$, and this proves the second containment in (3.36).

Step II: The kernel of $\omega_G^\delta \oplus \gamma_G^\delta$ is equal to

$$j_G^\delta : \Lambda^*(G^{\text{ab}}) \otimes H_{g,n} \hookrightarrow \text{RPic} \left(\text{Bun}_{G,g,n}^\delta \right).$$

Consider the diagram

$$\begin{array}{ccccc} \text{RPic} \left(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}} \right) & \xhookrightarrow{\text{ab}_\#^*} & \text{RPic} \left(\text{Bun}_{G,g,n}^\delta \right) & \xrightarrow{\theta_G^\delta} & \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathscr{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}G} \\ \downarrow \omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} \oplus \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}} & & \downarrow \omega_G^\delta \oplus \gamma_G^\delta & & \parallel \\ \text{NS}(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}) = \Lambda^*(G^{\text{ab}}) \oplus \text{Bil}^s(\Lambda(G^{\text{ab}})) & \xhookrightarrow{\overline{\Lambda_{\text{ab}}^*} \oplus B_{\text{ab}}^*} & \text{NS}(\text{Bun}_{G,g,n}^\delta) & \xrightarrow{\text{res}_{\mathscr{D}}^{\text{NS}}} & \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathscr{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}G} \end{array} \tag{3.39}$$

whose rows are exact by Corollary 3.4 and Proposition 3.11, and whose commutativity follows from (3.37) and (3.38). By applying the snake lemma to (3.39), we get that

$$\ker \left(\omega_{G^{ab}}^{\delta^{ab}} \oplus \gamma_{G^{ab}}^{\delta^{ab}} \right) \xrightarrow[\cong]{ab_{\#}^*} \ker \left(\omega_G^{\delta} \oplus \gamma_G^{\delta} \right), \tag{3.40}$$

$$\operatorname{coker} \left(\omega_{G^{ab}}^{\delta^{ab}} \oplus \gamma_{G^{ab}}^{\delta^{ab}} \right) \xrightarrow[\cong]{\overline{\Lambda_{ab}^*} \oplus B_{ab}^*} \operatorname{coker} \left(\omega_G^{\delta} \oplus \gamma_G^{\delta} \right). \tag{3.41}$$

By applying [14, Proposition 4.3.1] to G^{ab} , we get that the kernel of $\omega_{G^{ab}}^{\delta^{ab}} \oplus \gamma_{G^{ab}}^{\delta^{ab}}$ is equal to

$$j_{G^{ab}}^{\delta^{ab}} : \Lambda^*(G^{ab}) \otimes H_{g,n} \hookrightarrow \operatorname{RPic} \left(\operatorname{Bun}_{G^{ab},g,n}^{\delta^{ab}} \right).$$

By combining this with (3.40) and the fact that $j_G^{\delta} = ab_{\#}^* \circ j_{G^{ab}}^{\delta^{ab}}$, Step II follows.

Finally, the functoriality of the morphism $\omega_G^{\delta} \oplus \gamma_G^{\delta}$ (and hence of the sequence (3.34)) follows straightforwardly from (3.37) and (3.38).

Let us now prove (2). If $n > 0$ then $\omega_{G^{ab}}^{\delta^{ab}} \oplus \gamma_{G^{ab}}^{\delta^{ab}}$ is surjective by [14, Proposition 4.3.1], which, combined with (3.41), implies that $\omega_G^{\delta} \oplus \gamma_G^{\delta}$ is also surjective.

Assume now that $n = 0$ and call I_G^{δ} the subgroup of $\operatorname{NS}(\operatorname{Bun}_{G^{ab},g,n}^{\delta^{ab}})$ defined on the right-hand side of (3.35). By [14, Proposition 4.3.1], we know that $\operatorname{Im}(\omega_{G^{ab}}^{\delta^{ab}} \oplus \gamma_{G^{ab}}^{\delta^{ab}}) = I_{G^{ab}}^{\delta^{ab}}$. Using this and (3.41), in order to prove that $\operatorname{Im}(\omega_G^{\delta} \oplus \gamma_G^{\delta}) = I_G^{\delta}$, it is enough to prove that

$$\left\{ \begin{aligned} \operatorname{Im}(\omega_G^{\delta} \oplus \gamma_G^{\delta}) &\subseteq I_G^{\delta} \\ (\overline{\Lambda_{ab}^*} \oplus B_{ab}^*)^{-1}(I_G^{\delta}) &\subseteq I_{G^{ab}}^{\delta^{ab}}. \end{aligned} \right. \tag{3.42}$$

Furthermore, using that $\operatorname{RPic}(\operatorname{Bun}_{G^{ab},g,n}^{\delta^{ab}})$ is the pushout (3.9), the inclusions (3.42) are equivalent to the following two conditions:

$$\left((\omega_G^{\delta} \oplus \gamma_G^{\delta}) \circ \tau_G^{\delta} \right) \left(\operatorname{Sym}^2 \Lambda^*(T_G)^{\mathscr{W}G} \right) \subseteq I_G^{\delta}, \tag{3.43}$$

$$\left(\overline{\Lambda_{ab}^*} \oplus B_{ab}^* \right)^{-1} (I_G^{\delta}) = I_{G^{ab}}^{\delta^{ab}}. \tag{3.44}$$

Now, inclusion (3.43) follows from (3.38) together with the fact that b is even. Furthermore, if $(\chi, b) \in \operatorname{NS}(\operatorname{Bun}_{G^{ab},g,n}^{\delta^{ab}}) = \Lambda^*(G^{ab}) \oplus \operatorname{Bil}^s(\Lambda(G^{ab}))$, then we have for every $x \in \Lambda(T_G)$

$$\begin{aligned} &\overline{\Lambda_{ab}^*}(\chi)(x) - B_{ab}^*(b)(\delta, x) + (g-1)B_{ab}^*(b)(x, x) \\ &= \chi(x^{ab}) - b(\delta^{ab}, x^{ab}) + (g-1)b(x^{ab}, x^{ab}), \end{aligned}$$

where x^{ab} is the image of x into $\Lambda(G^{\text{ab}})$. Since $\Lambda(T_G)$ surjects onto $\Lambda(T_{G^{\text{ab}}})$, we deduce that

$$(\chi, b) \in I_{G^{\text{ab}}}^{\delta^{\text{ab}}} \Leftrightarrow \left(\overline{\Lambda_{\text{ab}}^*}, B_{\text{ab}}^*\right)^{-1} (\chi, b) \in I_G^{\delta},$$

which shows (3.44) and concludes the first assertion of part (2).

The last assertion follows from (3.41) and the analogous result for G^{ab} , see [14, Remark 4.3.2]. \square

4. The restriction homomorphism

The aim of this section is to describe the restriction homomorphism

$$\text{res}_G^{\delta}(C) : \text{RPic}\left(\text{Bun}_{G,g,n}^{\delta}\right) \rightarrow \text{Pic}\left(\text{Bun}_G^{\delta}(C)\right) \tag{4.1}$$

for any $(C, p_1, \dots, p_n) \in \mathcal{M}_{g,n}(k)$.

Before doing this, we need to recall the description of $\text{Pic}(\text{Bun}_G^{\delta}(C))$ obtained in [3].

Theorem 4.1 ([3, Theorem 5.3.1]). *Let C be a (irreducible, projective, smooth) curve of genus $g \geq 0$ over $k = \bar{k}$ and denote by J_C its Jacobian. Let G be a reductive group over k and fix $\delta \in \pi_1(G)$. Then there exists a (contravariantly) functorial exact sequence of Abelian groups*

$$0 \rightarrow \text{Hom}(\pi_1(G), J_C(k)) \xrightarrow{j_G^{\delta}(C)} \text{Pic}(\text{Bun}_G^{\delta}(C)) \xrightarrow{c_G^{\delta}(C)} \text{NS}\left(\text{Bun}_G^{\delta}(C)\right) \rightarrow 0, \tag{4.2}$$

where the Neron-Severi group $\text{NS}(\text{Bun}_G^{\delta}(C))$ is the group of all triples $(l_{\mathcal{R}}, b_{\mathcal{R}}, b)$ consisting of

- (i) $l_{\mathcal{R}} \in \Lambda^*(\mathcal{R}(G))$;
- (ii) $b_{\mathcal{R}} \in \text{Hom}^s(\Lambda(\mathcal{R}(G)) \otimes \Lambda(\mathcal{R}(G)), \text{End}(J_C))$, i.e., $b_{\mathcal{R}} : \Lambda(\mathcal{R}(G)) \otimes \Lambda(\mathcal{R}(G)) \rightarrow \text{End}(J_C)$ with the property that $b_{\mathcal{R}}(x_1 \otimes x_2)^{\dagger} = b_{\mathcal{R}}(x_2 \otimes x_1)$ where $\dagger : \text{End}(J_C) \rightarrow \text{End}(J_C)$ is the Rosati involution associated to the canonical polarization on J_C ;
- (iii) $b \in \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathcal{W}G} \cong \text{Sym}^2 \Lambda^*(T_G)^{\mathcal{W}G}$;

subject to the following compatibility conditions:

- (a) For some (equivalently any) lift $d^{\text{ss}} \in \Lambda(T_G)$ of the image δ^{ss} of δ in $\pi_1(G^{\text{ss}})$, the direct sum

$$l_{\mathcal{R}} \oplus b(d^{\text{ss}} \otimes -) : \Lambda(\mathcal{R}(G)) \oplus \Lambda(T_G) \rightarrow \mathbb{Z}$$

is integral on $\Lambda(T_G)$;

(b) *The orthogonal direct sum*

$$b_{\mathcal{R}} \perp (\text{id}_{J_C} \circ b) : (\Lambda(\mathcal{R}(G)) \oplus \Lambda(T_G)) \otimes (\Lambda(\mathcal{R}(G)) \oplus \Lambda(T_G)) \rightarrow \text{End}(J_C),$$

is integral on $\Lambda(T_G) \otimes \Lambda(T_G)$, where $\text{id}_{J_C} : \mathbb{Z} \rightarrow \text{End}(J_C)$ is the canonical map given by the addition on the Abelian variety J_C .

The (contravariant) functoriality of the exact sequence is described in [3, Theorem 5.3.1(iv)]. Note that if T is a torus, the group in (iii) is trivial and the conditions (a) and (b) are always satisfied. In particular, the projection on the first two factors give an isomorphism of groups

$$\text{NS} \left(\text{Bun}_T^\delta(C) \right) \cong \Lambda^*(T) \oplus \text{Hom}^s(\Lambda(T) \otimes \Lambda(T), \text{End}(J_C)),$$

and hence the elements $(l_T, b_T, 0)$ of $\text{NS}(\text{Bun}_T^\delta(C))$ can be seen as pairs $(l_T, b_T) \in \Lambda^*(T) \oplus \text{Hom}^s(\Lambda(T) \otimes \Lambda(T), \text{End}(J_C))$. For the rest of the paper, we adopt the latter presentation for the Neron-Severi group $\text{NS}(\text{Bun}_T^\delta(C))$ for the torus case.

In general, the Neron-Severi group $\text{NS}(\text{Bun}_G^\delta(C))$ can be described as follows.

Proposition 4.2 ([3, Proposition 5.2.11]). *With the above notation, there is an exact sequence*

$$0 \rightarrow \text{NS} \left(\text{Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C) \right) \xrightarrow{\text{ab}^* \cdot \text{NS}(C)} \text{NS} \left(\text{Bun}_G^\delta(C) \right) \xrightarrow{p} \text{Bil}^{s, \text{ev}}(\Lambda(T_G))^{\mathcal{W}_G},$$

$$(l_{\mathcal{R}}, b_{\mathcal{R}}, b) \mapsto b \tag{4.3}$$

$$(l_{G^{\text{ab}}}, b_{G^{\text{ab}}}) \mapsto ((l_{G^{\text{ab}}})|_{\Lambda(\mathcal{R}(G))}, (b_{G^{\text{ab}}})|_{\Lambda(\mathcal{R}(G)) \otimes \Lambda(\mathcal{R}(G))}, 0)$$

and the image of p is equal to

$$\text{Im}(p) = \begin{cases} \text{Bil}^{s, -\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|_{\Lambda(T_{G^{\text{ss}}})})^{\mathcal{W}_G} & \text{if } g \geq 1 \\ \{b \in \text{Bil}^{s, \text{ev}}(\Lambda(T_G))^{\mathcal{W}_G} : b^{\mathbb{Q}}(d^{\text{ss}} \otimes -) \text{ is integral on } \Lambda(T_{\mathcal{D}(G)})\} & \text{if } g = 0, \end{cases}$$

where $\text{Bil}^{s, -\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|_{\Lambda(T_{G^{\text{ss}}})})^{\mathcal{W}_G}$ is defined in Definition/Lemma 2.11 and d^{ss} is some (equivalently any) lift to $\Lambda(T_{G^{\text{ss}}})$ of $\delta^{\text{ss}} \in \pi_1(G^{\text{ss}})$.

We now describe the restriction homomorphism (4.1), using the description of $\text{RPic}(\text{Bun}_{G, g, n}^\delta)$ of Theorem 3.12 and the description of $\text{Pic}(\text{Bun}_G^\delta(C))$ of Theorem 4.1.

Theorem 4.3. *Assume that $g \geq 1$ and let $(C, p_1, \dots, p_n) \in \mathcal{M}_{g, n}(k)$ be a geometric point. Let G be a reductive group and fix $\delta \in \pi_1(G)$. Then the restriction*

homomorphism (4.1) sits into the functorial commutative diagram with exact rows

$$\begin{CD}
 \Lambda^*(G^{\text{ab}}) \otimes H_{g,n} @<j_G^\delta<< \text{RPic}(\text{Bun}_{G,g,n}^\delta) @>\omega_G^\delta \oplus \gamma_G^\delta>> \text{NS}(\text{Bun}_{G,g,n}^\delta) \\
 @V\text{res}_G^\delta(C)^oVV @VV\text{res}_G^\delta(C)V @VV\text{res}_G^\delta(C)^{\text{NS}}V \\
 \text{Hom}(\pi_1(G), J_C(k)) @<j_G^\delta(C)<< \text{Pic}(\text{Bun}_G^\delta(C)) @>c_G^\delta(C)>> \text{NS}(\text{Bun}_G^\delta(C)),
 \end{CD} \tag{4.4}$$

where

- $\text{res}_G^\delta(C)^o$ sends an element $[\Lambda(G^{\text{ab}}) \xrightarrow{f} H_{g,n}] \in \text{Hom}(\Lambda(G^{\text{ab}}), H_{g,n}) = \Lambda^*(G^{\text{ab}}) \otimes H_{g,n}$ into the element

$$\left[\pi_1(G) \xrightarrow{\pi_1^{\text{(ab)}}} \pi_1(G^{\text{ab}}) = \Lambda(G^{\text{ab}}) \xrightarrow{f} H_{g,n} \xrightarrow{\iota_C} J_C(k) \right] \in \text{Hom}(\pi_1(G), J_C(k)),$$

where

$$\begin{aligned}
 \iota_C &= \iota_{(C, p_1, \dots, p_n)} : H_{g,n} \rightarrow J_C(k) \\
 (m, \zeta) &\mapsto \omega_C^m \left(\sum_{i=1}^n \zeta_i p_i \right) \quad \text{if } g \geq 2, \\
 \zeta &\mapsto \mathcal{O}_C \left(\sum_{i=1}^n \zeta_i p_i \right) \quad \text{if } g = 1;
 \end{aligned}$$

- $\text{res}_G^\delta(C)^{\text{NS}}([\chi], b) = (\chi|_{\Lambda(\mathcal{R}(G))}, \text{id}_{J_C} \circ b|_{\Lambda(\mathcal{R}(G)) \otimes \Lambda(\mathcal{R}(G))}, b|_{\Lambda(T_G) \otimes \Lambda(T_G)})$.

Proof. The theorem has been proved for a torus in [14, Proposition 4.3.3]; and, in order to prove the case of a general reductive group G , we are going to use this result for G^{ab} and for a (fixed) maximal torus $\iota : T_G \hookrightarrow G$.

Observe that the two rows of (4.4) are exact by Theorems 3.12 and 4.1. Moreover, the two outer vertical arrows are well-defined: for $\text{res}_G^\delta(C)^o$ it is clear; for $\text{res}_G^\delta(C)^{\text{NS}}$ it follows from:

- (i) Given a lift $d \in \Lambda(T_G)$ of $\delta \in \pi_1(G)$, we can choose a representative $\chi^{b(d \otimes -)} \in \Lambda^*(T_G)$ of $[\chi]$ such that $(\chi^{b(d \otimes -)})|_{\Lambda(T_{\mathcal{D}(G)})} = b(d, -)|_{\Lambda(T_{\mathcal{D}(G)})}$ by (3.28) and this implies that

$$\chi|_{\Lambda(\mathcal{R}(G))} \oplus b(d \otimes -)|_{\Lambda(T_G)} : \Lambda(\mathcal{R}(G)) \oplus \Lambda(T_G) \rightarrow \mathbb{Z}$$

is integral on $\Lambda(T_G)$, namely it is the restriction of the chosen $\chi : \Lambda(T_G) \rightarrow \mathbb{Z}$;

- (ii) The orthogonal direct sum

$$\begin{aligned}
 &\text{id}_{J_C} \circ (b|_{\Lambda(\mathcal{R}(G)) \otimes \Lambda(\mathcal{R}(G))} \perp b|_{\Lambda(T_G) \otimes \Lambda(T_G)}) : (\Lambda(\mathcal{R}(G)) \oplus \Lambda(T_G)) \\
 &\otimes (\Lambda(\mathcal{R}(G)) \oplus \Lambda(T_G)) \rightarrow \text{End}(J_C),
 \end{aligned}$$

is integral on $\Lambda(T_G) \otimes \Lambda(T_G)$, since it is the restriction of $\text{id}_{J_C} \circ b$ by the claim in the proof of Proposition 2.4.

Hence it remains to prove the commutativity of the two squares in (4.4).

In order to prove the commutativity of the left square in (4.4), consider the following diagram:

$$\begin{array}{ccccc}
 & & & & j_G^\delta \\
 & & & & \curvearrowright \\
 & & \Lambda^*(G^{\text{ab}}) \otimes H_{g,n} & \xrightarrow{j_{G^{\text{ab}}}^\delta} & \text{RPic}(\text{Bun}_{G^{\text{ab}},g,n}^\delta) & \xrightarrow{\text{ab}_\#^*} & \text{RPic}(\text{Bun}_{G,g,n}^\delta) \\
 & & \downarrow \text{res}_{G^{\text{ab}}}^\delta(C)^\circ & & \downarrow \text{res}_{G^{\text{ab}}}^\delta(C) & & \downarrow \text{res}_G^\delta(C) \\
 \text{res}_G^\delta(C)^\circ \curvearrowleft & & \text{Hom}(\pi_1(G^{\text{ab}}), J_C(k)) & \xrightarrow{j_{G^{\text{ab}}}^\delta(C)} & \text{Pic}(\text{Bun}_{G^{\text{ab}}}(C)) & \xrightarrow{\text{ab}^{*,\text{NS}}(C)} & \text{Pic}(\text{Bun}_G^\delta(C)) \\
 & & \downarrow \text{Hom}(\pi_1(\text{ab}), -) & & \downarrow j_G^\delta(C) & & \downarrow j_G^\delta(C) \\
 & & \text{Hom}(\pi_1(G), J_C(k)) & & & &
 \end{array}
 \tag{4.5}$$

In the above diagram, every simple subdiagram commutes: the curved triangles commutes by the definition of j_G^δ and of $\text{res}_G^\delta(C)^\circ$; the left square commutes by [14, Proposition 4.3.3] applied to G^{ab} ; the right square commutes by the obvious functoriality of the restriction homomorphism; the lower triangle commutes by the functoriality of the morphism j_G^δ (see Theorem 4.1). By using all the above commutativity results, we deduce that $\text{res}_G^\delta(C) \circ j_G^\delta = j_G^\delta(C) \circ \text{res}_G^\delta(C)^\circ$, i.e., that the left square of (4.4) commutes.

In order to prove the commutativity of the right square in (4.4), choose a lift $d \in \Lambda(T_G)$ of $\delta \in \pi_1(G)$ and consider the diagram

$$\begin{array}{ccccc}
 \text{RPic}(\text{Bun}_{G,g,n}^\delta) & \xrightarrow{\omega_G^\delta \oplus \gamma_G^\delta} & \text{NS}(\text{Bun}_{G,g,n}^\delta) & & \\
 \downarrow \text{res}_G^\delta(C) & \searrow \iota_\#^* & \downarrow \text{res}_G^\delta(C)^{\text{NS}} & \searrow \iota^{*,\text{NS}} & \\
 & \text{RPic}(\text{Bun}_{T_G,g,n}^d) & \xrightarrow{\omega_{T_G}^d \oplus \gamma_{T_G}^d} & \text{NS}(\text{Bun}_{T_G,g,n}^d) & \\
 & \downarrow \text{res}_{T_G}^d(C) & & \downarrow \text{res}_{T_G}^d(C)^{\text{NS}} & \\
 \text{Pic}(\text{Bun}_G^\delta(C)) & \xrightarrow{c_G^\delta(C)} & \text{NS}(\text{Bun}_G^\delta(C)) & & \\
 & \searrow \iota_\#(C)^* & \downarrow \text{res}_G^\delta(C) & \searrow \iota^{*,\text{NS}}(C) & \\
 & \text{Pic}(\text{Bun}_{T_G}^d(C)) & \xrightarrow{c_{T_G}^d(C)} & \text{NS}(\text{Bun}_{T_G}^d(C)) &
 \end{array}
 \tag{4.6}$$

where $\iota^{*,\text{NS}}$ is the morphism of Definition/Lemma 3.10 and $\iota^{*,\text{NS}}(C)$ is the morphism of [3, Definition 5.2.5], and they are given by the formulas:

- $\iota^{*,\text{NS}}([\chi], b) = (\chi^{b(d \otimes -)}, b) \in \text{NS}(\text{Bun}_{T_G,g,n}^d)$, see (3.29);
- $\iota^{*,\text{NS}}(C)(l_{\mathcal{R}}, b_{\mathcal{R}}, b) = (l_{\mathcal{R}} \oplus b(d^{\text{ss}} \otimes -), b_{\mathcal{R}} \perp \text{id}_{J_C} \circ b) \in \text{NS}(\text{Bun}_{T_G}^d(C))$, where d^{ss} is the image of $d \in \Lambda(T_G)$ in $\Lambda(T_G^{\text{ss}})$, which is well-defined by conditions (4.1) and (4.1) of Theorem 4.1.

We have the following commutativity properties in the above diagram (4.6):

- (a) $\text{res}_{T_G}^d(C) \circ \iota_{\#}^* = \iota_{\#}(C)^* \circ \text{res}_G^{\delta}(C)$, which follows from the functoriality of the restriction homomorphism;
- (b) $\text{res}_{T_G}^d(C)^{\text{NS}} \circ \iota^{*,\text{NS}} = \iota^{*,\text{NS}}(C) \circ \text{res}_G^{\delta}(C)^{\text{NS}}$. In fact, for any $([\chi], b) \in \text{NS}(\text{Bun}_{G,g,n}^{\delta})$ we compute

$$\left\{ \begin{array}{l} (\text{res}_{T_G}^d(C)^{\text{NS}} \circ \iota^{*,\text{NS}})([\chi], b) = \text{res}_{T_G}^d(C)^{\text{NS}}(\chi^{b(d \otimes -)}, b) \\ \quad = (\chi^{b(d \otimes -)}, \text{id}_{J_C} \circ b) \\ (\iota^{*,\text{NS}}(C) \circ \text{res}_G^{\delta}(C)^{\text{NS}})([\chi], b) \\ \quad = \iota^{*,\text{NS}}(C)(\chi|_{\Lambda(\mathcal{A}(G))}, \text{id}_{J_C} \circ b|_{\Lambda(\mathcal{A}(G)) \otimes \Lambda(\mathcal{A}(G))}, b|_{\Lambda(T_G) \otimes \Lambda(T_G)}) \\ \quad = (\chi|_{\Lambda(\mathcal{A}(G))} \oplus b(d^{\text{ss}} \otimes -)|_{\Lambda(T_G)}, \text{id}_{J_C} \\ \quad \quad \circ (b|_{\Lambda(\mathcal{A}(G)) \otimes \Lambda(\mathcal{A}(G))} \perp b|_{\Lambda(T_G) \otimes \Lambda(T_G)})), \end{array} \right.$$

and the two results coincide by the fact that

$$b(d \otimes -)|_{\Lambda(T_G)} = b(d^{\text{ss}} \otimes -)|_{\Lambda(T_G)}$$

by the claim in the proof of Proposition 2.4, together with condition (ii);

- (c) $c_{T_G}^d(C) \circ \text{res}_{T_G}^d(C) = \text{res}_{T_G}^d(C)^{\text{NS}} \circ (\omega_{T_G}^d \oplus \gamma_{T_G}^d)$, which follows from Proposition [14, Proposition 4.3.3] applied to T_G ;
- (d) $\iota^{*,\text{NS}}(C) \circ c_G^{\delta}(C) = c_{T_G}^d(C) \circ \iota_{\#}(C)^*$, which follows from the functoriality of the exact sequence (4.2) together with [3, Definition 5.2.5, 5.2.7];
- (e) $\iota^{*,\text{NS}} \circ (\omega_G^{\delta} \oplus \gamma_G^{\delta}) = (\omega_{T_G}^d \oplus \gamma_{T_G}^d) \circ \iota_{\#}^*$, which follows from the functoriality of the homomorphism $\omega_G^{\delta} \oplus \gamma_G^{\delta}$ applied to the inclusion $\iota : T_G \hookrightarrow G$ (see Theorem 3.12(1)).

Using the above commutativity relations together with a diagram chase in (4.6), we conclude that

$$\iota^{*,\text{NS}}(C) \circ c_G^{\delta}(C) \circ \text{res}_G^{\delta}(C) = \iota^{*,\text{NS}}(C) \circ \text{res}_G^{\delta}(C)^{\text{NS}} \circ (\omega_G^{\delta} \oplus \gamma_G^{\delta}).$$

Since $\iota^{*,\text{NS}}(C)$ is injective (using that id_{J_C} is injective since $g \geq 1$), we can simplify $\iota^{*,\text{NS}}(C)$ from the above expression, and we get

$$c_G^{\delta}(C) \circ \text{res}_G^{\delta}(C) = \text{res}_G^{\delta}(C)^{\text{NS}} \circ (\omega_G^{\delta} \oplus \gamma_G^{\delta}),$$

which is exactly the commutativity of the right square in (4.4), and this concludes the proof. □

Corollary 4.4. *Keep the notation of Theorem 4.3. The kernel of the restriction homomorphism $\text{res}_G^{\delta}(C)$ is equal to*

$$\ker \left(\text{res}_G^{\delta}(C) \right) = \Lambda^*(G^{\text{ab}}) \otimes \ker \iota_C \subseteq \Lambda^*(G^{\text{ab}}) \otimes H_{g,n}.$$

Proof. This follows from the snake lemma applied to the exact sequence (4.4), using that $\text{res}_G^\delta(C)^{\text{NS}}$ is injective and the explicit description of $\text{res}_G^\delta(C)^o$. \square

We now give an alternative description of the restriction homomorphism onto the Neron-Severi group

$$\overline{\text{res}}_G^\delta(C) : \text{RPic}(\text{Bun}_{G,g,n}^\delta) \xrightarrow{\text{res}_G^\delta(C)} \text{Pic}(\text{Bun}_G^\delta(C)) \xrightarrow{c_G^\delta(C)} \text{NS}(\text{Bun}_G^\delta(C)), \quad (4.7)$$

for any $(C, p_1, \dots, p_n) \in \mathcal{M}_{g,n}(k)$, using the description of $\text{RPic}(\text{Bun}_{G,g,n}^\delta)$ in Corollary 3.4 and the description of $\text{NS}(\text{Bun}_G^\delta(C))$ in Proposition 4.2.

Theorem 4.5. *Assume that $g \geq 1$ and let $(C, p_1, \dots, p_n) \in \mathcal{M}_{g,n}(k)$ be a geometric point. Let G be a reductive group and fix $\delta \in \pi_1(G)$. The homomorphism (4.7) sits into the following functorial commutative diagram with exact rows:*

$$\begin{CD} \text{RPic}(\text{Bun}_{G^{ab},g,n}^{\delta^{ab}}) @<{\hookrightarrow}^{ab^\#}<< \text{RPic}(\text{Bun}_{G,g,n}^\delta) @>{\theta_G^\delta}>> \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} \\ @VV{\overline{\text{res}}_{G^{ab}}^{\delta^{ab}}(C)}V @VV{\text{res}_G^\delta(C)}V @VV{r_G}V \\ \text{NS}(\text{Bun}_{G^{ab}}^{\delta^{ab}}(C)) @<{\hookrightarrow}^{ab^*,\text{NS}(C)}<< \text{NS}(\text{Bun}_G^\delta(C)) @>{p}>> \text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} \end{CD} \quad (4.8)$$

where r_G is the injective homomorphism in Definition/Lemma 2.11.

Proof. The rows of (4.8) are exact by Corollary 3.4 and Proposition 4.2. We need to show the commutativity of the diagram (4.8).

The commutativity of the left square of (4.8) follows from the functoriality of the restriction homomorphism $\text{res}_G^\delta(C)$ and the functoriality of the homomorphism $c_G^\delta(C)$ (see Theorem 4.1).

Consider now the following diagram:

$$\begin{CD} \text{RPic}(\text{Bun}_{G,g,n}^\delta) @>{\theta_G^\delta}>> \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} \\ @VV{\omega_G^\delta \oplus \gamma_G^\delta}V @| \\ \overline{\text{res}}_G^\delta(C) @>{\text{res}_{\mathcal{D}}^{\text{NS}}}>> \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} \\ @VV{\text{res}_G^\delta(C)^{\text{NS}}}V @VV{r_G}V \\ \text{NS}(\text{Bun}_G^\delta(C)) @>{p}>> \text{Bil}^s(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} \cap \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathscr{W}_G}. \end{CD} \quad (4.9)$$

The commutativity of the right square of (4.8) follows from the following commutativity results:

- (a) The left triangle commutes because of the definition (4.7) of $\overline{\text{res}}_G^\delta(C)$ and of the commutativity of the diagram (4.4);

(b) The upper square commutes, *i.e.*, $\theta_G^\delta = \text{res}_{\mathcal{D}}^{\text{NS}} \circ (\omega_G^\delta \oplus \gamma_G^\delta)$.

In order to prove this, it is enough to show, using that $\text{RPic}(\text{Bun}_{G,g,n}^\delta)$ is the pushout (3.9), that

$$\theta_G^\delta \circ \text{ab}_\#^* = \text{res}_{\mathcal{D}}^{\text{NS}} \circ (\omega_G^\delta \oplus \gamma_G^\delta) \circ \text{ab}_\#^*, \tag{4.10}$$

$$\theta_G^\delta \circ \tau_G^\delta = \text{res}_{\mathcal{D}}^{\text{NS}} \circ (\omega_G^\delta \oplus \gamma_G^\delta) \circ \tau_G^\delta. \tag{4.11}$$

Condition (4.10) holds since $\theta_G^\delta \circ \text{ab}_\#^* = 0$ by (3.10) and $\text{res}_{\mathcal{D}}^{\text{NS}} \circ (\omega_G^\delta \oplus \gamma_G^\delta) \circ \text{ab}_\#^* = 0$ which follows from the fact that $(\omega_G^\delta \oplus \gamma_G^\delta) \circ \text{ab}_\#^* = \text{ab}^{*,\text{NS}} \circ (\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} \oplus \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}})$ (by the functoriality of the morphism $\omega_G^\delta \oplus \gamma_G^\delta$, see Theorem 3.12(1)) together with (3.33). Condition (4.11) follows from

$$\begin{cases} (\text{res}_{\mathcal{D}}^{\text{NS}} \circ (\omega_G^\delta \oplus \gamma_G^\delta) \circ \tau_G^\delta)(b) = \text{res}_{\mathcal{D}}^{\text{NS}}(b(\delta \otimes -), b) = b_{|\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_{\mathcal{D}(G)})} & \text{by (3.38)} \\ (\theta_G^\delta \circ \tau_G^\delta)(b) = b_{|\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_{\mathcal{D}(G)})} & \text{by Corollary 3.4;} \end{cases}$$

(c) The lower square commutes because, using the definitions of the maps involved, we have

$$(r_G \circ \text{res}_{\mathcal{D}}^{\text{NS}})([\chi], b) = b_{|\Lambda(T_G) \otimes \Lambda(T_G)} = (p \circ \text{res}_G^\delta(C)^{\text{NS}})([\chi], b). \quad \square$$

Corollary 4.6. *Keep the notation of Theorem 4.5 and assume furthermore that $\text{id}_{J_C} : \mathbb{Z} \xrightarrow{\cong} \text{End}(J_C)$ is an isomorphism. Then there exists a canonical short exact sequence*

$$0 \rightarrow \text{coker}(\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} \oplus \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}) \rightarrow \text{coker}(\overline{\text{res}_G^\delta(C)}) \rightarrow \text{coker}(r_G) \rightarrow 0. \tag{4.12}$$

In particular, if $n > 0$ then we have the canonical isomorphism

$$\text{coker}(\overline{\text{res}_G^\delta(C)}) \xrightarrow{\cong} \text{coker}(r_G).$$

Note that if k is uncountable and C is very general in $\mathcal{M}_g(k)$, then $\text{id}_{J_C} : \mathbb{Z} \xrightarrow{\cong} \text{End}(J_C)$ is an isomorphism, as it follows easily from [17].

Proof. By applying the snake lemma to (4.8) and using that r_G is injective, we get the exact sequence

$$0 \rightarrow \text{coker}(\overline{\text{res}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C)}) \rightarrow \text{coker}(\overline{\text{res}_G^\delta(C)}) \rightarrow \text{coker}(r_G) \rightarrow 0. \tag{4.13}$$

By Theorem 4.3, the morphism $\overline{\text{res}}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C)$ admits the following factorization:

$$\begin{aligned} \overline{\text{res}}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C) : \text{RPic}\left(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}\right) &\xrightarrow{\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} \oplus \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}} \text{NS}\left(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}\right) \\ &\xrightarrow{\text{res}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C)^{\text{NS}}} \text{NS}\left(\overline{\text{res}}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C)\right). \end{aligned}$$

By the assumption that id_{J_C} is an isomorphism together with the explicit description of $\text{res}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C)^{\text{NS}}$ in Theorem 4.3, we deduce that $\text{res}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C)^{\text{NS}}$ is an isomorphism. Hence, we get a canonical isomorphism

$$\text{coker}\left(\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} \oplus \gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}\right) \xrightarrow{\cong} \text{coker}\left(\overline{\text{res}}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C)\right). \tag{4.14}$$

By combining the short exact sequence (4.13) with the isomorphism (4.14), we conclude. \square

We end this section by describing the restriction homomorphism (4.1) in genus 0.

Remark 4.7. In genus 0, the only smooth curve is the projective line \mathbb{P}^1 . In this situation, it follows from the proof of [14, Proposition 5.4.1] that the restriction homomorphism

$$\text{res}_G^\delta(\mathbb{P}^1) : \text{RPic}\left(\text{Bun}_{G,0,n}^\delta\right) \rightarrow \text{Pic}\left(\text{Bun}_G^\delta(\mathbb{P}^1)\right)$$

is an isomorphism if $n > 0$ and injective if $n = 0$, for any $(\mathbb{P}^1, p_1, \dots, p_n) \in \mathcal{M}_{0,n}(k)$.

Using Theorem 4.1, it follows that if $n > 0$ then we have

$$\begin{aligned} &\text{RPic}\left(\text{Bun}_{G,0,n}^\delta\right) \\ &= \left\{ (l_{\mathcal{R}}, b) \in \Lambda^*(\mathcal{R}(G)) \times \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathscr{W}G} : l_{\mathcal{R}} \oplus b(d^{\text{ss}} \otimes -) \in \Lambda^*(T_G) \right\}, \end{aligned} \tag{4.15}$$

for some (equivalently any) lift $d^{\text{ss}} \in \Lambda(T_G)$ of the image δ^{ss} of δ in $\pi_1(G^{\text{ss}})$.

Moreover, using Proposition 4.2, we have the following exact sequence for $n > 0$:

$$\begin{aligned} \text{RPic}\left(\text{Bun}_{G^{\text{ab}},0,n}^{\delta^{\text{ab}}}\right) &\xrightarrow{\text{ab}_\#^*} \text{RPic}\left(\text{Bun}_{G,0,n}^\delta\right) \xrightarrow{\theta_G^\delta} \left\{ \begin{array}{l} b \in \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathscr{W}G} : \\ b^{\mathbb{Q}}(d^{\text{ss}} \otimes -) \text{ is integral on } \Lambda(T_{\mathcal{R}(G)}) \end{array} \right\} \\ (l_{\mathcal{R}}, b) &\mapsto b \\ l_{G^{\text{ab}}} &\mapsto (l_{G^{\text{ab}}}|_{\mathcal{R}(G)}, 0). \end{aligned} \tag{4.16}$$

5. The rigidification $\text{Bun}_{G,g,n}^\delta // \mathcal{Z}(G)$ and its Picard group

Since the center $\mathcal{Z}(G)$ of a reductive group G acts functorially on any G -bundle, we have that $\mathcal{Z}(G)$ sits functorially inside the automorphism group of any S -point $(\mathcal{C} \rightarrow S, \underline{\sigma}, E)$ of $\text{Bun}_{G,g,n}^\delta$ for any $\delta \in \pi_1(G)$. Hence we can form the rigidification

$$v_G^\delta : \text{Bun}_{G,g,n}^\delta \rightarrow \text{Bun}_{G,g,n}^\delta // \mathcal{Z}(G) := \mathfrak{Bun}_{G,g,n}^\delta, \tag{5.1}$$

which turns out to be a $\mathcal{Z}(G)$ -gerbe, *i.e.*, a gerbe banded by $\mathcal{Z}(G)$. The aim of this section is to study the Picard group of $\mathfrak{Bun}_{G,g,n}^\delta$ and the class of the $\mathcal{Z}(G)$ -gerbe v_G^δ .

From the Leray spectral sequence

$$E_{p,q}^2 = H^p \left(\mathfrak{Bun}_{G,g,n}^\delta, R^q(v_G^\delta)_*(\mathbb{G}_m) \right) \Rightarrow H^{p+q} \left(\text{Bun}_{G,g,n}^\delta, \mathbb{G}_m \right),$$

and using that $(v_G^\delta)_*(\mathbb{G}_m) = \mathbb{G}_m$ and that $R^1(v_G^\delta)_*(\mathbb{G}_m)$ is the constant sheaf $\Lambda^*(\mathcal{Z}(G)) = \text{Hom}(\mathcal{Z}(G), \mathbb{G}_m)$, we get the exact sequence

$$\begin{aligned} \text{Pic} \left(\mathfrak{Bun}_{G,g,n}^\delta \right) &\xrightarrow{(v_G^\delta)^*} \text{Pic} \left(\text{Bun}_{G,g,n}^\delta \right) \xrightarrow{\text{wt}_G^\delta} \Lambda^*(\mathcal{Z}(G)) \\ &\xrightarrow{\text{obs}_G^\delta} H^2 \left(\mathfrak{Bun}_{G,g,n}^\delta, \mathbb{G}_m \right) \xrightarrow{(v_G^\delta)^*} H^2 \left(\text{Bun}_{G,g,n}^\delta, \mathbb{G}_m \right). \end{aligned} \tag{5.2}$$

The homomorphism wt_G^δ (called the *weight homomorphism*) has the following geometric interpretation: given a line bundle \mathcal{L} on $\text{Bun}_{G,g,n}^\delta$, the character $\text{wt}_G^\delta(\mathcal{L}) \in \text{Hom}(\mathcal{Z}(G), \mathbb{G}_m)$ is such that, for any $\mathcal{E} := (\mathcal{C} \rightarrow S, \underline{\sigma}, E) \in \text{Bun}_{G,g,n}^\delta(S)$, we have the factorization

$$\text{wt}_G^\delta(S) : \mathcal{Z}(G)(S) \hookrightarrow \text{Aut}_{\text{Bun}_{G,g,n}^\delta(S)}(\mathcal{E}) \xrightarrow{\text{Aut}(\mathcal{L}_S)} \text{Aut}_{\mathcal{O}_S}(\mathcal{L}_S(\mathcal{E})) = \mathbb{G}_m(S),$$

where the first homomorphism is given by the canonical action of $\mathcal{Z}(G)$ on every G -gerbe, and the second homomorphism is induced by the functor of groupoids

$$\mathcal{L}_S : \text{Bun}_{G,g,n}^\delta(S) \rightarrow \{\text{Line bundles on } S\}$$

determined by \mathcal{L} .

The homomorphism obs_G^δ (called the *obstruction homomorphism*) has the following geometric interpretation: given any character $\lambda \in \text{Hom}(\mathcal{Z}(G), \mathbb{G}_m)$, the element $\text{obs}_G^\delta(\lambda)$ is the class in $H^2(\mathfrak{Bun}_{G,g,n}^\delta, \mathbb{G}_m)$ of the \mathbb{G}_m -gerbe $\lambda_*(v_G^\delta)$ obtained by pushing forward the $\mathcal{Z}(G)$ -gerbe v_G^δ along λ .

If we take the fiber of (5.1) over a geometric point $(C, p_1, \dots, p_n) \in \mathcal{M}_{g,n}(k)$, we get the $\mathcal{Z}(G)$ -gerbe

$$v_G^\delta(C) : \text{Bun}_G^\delta(C) \rightarrow \mathfrak{Bun}_G^\delta(C) := \text{Bun}_G^\delta(C) // \mathcal{Z}(G). \tag{5.3}$$

The Leray spectral sequence for \mathbb{G}_m relative to the $\mathcal{L}(G)$ -gerbe gives the exact sequence (analogously to (5.2))

$$\begin{aligned} \text{Pic} \left(\mathfrak{Bun}_G^\delta(C) \right) &\xrightarrow{v_G^\delta(C)^*} \text{Pic} \left(\text{Bun}_G^\delta(C) \right) \\ &\xrightarrow{\text{wt}_G^\delta(C)} \Lambda^*(\mathcal{L}(G)) \xrightarrow{\text{obs}_G^\delta(C)} H^2 \left(\mathfrak{Bun}_G^\delta(C), \mathbb{G}_m \right). \end{aligned} \tag{5.4}$$

The weight homomorphism $\text{wt}_G^\delta(C)$ and its cokernel, which coincides with the image of $\text{obs}_G^\delta(C)$ by (5.3), have been determined by Biswas-Hoffmann [5, Proposition 7.2], as we now recall (in a form which is slightly different from [5]).

Theorem 5.1 ([5]). *Let C be a (irreducible, projective, smooth) curve of genus $g \geq 1$ over $\bar{k} = k$. Let G be a reductive group G and fix $\delta \in \pi_1(G)$.*

(1) *The weight homomorphism $\text{wt}_G^\delta(C)$ factors as*

$$\begin{aligned} \text{wt}_G^\delta(C) : \text{Pic} \left(\text{Bun}_G^\delta(C) \right) &\xrightarrow{c_G^\delta(C)} \text{NS} \left(\text{Bun}_G^\delta(C) \right) \xrightarrow{\overline{\text{wt}}_G^\delta(C)} \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} \\ &(l_{\mathcal{A}}, b_{\mathcal{A}}, b) \mapsto [l_{\mathcal{A}} \oplus b(d^{\text{ss}} \otimes -)], \end{aligned} \tag{5.5}$$

where $c_G^\delta(C)$ is the homomorphism of Theorem 4.1 and $d^{\text{ss}} \in \Lambda(T_{G^{\text{ss}}})$ is any lift of the image δ^{ss} of δ in $\pi_1(G^{\text{ss}})$;

(2) *The homomorphism $\Lambda_{\mathcal{O}}^*$ of (2.17) induces an isomorphism*

$$\widetilde{\Lambda}_{\mathcal{O}}^* : \text{coker} \left(\text{wt}_G^\delta(C) \right) \xrightarrow{\cong} \text{coker} \left(\widetilde{\text{ev}}_{\mathcal{O}(G)}^\delta \right), \tag{5.6}$$

where $\widetilde{\text{ev}}_{\mathcal{O}(G)}^\delta$ is the homomorphism of Definition/Lemma 2.11.

Proof. Part (1): note that $\overline{\text{wt}}_G^\delta(C)$ is well-defined since $l_{\mathcal{A}} \oplus b(d^{\text{ss}} \otimes -)$ is integral on $\Lambda(T_G)$ by condition (4.1) of Theorem 4.1 and its class in $\frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})}$ is independent of the choice of the lift d^{ss} by Lemma 2.9. The equality $\text{wt}_G^\delta(C) = \overline{\text{wt}}_G^\delta(C) \circ c_G^\delta(C)$ is a consequence of the following factorization (see [5, Equation (5)]):

$$\begin{aligned} \text{wt}_G^\delta(C) : \text{Pic} \left(\text{Bun}_G^\delta(C) \right) &\xrightarrow{c_G^\delta(C)} \text{NS} \left(\text{Bun}_G^\delta(C) \right) \xrightarrow{\iota^{*,\text{NS}}(C)} \text{NS} \left(\text{Bun}_{T_G}^d(C) \right) \\ &\xrightarrow{p_1} \Lambda^*(T_G) \twoheadrightarrow \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} \end{aligned} \tag{5.7}$$

where $c_G^\delta(C)$ is the homomorphism of Theorem 4.1, $d \in \Lambda(T_G)$ is any lift of $\delta \in \pi_1(G)$, $\iota^{*,\text{NS}}(C)$ is the homomorphism defined in the second bullet below the diagram (4.6), $p_1 : \text{NS}(\text{Bun}_{T_G}^d(C)) \rightarrow \Lambda^*(T_G)$ is the projection onto the first factor.

Part (2): from part (1) and using that $c_G^\delta(C)$ is surjective by Theorem 4.1, we get that $\text{coker}(\text{wt}_G^\delta(C)) = \text{coker}(\overline{\text{wt}}_G^\delta(C))$. From the definition of $\overline{\text{wt}}_G^\delta(C)$, using Proposition 4.2 and the vertical exact sequence of (2.17), we get the commutative diagram with exact rows

$$\begin{array}{ccccc}
 \text{NS}(\text{Bun}_{G^{\text{ab}},g,n}^{\delta\text{ab}}(C)) & \xrightarrow{\text{c}^{\text{ab}^*.\text{NS}}(C)} & \text{NS}(\text{Bun}_{G,g,n}^\delta(C)) & \xrightarrow{P} & \text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}G} \\
 \downarrow \overline{\text{wt}}_{G^{\text{ab}}}^{\delta\text{ab}}(C) & & \downarrow \overline{\text{wt}}_G^\delta(C) & & \downarrow \overline{\text{ev}}_{\mathcal{D}(G)}^\delta \\
 \Lambda^*(G^{\text{ab}}) & \xrightarrow{\overline{\Lambda}_{\text{ab}}^*} & \Lambda^*(T_G) & \xrightarrow{\overline{\Lambda}_{\mathcal{D}}^*} & \Lambda^*(T_{\mathcal{D}(G)}) \\
 & & \Lambda^*(T_{G^{\text{ad}}}) & & \Lambda^*(T_{G^{\text{ad}}})
 \end{array} \tag{5.8}$$

From the definition, it follows that $\overline{\text{wt}}_{G^{\text{ab}}}^{\delta\text{ab}}(C)$ is the projection onto the first factor of $\text{NS}(\text{Bun}_{G^{\text{ab}},g,n}^{\delta\text{ab}}(C))$, and hence it is surjective. Therefore the snake lemma applied to (5.8) gives the conclusion. \square

Using the above result, we can now give an explicit expression for the weight homomorphism wt_G^δ .

Proposition 5.2. *Assume that $g \geq 1$. Fix a maximal torus $\iota : T_G \hookrightarrow G$ of the reductive group G and let $\delta \in \pi_1(G)$. The weight homomorphism wt_G^δ is equal to the following composition:*

$$\text{wt}_G^\delta : \text{Pic}(\text{Bun}_{G,g,n}^\delta) \rightarrow \text{RPic}(\text{Bun}_{G,g,n}^\delta) \xrightarrow{\omega_G^\delta} \Lambda^*(\mathcal{L}(G)) = \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})},$$

where ω_G^δ is the homomorphism of Definition/Lemma 3.7.

Proof. From the geometric description of the weight homomorphism, it follows that wt_G^δ can be computed on any fiber of Φ_G^δ over a geometric point $(C, p_1, \dots, p_n) \in \mathcal{M}_{g,n}(k)$, i.e., wt_G^δ factors as the composition

$$\begin{array}{ccc}
 \text{wt}_G^\delta : \text{Pic}(\text{Bun}_{G,g,n}^\delta) & \rightarrow & \text{RPic}(\text{Bun}_{G,g,n}^\delta) \\
 \text{res}_G^\delta(C) \downarrow & & \downarrow \text{wt}_G^\delta(C) \\
 \text{Pic}(\text{Bun}_G^\delta(C)) & \xrightarrow{\text{wt}_G^\delta(C)} & \Lambda^*(\mathcal{L}(G)),
 \end{array} \tag{5.9}$$

where $\text{res}_G^\delta(C)$ is the restriction homomorphism (4.1) and $\text{wt}_G^\delta(C)$ is the weight homomorphism of (5.4).

From Theorems 4.3 and 5.1, it follows that the composition $\text{wt}_G^\delta(C) \circ \text{res}_G^\delta(C)$ is equal to the morphism $\omega_G^\delta \oplus \gamma_G^\delta : \text{RPic}(\text{Bun}_{G,g,n}^\delta) \rightarrow \text{NS}(\text{Bun}_{G,g,n}^\delta)$ followed the composition

$$\text{NS}(\text{Bun}_{G,g,n}^\delta) \xrightarrow{\text{res}_G^\delta(C)^{\text{NS}}} \text{NS}(\text{Bun}_G^\delta(C)) \xrightarrow{\overline{\text{wt}}_G^\delta(C)} \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})}, \tag{5.10}$$

which sends an element $([\chi], b) \in \text{NS}(\text{Bun}_{G,g,n}^\delta)$ into

$$[\chi|_{\mathcal{L}(G)} \oplus b(d \otimes -)|_{\Lambda(T_G)}] = [\chi] \in \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}}}.$$

Therefore, the composition $\text{wt}_G^\delta(C) \circ \text{res}_G^\delta(C)$ is equal to the morphism ω_G^δ , and we conclude by (5.9). \square

Note that the forgetful morphism $\Phi_G^\delta : \text{Bun}_{G,g,n}^\delta \rightarrow \mathcal{M}_{g,n}$ factors as

$$\Phi_G^\delta : \text{Bun}_{G,g,n}^\delta \xrightarrow{v_G^\delta} \mathfrak{Bun}_{G,g,n}^\delta \xrightarrow{\Psi_G^\delta} \mathcal{M}_{g,n},$$

where the morphism Ψ_G^δ is fpqc (i.e., faithfully flat and locally quasi-compact) and cohomologically flat in degree zero (i.e., the natural morphism $(\Psi_G^\delta)^\sharp : \mathcal{O}_{\mathcal{M}_{g,n}} \rightarrow (\Psi_G^\delta)_*(\mathcal{O}_{\mathfrak{Bun}_{G,g,n}^\delta})$ is a universal isomorphism) since Φ_G^δ satisfies these properties (see Theorem 3.1(2)) and v_G^δ is smooth, surjective, and cohomologically flat in degree zero.

Hence, the pull-back morphism Ψ_G^δ is injective on Picard groups and we have an injective pull-back morphism on the relative Picard groups

$$\overline{(v_G^\delta)^*} : \text{RPic}(\mathfrak{Bun}_{G,g,n}^\delta) := \frac{\text{Pic}(\mathfrak{Bun}_{G,g,n}^\delta)}{(\Psi_G^\delta)^*(\text{Pic}(\mathcal{M}_{g,n}))} \hookrightarrow \text{RPic}(\text{Bun}_{G,g,n}^\delta) := \frac{\text{Pic}(\text{Bun}_{G,g,n}^\delta)}{(\Phi_G^\delta)^*(\text{Pic}(\mathcal{M}_{g,n}))}. \tag{5.11}$$

Therefore, combining (5.2) with Proposition 5.2, we get the new exact sequence

$$\begin{aligned} \text{RPic}(\mathfrak{Bun}_{G,g,n}^\delta) &\xrightarrow{\overline{(v_G^\delta)^*}} \text{RPic}(\text{Bun}_{G,g,n}^\delta) \xrightarrow{\omega_G^\delta} \Lambda^*(\mathcal{L}(G)) \\ &\xrightarrow{\text{obs}_G^\delta} H^2(\mathfrak{Bun}_{G,g,n}^\delta, \mathbb{G}_m) \xrightarrow{(v_G^\delta)^*} H^2(\text{Bun}_{G,g,n}^\delta, \mathbb{G}_m). \end{aligned} \tag{5.12}$$

We now want to compute the kernel of ω_G^δ , which can be identified with the Picard group of $\mathfrak{Bun}_{G,g,n}^\delta$, and the cokernel of ω_G^δ , which is an obstruction to the vanishing of the obstruction morphism. With this aim, we introduce the following group.

Definition 5.3. Let G be a reductive group with fixed maximal torus T_G and Weyl group \mathscr{W}_G , and fix $\delta \in \pi_1(G)$. Denote by

$$\text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) \subset \text{Bil}^{s, \mathscr{D}\text{-ev}}(\Lambda(T_G))^{\mathscr{W}_G}$$

the subgroup consisting of those $b \in \text{Bil}^{s, \mathscr{D}\text{-ev}}(\Lambda(T_G))^{\mathscr{W}_G}$ such that

$$0 = (\text{ev}_{\mathscr{D}(G)}^\delta \circ \text{res}_{\mathscr{D}})(b) = b(\delta \otimes -)|_{\Lambda(T_{\mathscr{D}(G)})} := [b(d \otimes -)|_{\Lambda(T_{\mathscr{D}(G)})}] \in \frac{\Lambda^*(T_{\mathscr{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}}}}, \tag{5.13}$$

where $d \in \Lambda(T_G)$ is any lift of δ , and $\text{ev}_{\mathscr{D}(G)}^\delta$ and $\text{res}_{\mathscr{D}}$ are the homomorphisms of Definition/Lemma 2.8.

The relation between the groups $\text{NS}(\mathfrak{Bun}_{G,g,n}^\delta)$ and $\text{NS}(\text{Bun}_{G,g,n}^\delta)$ is explained in the following

Proposition 5.4. *Let G be a reductive group with maximal torus T_G and Weyl group \mathcal{W}_G , and fix $\delta \in \pi_1(G)$. We have the following commutative diagram, with exact rows and columns:*

$$\begin{array}{ccccc}
 \text{Bil}^s(\Lambda(G^{\text{ab}})) = \text{NS}(\mathfrak{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}) & \xrightarrow{\widetilde{\text{ab}}^{*,\text{NS}}} & \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) & \xrightarrow{\widetilde{\text{res}}_{\mathcal{D}}^{\text{NS}}} & \ker((\text{ev}_{\mathcal{D}(G)}^\delta)) \\
 \downarrow \nu_{G^{\text{ab}}}^{\delta^{\text{ab}},\text{NS}} & & \downarrow \nu_G^{\delta,\text{NS}} & & \downarrow \\
 \Lambda^*(G^{\text{ab}}) \oplus \text{Bil}^s(\Lambda(G^{\text{ab}})) = \text{NS}(\text{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}) & \xrightarrow{\text{ab}^{*,\text{NS}}} & \text{NS}(\text{Bun}_{G,g,n}^\delta) & \xrightarrow{\text{res}_{\mathcal{D}}^{\text{NS}}} & \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^*}))^{\mathcal{W}_G} \\
 \downarrow \omega_{G^{\text{ab}}}^{\delta^{\text{ab}},\text{NS}} & & \downarrow \omega_G^{\delta,\text{NS}} & & \downarrow \text{ev}_{\mathcal{D}(G)}^\delta \\
 \Lambda^*(G^{\text{ab}}) \hookrightarrow & \xrightarrow{\overline{\Lambda}_{\text{ab}}^*} & \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} & \xrightarrow{\overline{\Lambda}_{\mathcal{D}}^*} & \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}})} \\
 & & \downarrow & & \downarrow \\
 & & \text{coker}(\omega_G^{\delta,\text{NS}}) & \xrightarrow{\cong} & \text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta)
 \end{array} \tag{5.14}$$

where the identification $\text{NS}(\mathfrak{Bun}_{G^{\text{ab}},g,n}^{\delta^{\text{ab}}}) = \text{Bil}^s(\Lambda(G^{\text{ab}}))$ follows from Definition 5.3, the morphisms $\widetilde{\text{ab}}^{*,\text{NS}}$ and $\widetilde{\text{res}}_{\mathcal{D}}^{\text{NS}}$ are the restrictions of, respectively, the morphisms $\text{ab}^{*,\text{NS}}$ and $\text{res}_{\mathcal{D}}^{\text{NS}}$ (which are defined in Proposition 3.11), the morphisms $\nu_{G^{\text{ab}}}^{\delta^{\text{ab}},\text{NS}}$ and $\nu_G^{\delta,\text{NS}}$ are induced by the inclusions onto the second factors, the morphisms $\omega_{G^{\text{ab}}}^{\delta^{\text{ab}},\text{NS}}$ and $\omega_G^{\delta,\text{NS}}$ are induced by the projections onto the first factors.

Proof. The second row is exact by Proposition 3.11, and the third row is exact by (2.17). Moreover, the second and third rows commute: the equality $\omega_G^{\delta,\text{NS}} \circ \text{ab}^{*,\text{NS}} = \overline{\Lambda}_{\text{ab}}^* \circ \omega_{G^{\text{ab}}}^{\delta^{\text{ab}},\text{NS}}$ follows from the fact that $\text{ab}^{*,\text{NS}} = \overline{\Lambda}_{\text{ab}}^* \oplus B_{\text{ab}}^*$ (as it can be deduced from Definition/Lemma 3.10), while the equality $\text{ev}_{\mathcal{D}(G)}^\delta \circ \text{res}_{\mathcal{D}}^{\text{NS}} = \overline{\Lambda}_{\mathcal{D}}^* \circ \omega_G^{\delta,\text{NS}}$ is exactly the condition (3.24). We now conclude by applying the snake lemma, and using that the kernel of $\omega_G^{\delta,\text{NS}}$ (respectively $\omega_{G^{\text{ab}}}^{\delta^{\text{ab}},\text{NS}}$) is equal to the image of $\nu_G^{\delta,\text{NS}}$ (respectively $\nu_{G^{\text{ab}}}^{\delta^{\text{ab}},\text{NS}}$) and that $\omega_{G^{\text{ab}}}^{\delta^{\text{ab}},\text{NS}}$ is surjective. \square

We can now give an explicit description of $\text{RPic}(\mathfrak{Bun}_{G,g,n}^\delta)$, which, via the morphism $\overline{(\nu_G^\delta)^*}$, can be identified with $\ker(\omega_G^\delta) \subseteq \text{RPic}(\text{Bun}_{G,g,n}^\delta)$, see (5.12).

Theorem 5.5. *Assume that $g \geq 1$. Let G be a reductive group with maximal torus T_G and Weyl group \mathcal{W}_G , and fix $\delta \in \pi_1(G)$.*

(1) *There is an exact sequence*

$$0 \rightarrow \Lambda^*(G^{\text{ab}}) \otimes H_{g,n} \xrightarrow{j_G^\delta} \text{RPic}(\mathfrak{Bun}_{G,g,n}^\delta) \xrightarrow{\overline{\gamma}_G^\delta} \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta), \quad (5.15)$$

where \overline{j}_G^δ and $\overline{\gamma}_G^\delta$ are uniquely determined by

$$(\overline{\nu}_G^\delta)^* \circ \overline{j}_G^\delta = j_G^\delta \quad \text{and} \quad \nu_G^{\delta,\text{NS}} \circ \overline{\gamma}_G^\delta = (\omega_G^\delta \oplus \gamma_G^\delta) \circ (\overline{\nu}_G^\delta)^*,$$

see Theorem 3.12(1) and Proposition 5.4;

(2) *The image of $\overline{\gamma}_G^\delta$ is equal to*

$$\text{Im}(\overline{\gamma}_G^\delta) = \begin{cases} \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) & \text{if } n \geq 1 \\ \left\{ b \in \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) : \begin{array}{l} b(\delta \otimes x) + (g-1)b(x \otimes x) \text{ is a multiple} \\ \text{of } 2g-2, \text{ for any } x \in \Lambda(T_G) \end{array} \right\} & \text{if } n = 0. \end{cases} \quad (5.16)$$

Remark 5.6. For $n = 0$, the divisibility condition in the right-hand side (5.16) depends only on the image x^{ab} of $x \in \Lambda(T_G)$ in $\Lambda(G^{\text{ab}})$. Indeed, $b(\delta \otimes x) = b^\mathbb{Q}(\delta \otimes x^{\text{ab}})$ by (5.13), while the parity of $b(x \otimes x)$ depends solely on x^{ab} since $b|_{\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_{\mathcal{D}(G)})}$ is even by definition and $b|_{\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(G^{\text{ab}})} \equiv 0$ by the claim in the proof of Proposition 2.4.

Proof. Consider the commutative diagram (of solid arrows)

$$\begin{array}{ccc} & \text{RPic}(\mathfrak{Bun}_{G,g,n}^\delta) \xrightarrow{\overline{\gamma}_G^\delta} \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) & \\ & \downarrow (\overline{\nu}_G^\delta)^* & \downarrow \nu_G^{\delta,\text{NS}} \\ \Lambda^*(G^{\text{ab}}) \otimes H_{g,n} \xrightarrow{j_G^\delta} \text{RPic}(\text{Bun}_{G,g,n}^\delta) \xrightarrow{\omega_G^\delta \oplus \gamma_G^\delta} \text{NS}(\text{Bun}_{G,g,n}^\delta) & & \\ & \downarrow \omega_G^\delta & \downarrow \omega_G^{\delta,\text{NS}} \\ & \Lambda^*(\mathcal{Z}^\circ(G)) \xlongequal{\quad} \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} & \end{array} \quad (5.17)$$

where the horizontal row is exact by Theorem 3.12(1), the left column is exact by (5.12) and the right column is exact by Proposition 5.4.

The above diagram implies the existence of the dotted arrows \overline{j}_G^δ and $\overline{\gamma}_G^\delta$. Then, the exact sequence (5.15) follows immediately. Moreover, we have

$$\nu_G^{\delta,\text{NS}}(\text{Im}(\overline{\gamma}_G^\delta)) = \text{Im}(\omega_G^\delta \oplus \gamma_G^\delta).$$

We now conclude using the description of $\text{Im}(\omega_G^\delta \oplus \gamma_G^\delta)$ in Theorem 3.12(2). \square

We now describe the cokernel of the homomorphism ω_G^δ , which, via the obstruction morphism obs_G^δ , can be identified with the kernel of the homomorphism

$$(\nu_G^\delta)^* : H^2 \left(\mathfrak{Bun}_{G,g,n}^\delta, \mathbb{G}_m \right) \rightarrow H^2 \left(\text{Bun}_{G,g,n}^\delta, \mathbb{G}_m \right),$$

see (5.2).

Theorem 5.7. *Assume that $g \geq 1$. Let G be a reductive group with maximal torus T_G and Weyl group \mathscr{W}_G , and fix $\delta \in \pi_1(G)$.*

(1) *If $n > 0$ then the homomorphism $\overline{\Lambda}_{\mathscr{D}}^*$ of (2.17) induces an isomorphism*

$$\widetilde{\Lambda}_{\mathscr{D}}^* : \text{coker} \left(\omega_G^\delta \right) \xrightarrow{\cong} \text{coker} \left(\text{ev}_{\mathscr{D}(G)}^\delta \right), \tag{5.18}$$

where $\text{ev}_{\mathscr{D}(G)}^\delta$ is the homomorphism of Definition/Lemma 2.8;

(2) *If $n = 0$ then there exists an exact sequence*

$$\begin{aligned} 0 \rightarrow \text{coker} \left(\overline{\gamma}_G^\delta \right) \xrightarrow{\partial_G^\delta} \text{Hom} \left(\Lambda(G^{\text{ab}}), \frac{\mathbb{Z}}{(2g-2)\mathbb{Z}} \right) \\ \xrightarrow{\widetilde{\Lambda}_{\text{ab}}^*} \text{coker} \left(\omega_G^\delta \right) \xrightarrow{\widetilde{\Lambda}_{\mathscr{D}}^*} \text{coker} \left(\text{ev}_{\mathscr{D}(G)}^\delta \right) \rightarrow 0, \end{aligned} \tag{5.19}$$

where $\text{coker}(\overline{\gamma}_G^\delta) = \frac{\text{NS}(\mathfrak{Bun}_{G,g,n}^\delta)}{\text{Im}(\gamma_G^\delta)}$ is the co-kernel of the homomorphism $\overline{\gamma}_G^\delta$ defined in Theorem 5.5(1), $\widetilde{\Lambda}_{\text{ab}}^*$ and $\widetilde{\Lambda}_{\mathscr{D}}^*$ are the homomorphisms induced by, respectively, $\overline{\Lambda}_{\text{ab}}^*$ and $\overline{\Lambda}_{\mathscr{D}}^*$ of (2.17), and ∂_G^δ is defined as follows:

$$\begin{aligned} \partial_G^\delta : \text{coker} \left(\overline{\gamma}_G^\delta \right) \rightarrow \text{Hom} \left(\Lambda(G^{\text{ab}}), \frac{\mathbb{Z}}{(2g-2)\mathbb{Z}} \right) \\ [b] \mapsto \{ x \mapsto [b(\delta \otimes \tilde{x}) + (1-g)b(\tilde{x} \otimes \tilde{x})] \}, \end{aligned} \tag{5.20}$$

where $b \in \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) \subset \text{Bil}^{s, \mathscr{D}\text{-ev}}(\Lambda(T_G))^{\mathscr{W}_G}$, and $\tilde{x} \in \Lambda(T_G)$ is any lift of $x \in \Lambda(G^{\text{ab}})$.

Proof. Consider the diagram

$$\begin{array}{ccccc} \Lambda^*(G^{\text{ab}}) \otimes \widehat{H}_{g,n} & \xrightarrow{i_G^\delta} & \text{RPic}(\text{Bun}_{G,g,n}^\delta) & \xrightarrow{\gamma_G^\delta} & \text{Bil}^{s, \mathscr{D}\text{-ev}}(\Lambda(T_G))^{\mathscr{W}_G} \\ \downarrow \widehat{\omega} := \omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} \circ i_{G^{\text{ab}}}^{\delta^{\text{ab}}} & & \downarrow \omega_G^\delta & & \downarrow \text{ev}_{\mathscr{D}(G)}^\delta \circ \text{res}_{\mathscr{D}} \\ \Lambda^*(G^{\text{ab}}) & \xrightarrow{\overline{\Lambda}_{\text{ab}}^*} & \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} & \xrightarrow{\overline{\Lambda}_{\mathscr{D}}^*} & \frac{\Lambda^*(T_{\mathscr{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}})} \end{array} \tag{5.21}$$

which has exact rows by (2.17) and (3.14). Furthermore, the diagram is commutative. Indeed, the left-hand square is commutative by the functoriality of ω_G^δ with respect to the inclusion $G^{\text{ab}} \hookrightarrow G$. On the other hand, the morphism $\omega_G^\delta \oplus \gamma_G^\delta$ factors through $\text{NS}(\text{Bun}_{G,g,n}^\delta)$. Then, the commutativity of the right-hand square follows by (5.14).

Using (3.15), [14, Remark 3.5.1] and Definition/Lemma 3.7(ii), we compute

$$\begin{aligned} \widehat{\omega}(\chi \otimes (m, \zeta)) &= \omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} \left(\left(\mathcal{L}_\chi, \omega_\pi^m \left(\sum \zeta_i \sigma_i \right) \right) \right) \\ &= \omega_{G^{\text{ab}}}^{\delta^{\text{ab}}} (\langle (\chi, 0), (0, \zeta) \rangle + m \langle (\chi, 0), (\chi, 0) \rangle - 2m \mathcal{L}(\chi, 0)) \quad (5.22) \\ &= |\zeta| \cdot \chi + 2m \chi(\delta^{\text{ab}}) \chi - 2m(\chi(\delta^{\text{ab}}) + 1 - g) \chi \\ &= (|\zeta| + m(2g - 2)) \chi \quad \text{if } g \geq 2. \end{aligned}$$

And similarly we get

$$\widehat{\omega}(\chi \otimes \zeta) = |\zeta| \cdot \chi \quad \text{if } g = 1. \quad (5.23)$$

Therefore, combining (5.22) and (5.23), we deduce that

$$\text{coker}(\widehat{\omega}) = \begin{cases} 0 & \text{if } n \geq 1 \\ \frac{\Lambda^*(G^{\text{ab}})}{(2g-2)\Lambda^*(G^{\text{ab}})} = \text{Hom} \left(\Lambda(G^{\text{ab}}), \frac{\mathbb{Z}}{(2g-2)\mathbb{Z}} \right) & \text{if } n = 0. \end{cases} \quad (5.24)$$

Using this, and applying the snake lemma to (5.21), we get that:

- If $n \geq 1$ then

$$\widetilde{\Lambda}_{\mathcal{D}}^* : \text{coker}(\omega_G^\delta) \xrightarrow{\cong} \text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta) \text{ is an isomorphism,}$$

which proves (1);

- If $n = 0$ then we have an exact sequence

$$\begin{aligned} \text{RPic}(\mathfrak{Bun}_{G,g,n}^\delta) \xrightarrow{\gamma_G^\delta} \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) \xrightarrow{\partial_G^\delta} \text{Hom} \left(\Lambda(G^{\text{ab}}), \frac{\mathbb{Z}}{(2g-2)\mathbb{Z}} \right) \\ \xrightarrow{\widetilde{\Lambda}_{\text{ab}}^*} \text{coker}(\omega_G^\delta) \xrightarrow{\widetilde{\Lambda}_{\mathcal{D}}^*} \text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta), \end{aligned} \quad (5.25)$$

where $\widetilde{\partial}_G^\delta$ is the boundary homomorphism.

Part (2) follows from the exact sequence (5.25) together with the following claim.

Claim: The boundary homomorphism $\widetilde{\partial}_G^\delta$ is given by

$$\begin{aligned} \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) \longrightarrow \text{Hom} \left(\Lambda(G^{\text{ab}}), \frac{\mathbb{Z}}{(2g-2)\mathbb{Z}} \right) \\ b \mapsto \left\{ x \mapsto [b(\delta \otimes \widetilde{x}) + (1-g)b(\widetilde{x} \otimes \widetilde{x})] \right\}, \end{aligned} \quad (5.26)$$

where $\widetilde{x} \in \Lambda(T_G)$ is any lift of $x \in \Lambda(G^{\text{ab}})$.

Let us first check that formula (5.26) is a well-defined homomorphism. We recall $\text{NS}(\mathfrak{Bun}_{G,g,n}^\delta)$ is a subgroup of $\text{Bil}^{s,\mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathcal{W}G}$ (see Definition 5.3). Then, the class

$$[b(\delta \otimes \tilde{x}) + (1 - g)b(\tilde{x} \otimes \tilde{x})] \in \frac{\mathbb{Z}}{(2g - 2)\mathbb{Z}}$$

is independent from the chosen lift \tilde{x} of x since $b(\delta \otimes -)|_{\Lambda(T_{\mathcal{D}(G)})} \equiv 0$ by (5.13), while the parity of $b(\tilde{x} \otimes \tilde{x})$ depends solely on x since $b|_{\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(T_{\mathcal{D}(G)})}$ is even by Definition 5.3 and $b|_{\Lambda(T_{\mathcal{D}(G)}) \otimes \Lambda(G^{\text{ab}})} \equiv 0$ by the claim in the proof of Proposition 2.4. Moreover, the map

$$x \mapsto [b(\delta \otimes \tilde{x}) + (1 - g)b(\tilde{x} \otimes \tilde{x})]$$

is linear since $b(\delta \otimes -)$ is linear and $b((\tilde{x} + \tilde{y}) \otimes (\tilde{x} + \tilde{y})) \equiv b(\tilde{x} \otimes \tilde{x}) + b(\tilde{y} \otimes \tilde{y}) \pmod{2}$.

It remains to check that the boundary homomorphism $\widetilde{\partial}_G^\delta$ is given by the above formula (5.26). By the definition of the boundary homomorphism, for any $b \in \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) = \ker(\text{ev}_{\mathcal{D}(G)}^\delta \circ \text{res}_{\mathcal{D}})$ the image $\widetilde{\partial}_G^\delta(b)$ is equal to the class $[\mu] \in \text{coker}(\widehat{\omega})$ of any element $\mu \in \Lambda^*(G^{\text{ab}})$ such that $\overline{\Lambda}_{\text{ab}}^*(\mu) = \omega_G^\delta(\widetilde{b})$, where $\widetilde{b} \in \text{RPic}(\text{Bun}_{G,g,n}^\delta)$ is any lift of b via the surjective homomorphism γ_G^δ . Using Theorem 3.3(2) and Definition/Lemma 3.5, there are the following three possibilities:

- (a) If $b \in \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathcal{W}G} \cap \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) \subseteq \text{Bil}^{s,\mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathcal{W}G}$, then we can choose $\widetilde{b} = \tau_G^\delta(b)$. Using Definition/Lemma 3.7(i) and the hypothesis that $b(\delta \otimes -)|_{\Lambda(T_{\mathcal{D}(G)})} \equiv 0$, we compute

$$\begin{aligned} \omega_G^\delta(\tau_G^\delta(b)) &= b(\delta \otimes -) = \overline{\Lambda}_{\text{ab}}^*(b^\mathbb{Q}(\delta \otimes -)|_{\Lambda(G^{\text{ab}})}) \\ &\Rightarrow \widetilde{\partial}_G^\delta(b) = [b^\mathbb{Q}(\delta \otimes -)|_{\Lambda(G^{\text{ab}})}]. \end{aligned}$$

This shows that $\widetilde{\partial}_G^\delta(b)$ is given by formula (5.26) applied to b , noticing that the second term in (5.26) goes away because b is even;

- (b) If $b = B_{\text{ab}}^*(\chi \otimes \chi' + \chi' \otimes \chi)$ for some $\chi, \chi' \in \Lambda^*(G^{\text{ab}})$, then we can choose $\widetilde{b} = \text{ab}_\#^*(\langle \chi, \chi' \rangle)$. Using Definition/Lemma 3.7(ii), we compute

$$\begin{aligned} \omega_G^\delta(\text{ab}_\#^*(\langle \chi, \chi' \rangle)) &= \overline{\Lambda}_{\text{ab}}^*(\chi(\delta^{\text{ab}})\chi' + \chi'(\delta^{\text{ab}})\chi) \\ &\Rightarrow \widetilde{\partial}_G^\delta(b) = [\chi(\delta^{\text{ab}})\chi' + \chi'(\delta^{\text{ab}})\chi]. \end{aligned}$$

This shows that $\widetilde{\partial}_G^\delta(b)$ is given by formula (5.26) applied to $b = B_{\text{ab}}^*(\chi \otimes \chi' + \chi' \otimes \chi)$ since, for any $x \in \Lambda(G^{\text{ab}})$, we have

$$\begin{aligned} & B_{\text{ab}}^*(\chi \otimes \chi' + \chi' \otimes \chi)(\delta \otimes \widetilde{x}) + (1-g)B_{\text{ab}}^*(\chi \otimes \chi' + \chi' \otimes \chi)(\widetilde{x} \otimes \widetilde{x}) \\ &= (\chi \otimes \chi' + \chi' \otimes \chi)(\delta^{\text{ab}} \otimes x) + (1-g)(\chi \otimes \chi' + \chi' \otimes \chi)(x \otimes x) \\ &= \chi(\delta^{\text{ab}})\chi'(x) + \chi'(\delta^{\text{ab}})\chi(x) + 2(1-g)\chi(x)\chi'(x) \\ &\equiv \chi(\delta^{\text{ab}})\chi'(x) + \chi'(\delta^{\text{ab}})\chi(x) \pmod{(2g-2)}; \end{aligned}$$

(c) If $b = B_{\text{ab}}^*(\chi \otimes \chi)$ for some $\chi \in \Lambda^*(G^{\text{ab}})$, then we can choose $\widetilde{b} = \text{ab}_\#^*(\mathcal{L}(\chi))$. Using Definition/Lemma 3.7(ii), we compute

$$\omega_G^\delta(\text{ab}_\#^*(\mathcal{L}(\chi))) = (\chi(\delta^{\text{ab}}) + 1-g)\overline{\Lambda_{\text{ab}}^*}(\chi) \Rightarrow \widetilde{\partial}_G^\delta(b) = (\chi(\delta^{\text{ab}}) + 1-g)[\chi].$$

This shows that $\widetilde{\partial}_G^\delta(b)$ is given by formula (5.26) applied to $b = B_{\text{ab}}^*(\chi \otimes \chi)$ since, for any $x \in \Lambda(G^{\text{ab}})$, we have

$$\begin{aligned} & B_{\text{ab}}^*(\chi \otimes \chi)(\delta \otimes \widetilde{x}) + (1-g)B_{\text{ab}}^*(\chi \otimes \chi + \chi)(\widetilde{x} \otimes \widetilde{x}) \\ &= (\chi \otimes \chi)(\delta^{\text{ab}} \otimes x) + (1-g)(\chi \otimes \chi)(x \otimes x) \\ &= \chi(\delta^{\text{ab}})\chi(x) + (1-g)\chi(x)^2 \\ &\equiv \chi(\delta^{\text{ab}})\chi(x) + (1-g)\chi(x) \pmod{(2g-2)}. \quad \square \end{aligned}$$

Remark 5.8. The exact sequence (5.19) (if $n = 0$) is canonically isomorphic to the exact sequence

$$\begin{aligned} 0 \rightarrow \text{coker}\left(\overline{\gamma_G^\delta}\right) &\xrightarrow{\widetilde{v}_G^{\delta, \text{NS}}} \text{coker}\left(\omega_G^\delta \oplus \gamma_G^\delta\right) \\ &\xrightarrow{\widetilde{\omega}_G^{\delta, \text{NS}}} \text{coker}\left(\omega_G^\delta\right) \rightarrow \text{coker}\left(\omega_G^{\delta, \text{NS}}\right) \rightarrow 0, \end{aligned} \tag{5.27}$$

which is obtained by quotienting out the exact sequence (see Proposition 5.4)

$$0 \rightarrow \text{NS}\left(\mathfrak{Bun}_{G,g,n}^\delta\right) \xrightarrow{v_G^{\delta, \text{NS}}} \text{NS}\left(\text{Bun}_{G,g,n}^\delta\right) \xrightarrow{\omega_G^{\delta, \text{NS}}} \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} \rightarrow \text{coker}\left(\omega_G^{\delta, \text{NS}}\right) \rightarrow 0,$$

by the exact sub-sequence

$$0 \rightarrow \text{Im}\left(\overline{\gamma_G^\delta}\right) \rightarrow \text{Im}\left(\omega_G^\delta \oplus \gamma_G^\delta\right) \rightarrow \text{Im}\left(\omega_G^\delta\right) \rightarrow 0 \rightarrow 0.$$

Indeed, Proposition 5.4 provides a canonical isomorphism $\text{coker}(\omega_G^{\delta, \text{NS}}) \xrightarrow{\cong} \text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta)$ which commutes with the two surjections from $\text{coker}(\omega_G^\delta)$ in (5.19) and (5.27).

Moreover, consider the diagram

$$\begin{CD}
 \Lambda^*(G^{ab}) \otimes \widehat{H}_{g,n} @>{i_{G^{ab}}}^{\delta^{ab}}>> \text{RPic}(\text{Bun}_{G^{ab},g,n}^{\delta^{ab}}) @>{\text{ab}^{\#}}>> \text{RPic}(\text{Bun}_{G,g,n}^{\delta}) @= \text{RPic}(\text{Bun}_{G,g,n}^{\delta}) \\
 @VV\widehat{\omega}:=\omega_{G^{ab}}^{\delta^{ab}} \circ \gamma_{G^{ab}}^{\delta^{ab}}V @VV\omega_{G^{ab}}^{\delta^{ab}} \oplus \gamma_{G^{ab}}^{\delta^{ab}}V @VV\omega_G^{\delta} \oplus \gamma_G^{\delta}V @VV\omega_G^{\delta}V \\
 \Lambda^*(G^{ab}) @>{i_1}>> \text{NS}(\text{Bun}_{G^{ab},g,n}^{\delta^{ab}}) @>{\text{ab}^{*,\text{NS}}}>> \text{NS}(\text{Bun}_{G,g,n}^{\delta}) @>{\omega_G^{\delta,\text{NS}}}>> \frac{\Lambda^*(T_G)}{\Lambda^*(G^{ab})}
 \end{CD} \tag{5.28}$$

which is commutative: the left square commutes since $\text{Im}(i_{G^{ab}}^{\delta^{ab}}) = \ker(\gamma_{G^{ab}}^{\delta^{ab}})$ by Theorem 3.6; the middle square is commutative by the functoriality of $\omega_G^{\delta} \oplus \gamma_G^{\delta}$ (see Theorem 3.12(1)) and the right square commutes by the definition of $\omega_G^{\delta,\text{NS}}$.

From the commutativity of (5.28), we get the homomorphisms

$$\begin{aligned}
 \text{Hom}\left(\Lambda(G^{ab}), \frac{\mathbb{Z}}{(2g-2)\mathbb{Z}}\right) &= \text{coker}(\widehat{\omega}) \xrightarrow{\widetilde{i}_1} \text{coker}\left(\omega_{G^{ab}}^{\delta^{ab}} \oplus \gamma_{G^{ab}}^{\delta^{ab}}\right) \\
 &\xrightarrow[\cong]{\widetilde{\text{ab}^{*,\text{NS}}}} \text{coker}\left(\omega_G^{\delta} \oplus \gamma_G^{\delta}\right) \xrightarrow[\cong]{\widetilde{\omega_G^{\delta,\text{NS}}}} \text{coker}\left(\omega_G^{\delta}\right)
 \end{aligned} \tag{5.29}$$

where $\widetilde{\text{ab}^{*,\text{NS}}}$ is an isomorphism by (3.41) and \widetilde{i}_1 is an isomorphism since $\omega_{G^{ab}}^{\delta^{ab}} \oplus \gamma_{G^{ab}}^{\delta^{ab}}$ induces an isomorphism between the cokernels of $i_{G^{ab}}^{\delta^{ab}}$ and of i_1 which are both canonically isomorphic to $\text{Bil}^s(\Lambda(G^{ab}))$ (see Theorem 3.6). Moreover, the composition of all the homomorphisms of (5.29) coincides with the homomorphism $\widetilde{\Lambda_{ab}^*}$ since the composition of all the homomorphisms in the bottom row of (5.28) is equal to $\widetilde{\Lambda_{ab}^*}$.

Therefore, we deduce that there exists an isomorphism $\phi = \widetilde{\text{ab}^{*,\text{NS}}} \circ \widetilde{i}_1$ from the second term of (5.19) into the second term of (5.27), which commutes with their homomorphisms $\widetilde{\Lambda_{ab}^*}$ and $\widetilde{\omega_G^{\delta,\text{NS}}}$ onto $\text{coker}(\omega_G^{\delta})$. This implies that ϕ also commutes with the kernel homomorphisms of $\widetilde{\Lambda_{ab}^*}$ and $\widetilde{\omega_G^{\delta,\text{NS}}}$, which are, respectively, $\widetilde{\partial_G^{\delta}}$ and $\widetilde{\nu_G^{\delta,\text{NS}}}$, and this completes the proof.

Remark 5.9. Assume that $g \geq 1$. Let $G = T$ be a torus and let $d \in \Lambda(T)$. Then clearly $\text{coker}(\text{ev}_{\mathcal{D}(T)}^d) = 0$, which implies that $\text{coker}(\omega_T^d) = 0$ if $n > 0$. On the other hand, if $n = 0$, then, using the explicit basis of $\text{RPic}(\text{Bun}_{T,g,n}^d)$ given in [14, Theorem 4.0.1(2)], it is possible to check that

$$\begin{aligned}
 \text{coker}\left(\overline{\gamma_T^d}\right) &\cong \frac{\mathbb{Z}}{\text{gcd}(2g-2, \div(d)+1-g)}\mathbb{Z} \oplus \left[\frac{\mathbb{Z}}{\text{gcd}(g-1, \div(d))\mathbb{Z}} \right]^{\oplus(\dim(T)-1)} \\
 \text{coker}\left(\omega_T^d\right) &\cong \frac{\mathbb{Z}}{\text{gcd}(2g-2, \div(d)+1-g)}\mathbb{Z} \oplus \left[\frac{\mathbb{Z}}{\text{gcd}(g-1, \div(d))\mathbb{Z}} \right]^{\oplus(\dim(T)-1)},
 \end{aligned}$$

where $\div(d)$ is the divisibility of d in the lattice $\Lambda(T)$, with the convention that $\text{coker}(\overline{\gamma_T^d}) = \{0\}$ if $g = 1$ and $d = 0$ (when the above expression for $\text{coker}(\overline{\gamma_T^d})$ is not well-defined).

We end this section by describing the relative Picard group of the rigidification $\mathfrak{Bun}_{G,g,n}^\delta$ and the cokernel of the weight homomorphism wt_G^δ , in genus $g = 0$.

Remark 5.10. Assume that $g = 0$ and $n \geq 1$. Using (4.15), it can be proved that the weight homomorphism wt_G^δ is equal to the composition

$$\text{wt}_G^\delta : \text{Pic}(\text{Bun}_{G,0,n}^\delta) \rightarrow \text{RPic}(\text{Bun}_{G,0,n}^\delta) \xrightarrow{\omega_G^\delta} \Lambda^*(\mathcal{L}(G)) = \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}}} \quad (5.30)$$

$$(l_{\mathcal{L}}, b) \mapsto [l_{\mathcal{L}} \oplus b(d^{\text{ss}} \otimes -)].$$

Therefore, using the exact sequences (4.16) and (2.17) and the fact that $\omega_{G^{\text{ab}}}^\delta$ is an isomorphism, it follows that (for some, or equivalently any, lift $d^{\text{ss}} \in \Lambda(T_G)$ of the image δ^{ss} of δ in $\pi_1(G^{\text{ss}})$):

- (i) The homomorphism θ_G^δ induces an isomorphism

$$\text{RPic}(\mathfrak{Bun}_{G,0,n}^\delta) \xrightarrow{\cong} \{b \in \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathcal{W}_G} : b(d^{\text{ss}} \otimes -) \in \Lambda^*(T_{G^{\text{ad}}})\};$$

- (ii) The cokernel of ω_G^δ (and hence of the weight homomorphism wt_G^δ) is equal to the cokernel of the homomorphism

$$\left. \begin{array}{l} \{b \in \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathcal{W}_G} : \\ b^{\mathbb{Q}}(d^{\text{ss}} \otimes -) \text{ is integral on } \Lambda(T_{\mathcal{D}(G)}) \} \end{array} \right\} \rightarrow \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}}}$$

$$b \mapsto [b(d^{\text{ss}} \otimes -)].$$

6. The universal moduli space $M_{G,g,n}^{\text{ss}}$ and its divisor class group

In this section we will describe the divisor class group of the universal moduli space $M_{G,g,n}^{\delta,\text{ss}}$ of semistable G -bundles (over n -marked smooth curves of genus g) in terms of the Picard group of $\mathfrak{Bun}_{G,g,n}$. Before presenting the results, we need some preparation.

Definition 6.1.

- (i) Let $P \rightarrow C$ be a G -bundle over a k -curve C . We say that P is (semi)stable if for any reduction F to any parabolic subgroup $P \subseteq G$, we have

$$\text{deg}(\text{ad}(F)) \underset{(\leq)}{<} 0,$$

where $\text{ad}(F) := (F \times \mathfrak{p})/P$ is the adjoint bundle of F , i.e., the vector bundle on C induced by F via the adjoint representation $P \rightarrow GL(\mathfrak{p})$. We say that P is *regularly stable*, if either G is a torus or P is stable and $\text{Aut}(P) = \mathcal{L}(G)$;

- (ii) Denote by $\text{Bun}_{G,g,n}^{\delta,ss}$, respectively $\text{Bun}_{G,g,n}^{\delta,rs}$, the locus in $\text{Bun}_{G,g,n}^{\delta}$ consisting of G -bundles over families of n -marked curves of genus g whose geometric fibers are semistable, respectively regularly stable, and by $\mathfrak{Bun}_{G,g,n}^{\delta,ss}$, respectively $\mathfrak{Bun}_{G,g,n}^{\delta,rs}$, its image in $\mathfrak{Bun}_{G,g,n}^{\delta}$.

We collect in the following Proposition the properties of the loci $\text{Bun}_{G,g,n}^{\delta,ss}$ and $\text{Bun}_{G,g,n}^{\delta,rs}$ (respectively $\mathfrak{Bun}_{G,g,n}^{\delta,ss}$ and $\mathfrak{Bun}_{G,g,n}^{\delta,rs}$).

Proposition 6.2.

- (i) *The loci $\text{Bun}_{G,g,n}^{\delta,ss} \subseteq \text{Bun}_{G,g,n}^{\delta}$ and $\mathfrak{Bun}_{G,g,n}^{\delta,ss} \subseteq \mathfrak{Bun}_{G,g,n}^{\delta}$ are open (smooth) substacks of finite type over $\mathcal{M}_{g,n}$;*
- (ii) *If G is a torus then $\text{Bun}_{G,g,n}^{\delta,rs} = \text{Bun}_{G,g,n}^{\delta,ss} = \text{Bun}_{G,g,n}^{\delta}$ and $\mathfrak{Bun}_{G,g,n}^{\delta,rs} = \mathfrak{Bun}_{G,g,n}^{\delta,ss} = \mathfrak{Bun}_{G,g,n}^{\delta}$;*
- (iii) *If G is not a torus then the complements $\text{Bun}_{G,g,n}^{\delta} \setminus \text{Bun}_{G,g,n}^{\delta,ss}$ and $\mathfrak{Bun}_{G,g,n}^{\delta} \setminus \mathfrak{Bun}_{G,g,n}^{\delta,ss}$ have codimension at least g ;*
- (iv) *If G is not a torus and one of the following holds*
 - (a) $\text{char}(k) > 0$ and $g \geq 4$;
 - (b) $\text{char}(k) = 0$ and $g \geq 2$, with the exception of the case $g = 2$ and G having a non-trivial homomorphism into PGL_2 ;*then the complements $\text{Bun}_{G,g,n}^{\delta,rs} \setminus \text{Bun}_{G,g,n}^{\delta,ss}$ and $\mathfrak{Bun}_{G,g,n}^{\delta,rs} \setminus \mathfrak{Bun}_{G,g,n}^{\delta,ss}$ have codimension at least two.*

Proof. The properties for $\text{Bun}_{G,g,n}^{\delta,ss}$ have been proved in [14, Proposition 3.2.3, 3.2.5]. The properties for $\text{Bun}_{G,g,n}^{\delta,rs}$ have been proved in [5, Theorem 2.5] for $\text{char}(k) > 0$ and in [9, Theorem II.6] for $\text{char}(k) = 0$. The properties for $\mathfrak{Bun}_{G,g,n}^{\delta,ss}$ and $\mathfrak{Bun}_{G,g,n}^{\delta,rs}$ follow since $\nu_G^{\delta} : \text{Bun}_{G,g,n}^{\delta} \rightarrow \mathfrak{Bun}_{G,g,n}^{\delta}$ is a $\mathcal{L}(G)$ -gerbe. □

Corollary 6.3.

- (i) *If either G is a torus or $g \geq 2$, the restriction homomorphisms*

$$\text{Pic} \left(\text{Bun}_{G,g,n}^{\delta} \right) \rightarrow \text{Pic} \left(\text{Bun}_{G,g,n}^{\delta,ss} \right) \text{ and } \text{Pic} \left(\mathfrak{Bun}_{G,g,n}^{\delta} \right) \rightarrow \text{Pic} \left(\mathfrak{Bun}_{G,g,n}^{\delta,ss} \right)$$

are bijective;

- (ii) *If one of the following holds*
 - (a) G is a torus;
 - (b) G is not a torus, $\text{char}(k) > 0$ and $g \geq 4$;
 - (c) G is not a torus, $\text{char}(k) = 0$ and $g \geq 2$, with the exception of the case $g = 2$ and G having a non-trivial homomorphism into PGL_2 ;

then the restriction homomorphisms

$$\text{Pic} \left(\text{Bun}_{G,g,n}^{\delta,ss} \right) \rightarrow \text{Pic} \left(\text{Bun}_{G,g,n}^{\delta,rs} \right) \text{ and } \text{Pic} \left(\mathfrak{Bun}_{G,g,n}^{\delta,ss} \right) \rightarrow \text{Pic} \left(\mathfrak{Bun}_{G,g,n}^{\delta,rs} \right)$$

are bijective.

Proof. Under the assumptions in (i), Proposition 6.2 implies that either the complements $\text{Bun}_{G,g,n}^{\delta} \setminus \text{Bun}_{G,g,n}^{\delta,ss}$ and $\mathfrak{Bun}_{G,g,n}^{\delta} \setminus \mathfrak{Bun}_{G,g,n}^{\delta,ss}$ are empty or they have codimension at least two. Then the conclusion follows since $\text{Bun}_{G,g,n}^{\delta}$ (and hence $\mathfrak{Bun}_{G,g,n}^{\delta}$) is smooth by Theorem 3.1(3), see, e.g., [14, Lemma 2.3.1]. The same argument applies to point (ii). \square

We now make the following:

Assumption 6.4. There exists an adequate moduli space

$$\pi : \mathfrak{Bun}_{G,g,n}^{\delta,ss} \rightarrow M_{G,g,n}^{\delta,ss} \tag{6.1}$$

in the sense of Alper [2] (which is the same as a good moduli space if $\text{char}(k) = 0$, see [2, Proposition 5.1.4]).

Although we expect that Assumption 6.4 should always hold true, we do not know of a reference in the literature where this is proved in full generality. As far as we know, the cases covered in the literature are the following:

- $G = T$ torus, in which case π is a coarse moduli space in the sense of Keel-Mori;
- $G = GL_r$, by [25];
- $\text{char}(k) = 0$, $g \geq 2$ and $n = 0$, by [8];
- $\text{char}(k) = 0$ and $\mathcal{M}_{g,n}$ is a variety (which happens if and only if $n > 2g + 2$), by [20].

Furthermore, in these cases, the adequate moduli space is a quasi-projective variety.

Remark 6.5. In characteristic zero and $g \geq 2$, the remaining cases should follow by a slight modification of the argument in [8]. In positive characteristic over a fixed curve, the problem has been solved in [15, 16]. We are not aware, if the same results hold in the universal setting.

We are now ready to compare the divisor class group $\text{Cl}(M_{G,g,n}^{\delta,ss})$ of the algebraic space $M_{G,g,n}^{\delta,ss}$ (see [26, Tag 0EDQ]) with the Picard group $\text{Pic}(\mathfrak{Bun}_{G,g,n}^{\delta})$.

We denote by $M_{G,g,n}^{\delta,rs}$ the (open) subset $\pi(\text{Bun}_{G,g,n}^{\delta,rs}) \subset M_{G,g,n}^{\delta,ss}$. Since $\mathfrak{Bun}_{G,g,n}^{\delta}$ is smooth, the algebraic space $M_{G,g,n}^{\delta,rs}$ is normal, see [2, Proposition

5.4.1]. Then, the complement of the smooth locus $(M_{G,g,n}^{\delta,rs})_{\text{sm}}$ of $M_{G,g,n}^{\delta,rs}$ has codimension at least two. In particular, the restriction homomorphism gives an isomorphism

$$\text{Cl}\left(M_{G,g,n}^{\delta,rs}\right) \xrightarrow{\cong} \text{Cl}\left(\left(M_{G,g,n}^{\delta,rs}\right)_{\text{sm}}\right) = \text{Pic}\left(\left(M_{G,g,n}^{\delta,rs}\right)_{\text{sm}}\right).$$

By definition, the pull-back along the adequate moduli space π gives an injective homomorphism

$$\text{Pic}\left(\left(M_{G,g,n}^{\delta,rs}\right)_{\text{sm}}\right) \hookrightarrow \text{Pic}\left(\pi^{-1}\left(\left(M_{G,g,n}^{\delta,rs}\right)_{\text{sm}}\right)\right) \xrightarrow[\text{res}^{-1}]{\cong} \text{Pic}\left(\mathfrak{Bun}_{G,g,n}^{\delta,rs}\right).$$

The last isomorphism is the inverse of the restriction morphism res , which is an isomorphism since $\mathfrak{Bun}_{G,g,n}^{\delta,rs}$ is smooth and that the complement of $\pi^{-1}\left(\left(M_{G,g,n}^{\delta,rs}\right)_{\text{sm}}\right)$ has codimension at least two (since $\mathfrak{Bun}_{G,g,n}^{\delta,rs}$ has finite inertia and, so, the inverse image along π preserves the codimension). Putting all together, we get an injective homomorphism

$$\tilde{\pi}^* : \text{Cl}\left(M_{G,g,n}^{\delta,rs}\right) \hookrightarrow \text{Pic}\left(\mathfrak{Bun}_{G,g,n}^{rs}\right). \tag{6.2}$$

Theorem 6.6. *Let $g + n \geq 3$ (i.e., $\mathcal{M}_{g,n}$ is generically a variety). Suppose that Assumption 6.4 holds true and that one of the followings conditions holds:*

- (i) G is a torus;
- (ii) G is not a torus, $\text{char}(k) > 0$, $g \geq 4$;
- (iii) G is not a torus, $\text{char}(k) = 0$, $g \geq 2$, with the exception of the case $g = 2$ and G having a non-trivial homomorphism into PGL_2 .

Then we have the following isomorphisms:

$$\begin{aligned} \text{Cl}\left(M_{G,g,n}^{\delta,ss}\right) &\xrightarrow[\text{res}]{\cong} \text{Cl}\left(M_{G,g,n}^{\delta,rs}\right) \xrightarrow[\tilde{\pi}^*]{\cong} \text{Pic}\left(\mathfrak{Bun}_{G,g,n}^{\delta,rs}\right) \\ &\xrightarrow[\text{res}^{-1}]{\cong} \text{Pic}\left(\mathfrak{Bun}_{G,g,n}^{\delta,ss}\right) \xrightarrow[\text{res}^{-1}]{\cong} \text{Pic}\left(\mathfrak{Bun}_{G,g,n}^{\delta}\right) \end{aligned}$$

where res are the obvious restriction homomorphisms and $\tilde{\pi}^*$ is the homomorphism (6.2).

Proof. The last two isomorphisms follow by Corollary 6.3. Observe that

$$\text{cod}_{M_{G,g,n}^{\delta,ss}}\left(M_{G,g,n}^{\delta,ss} \setminus M_{G,g,n}^{\delta,rs}\right) \geq \text{cod}_{\text{Bun}_{G,g,n}^{\delta,ss}}\left(\text{Bun}_{G,g,n}^{\delta,ss} \setminus \text{Bun}_{G,g,n}^{\delta,rs}\right) \geq 2$$

where the last inequality follows by Proposition 6.2. In particular, the first homomorphism res is bijective.

In order to prove that $\tilde{\pi}^*$ is an isomorphism, let $\text{Bun}_{G,g,n}^{\delta,o-rs} \subset \text{Bun}_{G,g,n}^{\delta,rs}$ be the open substack whose k -points are triples $(C, \underline{\sigma}, P) \in \text{Bun}_{G,g,n}^{\delta}(k)$ such that

- $\text{Aut}(C, \underline{\sigma}) = \{1\}$;
- $\text{Aut}(P) = \mathcal{L}(G)$.

We denote by $M_{G,g,n}^{\delta,o-rs}$ the open subset $\pi(\text{Bun}_{G,g,n}^{\delta,o-rs}) \subset M_{G,g,n}^{\delta,rs}$ and by $\mathfrak{Bun}_{G,g,n}^{\delta,o-rs}$ the open substack $v_G^\delta(\text{Bun}_{G,g,n}^{\delta,o-rs}) \subset \mathfrak{Bun}_{G,g,n}^{\delta,rs}$.

We now assume $g + n > 3$. In this range, the locus of n -marked curves without non-trivial automorphisms has codimension at least two in $\mathcal{M}_{g,n}$ (see [1, Chapter XII, Proposition (2.5)]), and we deduce that

$$\begin{aligned} \text{cod}_{\mathfrak{Bun}_{G,g,n}^{\delta,rs}} \left(\mathfrak{Bun}_{G,g,n}^{\delta,rs} \setminus \mathfrak{Bun}_{G,g,n}^{\delta,o-rs} \right) &\geq 2 \\ \text{and } \text{cod}_{M_{G,g,n}^{\delta,rs}} \left(M_{G,g,n}^{\delta,rs} \setminus M_{G,g,n}^{\delta,o-rs} \right) &\geq 2. \end{aligned} \tag{6.3}$$

In particular, the restriction homomorphisms

$$\text{Pic} \left(\mathfrak{Bun}_{G,g,n}^{\delta,rs} \right) \rightarrow \text{Pic} \left(\mathfrak{Bun}_{G,g,n}^{\delta,o-rs} \right) \text{ and } \text{Cl} \left(M_{G,g,n}^{\delta,rs} \right) \rightarrow \text{Cl} \left(M_{G,g,n}^{\delta,o-rs} \right) \tag{6.4}$$

are bijective. Hence, it is enough to show that the restriction

$$\widetilde{\pi}_{|\text{Cl}(M_{G,g,n}^{\delta,o-rs})}^* : \text{Cl} \left(M_{G,g,n}^{\delta,o-rs} \right) \hookrightarrow \text{Pic} \left(\mathfrak{Bun}_{G,g,n}^{\delta,o-rs} \right)$$

is an isomorphism. This follows from the fact that the adequate moduli space (6.1) restricted to those open substacks

$$\pi|_{\mathfrak{Bun}_{G,g,n}^{\delta,o-rs}} : \mathfrak{Bun}_{G,g,n}^{\delta,o-rs} \rightarrow M_{G,g,n}^{\delta,o-rs} \tag{6.5}$$

is an isomorphism. Indeed, by definition $\text{Bun}_{G,g,n}^{\delta,o-rs}$ is an algebraic space. In particular, the adequate moduli space (6.5) is a coarse moduli space in the sense of Keel-Mori, see [2, Theorem 8.3.2]. Since the coarse moduli space is universal for maps to algebraic spaces, we must have that $\pi|_{\mathfrak{Bun}_{G,g,n}^{\delta,o-rs}}$ is an isomorphism.

It remains the case $g + n = 3$, *i.e.*, $(g, n) = (3, 0), (2, 1)$. Under these assumptions, the locus of n -marked curves with non-trivial automorphisms has one (irreducible) divisor component D (see [1, Chapter XII, Proposition (2.5)]). Since the morphism $\Phi_G^\delta : \mathfrak{Bun}_{G,g,n}^{\delta,rs} \rightarrow \mathcal{M}_{g,n}$ is smooth with irreducible fibers, we deduce that the restriction homomorphisms (6.4) are surjective and their kernels are freely generated by the irreducible divisors $(\Phi_G^\delta)^{-1}(D)$ and $\pi((\Phi_G^\delta)^{-1}(D))$, respectively. With this in mind, the theorem follows by repeating the argument of the previous case. □

7. Examples

The aim of this section is to make explicit the results of this paper for the reductive groups G such that the semisimple factor \mathfrak{g}^{ss} of the Lie algebra \mathfrak{g} of G (see (2.4)) is simple. We will therefore distinguish several cases according to the type of the

simple Lie algebra \mathfrak{g}^{ss} . For each of these cases, we will first compute the lattices of W_G -symmetric bilinear forms appearing in Proposition 2.4 and Definition/Lemma 2.11

$$\begin{aligned} \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} &\xrightarrow{r_G} \text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} \\ &\subseteq \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathscr{W}_G}, \end{aligned} \tag{7.1}$$

which have rank one by Corollary 2.7. Then we will compute, for any $\delta \in \pi_1(G)$, the cokernels of the morphisms

$$\begin{aligned} \text{ev}_{\mathcal{D}(G)}^\delta : \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} &\longrightarrow \Lambda^*(\mathcal{L}(\mathcal{D}(G))) = \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{\text{ad}})}, \\ \tilde{\text{ev}}_{\mathcal{D}(G)}^\delta : \text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} &\longrightarrow \Lambda^*(\mathcal{L}(\mathcal{D}(G))) = \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{\text{ad}})} \end{aligned}$$

appearing in Definition/Lemmas 2.8 and 2.11. Note that, since $\text{ev}_{\mathcal{D}(G)}^\delta$ is the restriction of $\tilde{\text{ev}}_{\mathcal{D}(G)}^\delta$, there is a surjection

$$\text{coker}\left(\text{ev}_{\mathcal{D}(G)}^\delta\right) \twoheadrightarrow \text{coker}\left(\tilde{\text{ev}}_{\mathcal{D}(G)}^\delta\right),$$

whose kernel is either trivial or isomorphic to $\mathbb{Z}/2\mathbb{Z}$ by Definition/Lemma 2.11(i) and the fact that the lattices (7.1) have rank one.

7.1. Type A_{n-1} ($n \geq 2$)

Let us first recall some properties of the root system A_{n-1} .

Consider the vector space \mathbb{R}^n endowed with the standard scalar product $(-, -)$ and with the canonical basis $\{\epsilon_1, \dots, \epsilon_n\}$. Consider the subvector space

$$V(A_{n-1}) := \left\{ \xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n : \sum_i \xi_i = 0 \right\} \subset \mathbb{R}^n.$$

We will freely identify $V(A_{n-1})$ with its dual vector space by means of the (restriction of the) standard scalar product $(-, -)$. The root (respectively coroot) lattice $Q(A_{n-1})$ (respectively $Q(A_{n-1}^\vee)$) and the weight (respectively coweight) lattice $P(A_{n-1})$ (respectively $P(A_{n-1}^\vee)$) of A_{n-1} are given by

$$\begin{aligned} Q(A_{n-1}) &= Q(A_{n-1}^\vee) = V(A_{n-1}) \cap \mathbb{Z}^n \subset P(A_{n-1}) \\ &= P(A_{n-1}^\vee) = V(A_{n-1}) \cap \mathbb{Z}^n + \left\langle \omega_1 := \epsilon_1 - \frac{\sum_i \epsilon_i}{n} \right\rangle. \end{aligned}$$

It follows that group $P(A_{n-1})/Q(A_{n-1})$ is cyclic of order n and it is generated by ω_1 . The Weyl group $\mathscr{W}(A_{n-1})$ of A_{n-1} is equal to S_n and it acts on the above lattices by permuting the coordinates of $V(A_{n-1}) \subset \mathbb{R}^n$.

A semisimple group H which is almost-simple of type A_{n-1} is isomorphic to SL_n / μ_r , for some (unique) $r \in \mathbb{N}$ such that $r|n$. In particular, $H = \mathrm{SL}_n$ and $H^{\mathrm{ad}} = \mathrm{PSL}_n$. By choosing the standard maximal tours T_H of H consisting of diagonal matrices, we get the canonical identifications

$$\begin{aligned} \Lambda(T_{\mathrm{SL}_n}) &= Q(A_{n-1}^\vee) \subseteq \Lambda(T_{\mathrm{SL}_n / \mu_r}) \\ &= Q(A_{n-1}^\vee) + \left\langle \frac{n}{r} \omega_1 \right\rangle \subseteq \Lambda(T_{\mathrm{PSL}_n}) = P(A_{n-1}^\vee), \\ \Lambda^*(T_{\mathrm{SL}_n}) &= P(A_{n-1}) \supseteq \Lambda^*(T_{\mathrm{SL}_n / \mu_r}) \\ &= Q(A_{n-1}) + \langle r \omega_1 \rangle \supseteq \Lambda^*(T_{\mathrm{PSL}_n}) = Q(A_{n-1}). \end{aligned} \tag{7.2}$$

It follows that the fundamental group of SL_n / μ_r is equal to

$$\pi_1(H) = \frac{\Lambda(T_{\mathrm{SL}_n / \mu_r})}{\Lambda(T_{\mathrm{SL}_n})} = \left\langle \frac{n}{r} \omega_1 \right\rangle \cong \mathbb{Z} / r\mathbb{Z}, \tag{7.3}$$

while the character group of the center $\mathcal{Z}(\mathrm{SL}_n / \mu_r) = \mu_n / \mu_r \cong \mu_{n/r}$ is equal to

$$\Lambda^*(\mathcal{Z}(\mathrm{SL}_n / \mu_r)) = \frac{\Lambda^*(T_{\mathrm{SL}_n / \mu_r})}{\Lambda^*(T_{\mathrm{PSL}_n})} = \langle r \omega_1 \rangle \cong \mathbb{Z} / \frac{n}{r} \mathbb{Z}. \tag{7.4}$$

From now on, we will consider the following

Set-up: Let G be a reductive group such that $\mathcal{D}(G) = \mathrm{SL}_n / \mu_r$ and $G^{\mathrm{ss}} = \mathrm{SL}_n / \mu_s$, with $1 \leq r|s|n$. Equivalently, G is the product of a torus and one of the reductive groups (see [7, (2.1)])

$$\begin{cases} \mathrm{SL}_n / \mu_r & \text{if } r = s \\ C_{\mu_s / \mu_r}(\mathrm{SL}_n / \mu_r) := \frac{\mathrm{SL}_n / \mu_r \times \mathbb{G}_m}{\mu_s / \mu_r} & \text{if } r \neq s, \end{cases}$$

where $\mu_s / \mu_r \cong \mu_{s/r}$ is embedded diagonally in $\mathrm{SL}_n / \mu_r \times \mathbb{G}_m$.

Lemma 7.1. *Let G be a reductive group as in the above set-up. Then we have that:*

- (i) $\mathrm{Bil}^{s, \mathrm{ev}}(\Lambda(T_G))^{\mathcal{W}_G} = \langle (-, -) \rangle$;
- (ii) $\mathrm{Bil}^{s, -\mathrm{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\mathrm{ss}}}))^{\mathcal{W}_G} = \left\langle \frac{\mathrm{lcm}(r,s,n)}{n} (-, -) \right\rangle$;
- (iii) $\mathrm{Bil}^{s, \mathrm{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\mathrm{ss}}}))^{\mathcal{W}_G} = \begin{cases} \left\langle 2 \frac{\mathrm{lcm}(r,s,n)}{n} (-, -) \right\rangle & \text{if } v_2(r) = v_2(s) \geq \frac{v_2(n)}{2} > 0 \\ \left\langle \frac{\mathrm{lcm}(r,s,n)}{n} (-, -) \right\rangle & \text{otherwise;} \end{cases}$

here $v_2(N)$ is the 2-adic valuation of a number $N \in \mathbb{N}$, i.e., $2^{v_2(N)} | N$ but $2^{v_2(N)+1} \nmid N$.

In particular,

$$\text{coker}(r_G) = \begin{cases} \mathbb{Z}/2\mathbb{Z} & \text{if } v_2(r) = v_2(s) \geq \frac{v_2(n)}{2} > 0 \\ 0 & \text{otherwise.} \end{cases}$$

Part (ii), combined with [22], recovers [7, Section 3] if $G = \text{SL}_n$ or PGL_n and [21, Theorem 5.7] if $G = \text{SL}_n / \mu_r$ for some $1 \leq r \mid n$.

Proof. First of all, the three lattices have dimension one by Corollary 2.7.

Part (i): the symmetric bilinear form $(-, -) \in \text{Bil}^s(\Lambda(T_G))$ is $\mathscr{W}(A_{n-1}) = S_n$ -invariant and it is even since $\Lambda(T_G) = Q(A_{n-1}^\vee)$ is generated by the elements $\epsilon_i - \epsilon_j$ (for $1 \leq i \neq j \leq n$) and we have that

$$(\epsilon_i - \epsilon_j, \epsilon_i - \epsilon_j) = 2.$$

This also shows that $(-, -)$ is a generator of $\text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathscr{W}_G}$ since 2 is the smallest non-zero even integer.

Part (ii): consider an element of $\text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathscr{W}_G}$, which, by part (i), is of the form $\alpha(-, -)$ for $\alpha \in \mathbb{Z}$. Using (7.2), we get that the element $\alpha(-, -)$ belongs to $\text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathscr{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G}$ if and only if

$$\mathbb{Z} \ni \alpha \left(\frac{n}{r} \omega_1, \frac{n}{s} \omega_1 \right) = \alpha \frac{n^2}{rs} (\omega_1, \omega_1) = \alpha \frac{n^2}{rs} \frac{n-1}{n} \iff \alpha \frac{n}{rs} \in \mathbb{Z} \iff \frac{\text{lcm}(rs, n)}{n} \mid \alpha,$$

where in the second equivalence we have used that r and s are coprime with $n - 1$ since they divide n by assumption.

Part (iii): since $\text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathscr{D}(G)}) \mid \Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G}$ has rank one, by Definition/Lemma 2.11(i) we have that the inclusion r_G has index at most two. Using (7.2) and part (ii), we get that r_G has index two if and only if

$$\begin{aligned} \frac{\text{lcm}(rs, n)}{n} \left(\frac{n}{r} \omega_1, \frac{n}{r} \omega_1 \right) &= \frac{\text{lcm}(rs, n)}{r^2} (n-1) \quad \text{is odd} \\ &\iff \begin{cases} n \text{ is even} \\ v_2(\text{lcm}(rs, n)) \leq 2v_2(r). \end{cases} \end{aligned}$$

The second condition is equivalent to (using that $r \mid s$)

$$\begin{aligned} \max(v_2(rs), v_2(n)) = v_2(\text{lcm}(rs, n)) &\leq 2v_2(r) \\ &\iff \begin{cases} v_2(r) + v_2(s) \leq 2v_2(r) \\ v_2(n) \leq 2v_2(r), \end{cases} \iff \begin{cases} v_2(s) = v_2(r) \\ v_2(n) \leq 2v_2(r), \end{cases} \end{aligned}$$

and this concludes the proof. □

Lemma 7.2. *Let G be a reductive group as in the above set-up. Consider an element $\delta \in \pi_1(G)$ and denote by $\Delta^{ss} \in \mathbb{Z}/s\mathbb{Z}$ the element corresponding to $\delta^{ss} \in \pi_1(G^{ss}) = \pi_1(\mathrm{SL}_n/\mu_s)$ under the isomorphism $\pi_1(\mathrm{SL}_n/\mu_s) \cong \mathbb{Z}/s\mathbb{Z}$ of (7.3). Then we have that*

- (i) $\mathrm{coker}(\tilde{\mathrm{ev}}_{\mathcal{D}(G)}^\delta) \cong \frac{\mathbb{Z}}{\mathrm{gcd}\left(\frac{\Delta^{ss} \mathrm{lcm}(rs, n)}{rs}, \frac{n}{r}\right)\mathbb{Z}};$
- (ii) $\mathrm{coker}(\mathrm{ev}_{\mathcal{D}(G)}^\delta) \cong \begin{cases} \frac{\mathbb{Z}}{\mathrm{gcd}\left(\frac{2\Delta^{ss} \mathrm{lcm}(rs, n)}{rs}, \frac{n}{r}\right)\mathbb{Z}} & \text{if } v_2(r) = v_2(s) \geq \frac{v_2(n)}{2} > 0 \\ \frac{\mathbb{Z}}{\mathrm{gcd}\left(\frac{\Delta^{ss} \mathrm{lcm}(rs, n)}{rs}, \frac{n}{r}\right)\mathbb{Z}} & \text{otherwise.} \end{cases}$

In particular,

$$\left| \mathrm{coker}\left(\mathrm{ev}_{\mathcal{D}(G)}^\delta\right) \right| = \begin{cases} \left| \mathrm{coker}\left(\tilde{\mathrm{ev}}_{\mathcal{D}(G)}^\delta\right) \right| + 1 & \text{if } \begin{cases} v_2(r) = v_2(s) \geq \frac{v_2(n)}{2} > 0 \\ v_2(\Delta^{ss}) < v_2(n) - v_2(r) \end{cases} \\ \left| \mathrm{coker}\left(\tilde{\mathrm{ev}}_{\mathcal{D}(G)}^\delta\right) \right| & \text{otherwise.} \end{cases}$$

Proof. Let us first prove part (7.2). Lemma 7.1(ii) implies that

$$\mathrm{Bil}^{s, -\mathrm{ev}}\left(\Lambda(T_{\mathcal{D}(G)}) \mid \Lambda(T_{G^{ss}})\right)^{\mathcal{W}G}$$

is freely generated by $\frac{\mathrm{lcm}(rs, n)}{n}(-, -)$. Therefore part (7.2) follows from the following

Claim: We have that

$$\tilde{\mathrm{ev}}_{\mathcal{D}(G)}^\delta\left(\frac{\mathrm{lcm}(rs, n)}{n}(-, -)\right) = \frac{\Delta^{ss} \mathrm{lcm}(rs, n)}{rs} \in \mathbb{Z}/\frac{n}{r}\mathbb{Z} \cong \Lambda^*(\mathcal{Z}(\mathcal{D}(G))),$$

where the last isomorphism follows from (7.4) together with our assumption that $\mathcal{D}(G) = \mathrm{SL}_n/\mu_r$.

Indeed, using that $\pi_1(G^{ss}) = \pi_1(\mathrm{SL}_n/\mu_s)$ is generated by $\frac{n}{s}\omega_1$ by (7.3) and $\Lambda^*(\mathcal{Z}(\mathcal{D}(G))) = \Lambda^*(\mathcal{Z}(\mathrm{SL}_n/\mu_r))$ is generated by $r\omega_1$ by (7.4), the claim follows from the obvious identity

$$\frac{\mathrm{lcm}(rs, n)}{n}\left(\Delta^{ss}\frac{n}{s}\omega_1, -\right) = \Delta^{ss}\frac{\mathrm{lcm}(rs, n)}{rs}(r\omega_1, -) \in \Lambda^*(T_{\mathrm{SL}_n/\mu_r}).$$

Part (7.2) is proved in the same way using the description of

$$\mathrm{Bil}^{s, \mathrm{ev}}\left(\Lambda(T_{\mathcal{D}(G)}) \mid \Lambda(T_{G^{ss}})\right)^{\mathcal{W}G}$$

contained in Lemma 7.1(iii).

The last assertion is straightforward. □

7.2. Types B_l and C_l (with $l \geq 2$)

Let us first recall some properties of the dual root systems B_l and C_l .

Consider the vector space $V(B_l) = V(C_l) = \mathbb{R}^l$ endowed with the standard scalar product $(-, -)$ and with the canonical basis $\{\epsilon_1, \dots, \epsilon_l\}$. We will freely identify \mathbb{R}^l with its dual vector space by means of the (restriction of the) standard scalar product $(-, -)$. The (co)root and (co)weight lattices of B_l and C_l are given by

$$Q(B_l) = Q(C_l^\vee) = \mathbb{Z}^l \subset P(B_l) = P(C_l^\vee) = \mathbb{Z}^l + \left\langle \omega_l := \frac{\sum_i \epsilon_i}{2} \right\rangle,$$

$$Q(C_l) = Q(B_l^\vee) = \left\{ \xi \in \mathbb{Z}^l : (\xi, \xi) \text{ is even} \right\} \subset P(C_l) = P(B_l^\vee) = \mathbb{Z}^l.$$

It follows that the group $P(B_l)/Q(B_l)$ is cyclic of order 2 generated by ω_l , while $P(C_l)/Q(C_l)$ is cyclic of order two generated by, say, ϵ_1 . The Weyl group of B_l and C_l is equal to $\mathcal{W}(B_l) = \mathcal{W}(C_l) = (\mathbb{Z}/2\mathbb{Z})^l \rtimes S_l$ and it acts on the above lattices in such a way that S_l permutes the coordinates of \mathbb{R}^l while $(\mathbb{Z}/2\mathbb{Z})^l$ changes the signs of all the coordinates.

A semisimple group H which is almost-simple of type B_l (respectively C_l) is isomorphic to either the (simply-connected) spin group Spin_{2l+1} (respectively the symplectic group Sp_{2l}) or the (adjoint) orthogonal group SO_{2l+1} (respectively the projective symplectic group PSp_{2l}). By choosing the standard maximal tours T_H of H consisting of diagonal matrices, we get the canonical identifications

$$\begin{aligned} \Lambda(T_{\text{Spin}_{2l+1}}) &= \Lambda^*(T_{\text{PSp}_{2l}}) = \left\{ \xi \in \mathbb{Z}^l : (\xi, \xi) \text{ is even} \right\} \subset \Lambda(T_{\text{SO}_{2l+1}}) \\ &= \Lambda^*(T_{\text{Sp}_{2l}}) = \mathbb{Z}^l, \\ \Lambda^*(T_{\text{Spin}_{2l+1}}) &= \Lambda(T_{\text{PSp}_{2l}}) = \mathbb{Z}^l + \left\langle \omega_l := \frac{\sum_i \epsilon_i}{2} \right\rangle \supset \Lambda^*(T_{\text{SO}_{2l+1}}) \\ &= \Lambda(T_{\text{Sp}_{2l}}) = \mathbb{Z}^l. \end{aligned} \tag{7.5}$$

It follows that the fundamental group of H is equal to

$$\pi_1(H) = \begin{cases} \{0\} & \text{if } H = \text{Spin}_{2l+1}, \text{Sp}_{2l} \\ \frac{\Lambda(T_{\text{SO}_{2l+1}})}{\Lambda(T_{\text{Spin}_{2l+1}})} = \langle \epsilon_1 \rangle \cong \mathbb{Z}/2\mathbb{Z} & \text{if } H = \text{SO}_{2l+1} \\ \frac{\Lambda(T_{\text{PSp}_{2l}})}{\Lambda(T_{\text{Sp}_{2l}})} = \langle \omega_l \rangle \cong \mathbb{Z}/2\mathbb{Z} & \text{if } H = \text{PSp}_{2l}, \end{cases} \tag{7.6}$$

while the character group of the center $\mathcal{Z}(H)$ is equal to

$$\Lambda^*(\mathcal{Z}(H)) = \begin{cases} \{0\} & \text{if } H = \text{SO}_{2l+1}, \text{PSp}_{2l} \\ \frac{\Lambda^*(T_{\text{Spin}_{2l+1}})}{\Lambda^*(T_{\text{SO}_{2l+1}})} = \langle \omega_l \rangle \cong \mathbb{Z}/2\mathbb{Z} & \text{if } H = \text{Spin}_{2l+1} \\ \frac{\Lambda^*(T_{\text{Sp}_{2l}})}{\Lambda^*(T_{\text{PSp}_{2l}})} = \langle \epsilon_1 \rangle \cong \mathbb{Z}/2\mathbb{Z} & \text{if } H = \text{Sp}_{2l}. \end{cases} \tag{7.7}$$

From now on, we will consider the following:

Set-up: Let G be a reductive group such that $\mathcal{D}(G)$ (or equivalently G^{ss}) is almost-simple of type B_l or C_l . Equivalently, G is the product of a torus and one of the following reductive groups (see [7, Example 2.3]):

$$\mathrm{Spin}_{2l+1}, \mathrm{SO}_{2l+1}, C_{\mu_2}(\mathrm{Spin}_{2l+1}) = \mathrm{CSpin}_{2l+1}, \mathrm{Sp}_{2l}, \mathrm{PSp}_{2l}, C_{\mu_2}(\mathrm{Sp}_{2l}) = \mathrm{CSp}_{2l},$$

where CSpin_{2l+1} is the Clifford group of order $2l + 1$ and CSp_{2l} is the group of symplectic similitudes of order $2l$.

Lemma 7.3. *Let G be a reductive group as in the above set-up. Then we have that:*

$$(i) \quad \mathrm{Bil}^{s,\mathrm{ev}}(\Lambda(T_G))^{\mathscr{W}_G} = \begin{cases} \langle(-, -) \rangle & \text{if } G = \mathrm{Spin}_{2l+1} \\ \langle 2(-, -) \rangle & \text{if } G = \mathrm{Sp}_{2l}; \end{cases}$$

$$(ii) \quad \mathrm{Bil}^{s,-\mathrm{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{ss}}))^{\mathscr{W}_G} \\ = \begin{cases} \langle(-, -) \rangle & \text{if } \mathcal{D}(G) = \mathrm{Spin}_{2l+1} \text{ or } \mathrm{SO}_{2l+1} \\ \langle 2(-, -) \rangle & \text{if either } \mathcal{D}(G) = \mathrm{Sp}_{2l} \\ & \text{or } \mathcal{D}(G) = \mathrm{PSp}_{2l} \text{ and } l \text{ is even} \\ \langle 4(-, -) \rangle & \text{if } \mathcal{D}(G) = \mathrm{PSp}_{2l} \text{ and } l \text{ is odd;} \end{cases}$$

$$(iii) \quad \mathrm{Bil}^{s,\mathrm{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{ss}}))^{\mathscr{W}_G} \\ = \begin{cases} \langle(-, -) \rangle & \text{if } \mathcal{D}(G) = \mathrm{Spin}_{2l+1} \\ \langle 2(-, -) \rangle & \text{if either } \mathcal{D}(G) = \mathrm{SO}_{2l+1} \text{ or } \mathcal{D}(G) = \mathrm{Sp}_{2l} \\ & \text{or } \mathcal{D}(G) = \mathrm{PSp}_{2l} \text{ and } 4 \mid l \\ \langle 4(-, -) \rangle & \text{if } \mathcal{D}(G) = \mathrm{PSp}_{2l} \text{ and } l \equiv 2 \pmod{4} \\ \langle 8(-, -) \rangle & \text{if } \mathcal{D}(G) = \mathrm{PSp}_{2l} \text{ and } l \text{ is odd.} \end{cases}$$

In particular,

$$\mathrm{coker}(r_G) = \begin{cases} \mathbb{Z}/2\mathbb{Z} & \text{if either } \mathcal{D}(G) = \mathrm{SO}_{2l+1} \text{ or } \mathcal{D}(G) = \mathrm{PSp}_{2l} \text{ and } 4 \nmid l \\ 0 & \text{otherwise.} \end{cases}$$

Part (ii), combined with [22], recovers [7, Section 4] if $G = \mathrm{Sp}_{2l}$ or PSp_{2l} and [7, (5.1)] if $G = \mathrm{Spin}_{2l+1}$ or SO_{2l+1} .

Proof. First of all, the three lattices have dimension one by Corollary 2.7.

Part (i): the symmetric bilinear form $(-, -) \in \mathrm{Bil}^s(\Lambda(T_G))$ is $\mathscr{W}(B_l) = \mathscr{W}(C_l)$ -invariant. From (7.5), it follows that if $G = \mathrm{Spin}_{2l+1}$ then $(-, -)$ is even and it generates $\mathrm{Bil}^{s,\mathrm{ev}}(\Lambda(T_{\mathrm{Spin}_{2l+1}}))^{\mathscr{W}(B_l)}$; while if $G = \mathrm{Sp}_{2l}$ then $2(-, -)$ is even and it generates $\mathrm{Bil}^{s,\mathrm{ev}}(\Lambda(T_{\mathrm{Sp}_{2l}}))^{\mathscr{W}(C_l)}$.

Part (ii): if $\mathcal{D}(G)$ is almost simple of type B_l , then from (7.5) it follows that $(-, -)$ is integral on $\Lambda(T_{\text{SO}_{2l+1}}) \otimes \Lambda(T_{\text{SO}_{2l+1}}) = \mathbb{Z}^l \otimes \mathbb{Z}^l$, which implies that $\text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}G} = \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathcal{W}G} = \langle(-, -)\rangle$ by part (i).

If, instead, $\mathcal{D}(G)$ is almost simple of type C_l , then from (7.5) it follows that $2(-, -)$ is always integral on $\Lambda(T_{\text{Sp}_{2l}}) \otimes \Lambda(T_{\text{PSp}_{2l}})$ while $2\alpha(-, -)$ (for $\alpha \in \mathbb{Z}$) it is integral on $\Lambda(T_{\text{PSp}_{2l}}) \otimes \Lambda(T_{\text{PSp}_{2l}})$ if and only if

$$\mathbb{Z} \ni 2\alpha \left(\frac{\sum_i \epsilon_i}{2}, \frac{\sum_i \epsilon_i}{2} \right) = 2\alpha \frac{l}{4} \iff 2 \mid \alpha \cdot l,$$

from which the conclusion follows.

Part (iii): from (7.5), it follows that $(-, -)$ is even on $\Lambda(T_{\text{Spin}_{2l+1}})$ while $\alpha(-, -)$ is even on $\Lambda(T_{\text{SO}_{2l+1}})$ if and only if α is even, which gives the conclusion if $\mathcal{D}(G)$ is almost simple of type B_l .

Again from (7.5) it follows that $2(-, -)$ is even on $\Lambda(T_{\text{Sp}_{2l}})$, while $\alpha 2(-, -)$ is even on $\Lambda(T_{\text{PSp}_{2l}})$ if and only if

$$2\mathbb{Z} \ni 2\alpha \left(\frac{\sum_i \epsilon_i}{2}, \frac{\sum_i \epsilon_i}{2} \right) = 2\alpha \frac{l}{4} \iff 4 \mid \alpha \cdot l \iff \begin{cases} 1 \mid \alpha & \text{if } 4 \mid l \\ 2 \mid \alpha & \text{if } l \equiv 2 \pmod{4} \\ 4 \mid \alpha & \text{if } 2 \nmid l, \end{cases}$$

from which the conclusion follows in the case when $\mathcal{D}(G)$ is almost simple of type C_l . □

Lemma 7.4. *Let G be a reductive group as in the above set-up. Consider an element $\delta \in \pi_1(G)$ with image $\delta^{\text{ss}} \in \pi_1(G^{\text{ss}})$. Then we have that*

$$\begin{aligned} \text{coker}(\tilde{\text{ev}}_{\mathcal{D}(G)}^\delta) &= \text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta) \\ &= \begin{cases} 0 & \text{if either } \mathcal{D}(G) = \text{SO}_{2l+1} \text{ or } \mathcal{D}(G) = \text{PSp}_{2l} \\ & \text{or } \mathcal{D}(G) = \text{Sp}_{2l} \text{ and } \delta^{\text{ss}} \neq 0 \text{ and } 2 \nmid l \\ \mathbb{Z}/2\mathbb{Z} & \text{otherwise.} \end{cases} \end{aligned}$$

The computation of $\text{coker}(\tilde{\text{ev}}_{\mathcal{D}(G)}^\delta)$ for $G = \text{Sp}_{2l}$ or PSp_{2l} can be found in [6, Section 8.2] and for $G = \text{Spin}_{2l+1}$ or SO_{2l+1} in [6, Section 8.3].

Proof. If $\mathcal{D}(G)$ is of type B_l , then $\tilde{\text{ev}}_{\mathcal{D}(G)}^\delta \equiv 0$ since

$$\text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}G} = \langle(-, -)\rangle$$

by Lemma 7.3(ii) and $(-, -)$ is integral on $\Lambda(T_{\text{SO}_{2l+1}}) \otimes \Lambda(T_{\text{SO}_{2l+1}})$ by (7.5). Therefore, the conclusion follows from (7.7).

Assume now that $\mathcal{D}(G)$ is of type C_l . The map $\tilde{ev}_{\mathcal{D}(G)}^\delta$ (and hence also $ev_{\mathcal{D}(G)}^\delta$) is obviously zero if either $\delta^{ss} = 0$ (which is always the case if $G^{ss} = Sp_{2l}$ by (7.6)) or $\mathcal{D}(G) = PSp_{2l}$. In the remaining cases,

$$Bil^{s,-ev}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{ss}}))^{/G} = Bil^{s,ev}(\Lambda(T_{\mathcal{D}(G)})) = \langle 2(-, -) \rangle$$

by Lemma 7.3 and hence

$$\tilde{ev}_{\mathcal{D}(G)}^\delta = ev_{\mathcal{D}(G)}^\delta \equiv 0 \iff \mathbb{Z} \ni 2 \left(\frac{\sum_i \epsilon_i}{2}, \frac{\sum_i \epsilon_i}{2} \right) = 2 \frac{l}{4} \iff l \text{ is even.}$$

We conclude using (7.7). □

7.3. Type D_l (with $l \geq 3$)

Let us first recall some properties of the root system D_l .

Consider the vector space \mathbb{R}^l endowed with the standard scalar product $(-, -)$ and with the canonical basis $\{\epsilon_1, \dots, \epsilon_l\}$. We will freely identify \mathbb{R}^l with its dual vector space by means of the (restriction of the) standard scalar product $(-, -)$. The root (respectively coroot) lattices and the weight (respectively coweight) lattices of D_l are given by

$$Q(D_l) = Q(D_l^\vee) = \left\{ \xi \in \mathbb{Z}^l : (\xi, \xi) \text{ is even} \right\} \subset P(D_l) = P(D_l^\vee) = \mathbb{Z}^l + \left\langle \frac{\sum_i \epsilon_i}{2} \right\rangle.$$

We set

$$\omega_l := \frac{\epsilon_1 + \dots + \epsilon_l}{2} \quad \text{and} \quad \omega_{l-1} := \frac{\epsilon_1 + \dots + \epsilon_{l-1} - \epsilon_l}{2}.$$

The group $P(D_l)/Q(D_l)$ is equal to

$$\begin{aligned} P(D_l)/Q(D_l) &= \{2\epsilon_1 = 0, \epsilon_1, \omega_l, \omega_{l-1}\} \\ &\cong \begin{cases} \mathbb{Z}/4\mathbb{Z} = \langle \omega_l \rangle = \langle \omega_{l-1} \rangle & \text{if } l \text{ is odd} \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} & \text{if } l \text{ is even.} \end{cases} \end{aligned}$$

The Weyl group of D_l is equal to

$$\mathcal{W}(D_l) = (\mathbb{Z}/2\mathbb{Z})^{l-1} \times S_l = \left\{ (\xi = (\xi_1, \dots, \xi_l), \sigma) \in (\mathbb{Z}/2\mathbb{Z})^l \times S_l : \prod_i \xi_i = 1 \right\},$$

and it acts on the above lattices in such a way that S_l permutes the coordinates of \mathbb{R}^l while $(\mathbb{Z}/2\mathbb{Z})^{l-1} \leq (\mathbb{Z}/2\mathbb{Z})^l$ changes the signs of all the coordinates.

A semisimple group H which is almost-simple of type D_l is isomorphic to either the (simply-connected) spin group $Spin_{2l}$, or the orthogonal group SO_{2l} , or

the (adjoint) projective orthogonal group PSO_{2l} or, if l is even, to that of the two semisimple groups

$$\Omega_{2l}^{\pm 1} := \text{Spin}_{2l} / \langle \omega_{l-\frac{1}{2} \pm \frac{1}{2}} \rangle.$$

Note that the two groups $\Omega_{2l}^{\pm 1}$ are (abstractly) isomorphic; the isomorphism is induced by the automorphism of the Dynkin diagram D_l that exchanges the last two nodes.

By choosing the standard maximal tours T_H of H consisting of diagonal matrices, we get the canonical identifications

$$\begin{aligned} \Lambda(T_{\text{Spin}_{2l}}) &= \left\{ \xi \in \mathbb{Z}^l : (\xi, \xi) \text{ is even} \right\} \subset \Lambda(T_{\text{SO}_{2l}}) \\ &= \mathbb{Z}^l \subset \Lambda(T_{\text{PSO}_{2l}}) = \mathbb{Z}^l + \left\langle \frac{\sum_i \epsilon_i}{2} \right\rangle, \\ \Lambda^*(T_{\text{Spin}_{2l}}) &= \mathbb{Z}^l + \left\langle \frac{\sum_i \epsilon_i}{2} \right\rangle \supset \Lambda^*(T_{\text{SO}_{2l}}) \\ &= \mathbb{Z}^l \supset \Lambda^*(T_{\text{SO}_{2l}}) = \left\{ \xi \in \mathbb{Z}^l : (\xi, \xi) \text{ is even} \right\}, \\ \Lambda\left(T_{\Omega_{2l}^{\pm 1}}\right) &= \Lambda^*\left(T_{\Omega_{2l}^{\pm 1}}\right) = \left\{ \xi \in \mathbb{Z}^l : (\xi, \xi) \text{ is even} \right\} + \left\langle \omega_{l-\frac{1}{2} \pm \frac{1}{2}} \right\rangle. \end{aligned} \tag{7.8}$$

It follows that the fundamental group of H is equal to

$$\pi_1(H) = \begin{cases} \{0\} & \text{if } H = \text{Spin}_{2l} \\ \frac{\Lambda(T_{\text{SO}_{2l}})}{\Lambda(T_{\text{Spin}_{2l}})} = \langle \epsilon_1 \rangle \cong \mathbb{Z}/2\mathbb{Z} & \text{if } H = \text{SO}_{2l} \\ \frac{\Lambda(T_{\Omega_{2l}^{\pm 1}})}{\Lambda(T_{\text{Spin}_{2l}})} = \langle \omega_{l-\frac{1}{2} \pm \frac{1}{2}} \rangle \cong \mathbb{Z}/2\mathbb{Z} & \text{if } H = \Omega_{2l}^{\pm 1} \\ \frac{\Lambda(T_{\text{PSO}_{2l}})}{\Lambda(T_{\text{Spin}_{2l}})} = \{0, \epsilon_1, \omega_l, \omega_{l-1}\} & \text{if } H = \text{PSO}_{2l}, \end{cases} \tag{7.9}$$

while the character group of the center $\mathcal{Z}(H)$ is equal to

$$\Lambda^*(\mathcal{Z}(H)) = \begin{cases} \{0\} & \text{if } H = \text{PSO}_{2l} \\ \frac{\Lambda^*(T_{\text{SO}_{2l}})}{\Lambda^*(T_{\text{PSO}_{2l}})} = \langle \epsilon_1 \rangle \cong \mathbb{Z}/2\mathbb{Z} & \text{if } H = \text{SO}_{2l} \\ \frac{\Lambda^*(T_{\Omega_{2l}^{\pm 1}})}{\Lambda^*(T_{\text{PSO}_{2l}})} = \langle \omega_{l-\frac{1}{2} \pm \frac{1}{2}} \rangle \cong \mathbb{Z}/2\mathbb{Z} & \text{if } H = \Omega_{2l}^{\pm 1} \\ \frac{\Lambda^*(T_{\text{Spin}_{2l}})}{\Lambda^*(T_{\text{PSO}_{2l}})} = \{0, \epsilon_1, \omega_l, \omega_{l-1}\} & \text{if } H = \text{Spin}_{2l}. \end{cases} \tag{7.10}$$

From now on, we will consider the following:

Set-up: Let G be a reductive group such that $\mathcal{D}(G)$ (or equivalently G^{ss}) is almost simple of type D_l . Equivalently, G is the product of a torus and one of the reductive

groups (see [7, Example 2.3])

$$\begin{cases} \text{Spin}_{2l}, \text{SO}_{2l}, \text{PSO}_{2l}, C_{\langle \epsilon_1 \rangle}(\text{Spin}_{2l}) = \text{CSpin}_{2l}, \\ C_{\mu_2}(\text{SO}_{2l}) = \text{CSO}_{2l}, C_{\mathcal{D}(\text{Spin}_{2l})}(\text{Spin}_{2l}), \\ \Omega_{2l}^{\pm}, C_{(\omega_{l-\frac{1}{2} \pm \frac{1}{2}})}(\text{Spin}_{2l}), C_{\mu_2}(\Omega_{2l}^{\pm}) \quad \text{if } l \text{ is even,} \end{cases}$$

where CSpin_{2l} is the Clifford group of order $2l$ and CSO_{2l} is the group of special orthogonal similitudes of order $2l$.

Lemma 7.5. *Let G be a reductive group as in the above set-up. Then we have that:*

(i) $\text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathcal{H}G} = \langle(-, -)\rangle;$

(ii) $\text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathcal{H}G}$

$$= \begin{cases} \langle(-, -)\rangle & \text{if either } \mathcal{D}(G) = \text{Spin}_{2l} \\ & \text{or } \mathcal{D}(G) = G^{\text{ss}} = \text{SO}_{2l} \\ & \text{or } \mathcal{D}(G) = G^{\text{ss}} = \Omega_{2l}^{\pm} \text{ and } 4 \mid l \\ \langle 2(-, -)\rangle & \text{if either } \mathcal{D}(G) = \text{SO}_{2l} \text{ and } G^{\text{ss}} = \text{PSO}_{2l} \\ & \text{or } \mathcal{D}(G) = \text{PSO}_{2l} \text{ and } 2 \mid l \\ & \text{or } \mathcal{D}(G) = G^{\text{ss}} = \Omega_{2l}^{\pm} \text{ and } l \equiv 2 \pmod{4} \\ & \text{or } \mathcal{D}(G) = \Omega_{2l}^{\pm} \text{ and } G^{\text{ss}} = \text{PSO}_{2l} \\ \langle 4(-, -)\rangle & \text{if } \mathcal{D}(G) = \text{PSO}_{2l} \text{ and } 2 \nmid l; \end{cases}$$

(iii) $\text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathcal{H}G}$

$$= \begin{cases} \langle(-, -)\rangle & \text{if } \mathcal{D}(G) = \text{Spin}_{2l} \\ \langle 2(-, -)\rangle & \text{if either } \mathcal{D}(G) = \text{SO}_{2l} \\ & \text{or } \mathcal{D}(G) = \text{PSO}_{2l} \text{ and } 4 \mid l \\ & \text{or } \mathcal{D}(G) = \Omega_{2l}^{\pm} \text{ and } 4 \mid l \\ \langle 4(-, -)\rangle & \text{if } \mathcal{D}(G) = \text{PSO}_{2l} \text{ and } l \equiv 2 \pmod{4} \\ & \text{or } \mathcal{D}(G) = \Omega_{2l}^{\pm} \text{ and } l \equiv 2 \pmod{4} \\ \langle 8(-, -)\rangle & \text{if } \mathcal{D}(G) = \text{PSO}_{2l} \text{ and } 2 \nmid l. \end{cases}$$

In particular,

$$\text{coker}(r_G) = \begin{cases} \mathbb{Z}/2\mathbb{Z} & \text{if either } \mathcal{D}(G) = G^{\text{ss}} = \text{SO}_{2l} \\ & \text{or } \mathcal{D}(G) = G^{\text{ss}} = \Omega_{2l}^{\pm} \text{ and } 4 \mid l \\ & \text{or } \mathcal{D}(G) = \Omega_{2l}^{\pm} \text{ and } l \equiv 2 \pmod{4} \\ & \text{or } \mathcal{D}(G) = \text{PSO}_{2l} \text{ and } 4 \nmid l \\ 0 & \text{otherwise.} \end{cases}$$

Part (ii), combined with [22], recovers [7, Section 5] if $G = \text{Spin}_{2l}$ or SO_{2l} or PSO_{2l} .

Proof. First of all, the three lattices have dimension one by Corollary 2.7.

Part (i): the symmetric bilinear form $(-, -) \in \text{Bil}^s(\Lambda(T_{\text{Spin}_{2l}}))$ is $\mathscr{W}(D_l)$ -invariant and, from (7.5), it follows that $(-, -)$ is even and generates

$$\text{Bil}^{s,\text{ev}}(\Lambda(T_{\text{Spin}_{2l}}))^{\mathscr{W}(D_l)}.$$

Part (ii): from (7.5), we get that (for any α):

- $(-, -)$ is integral on $\Lambda(T_{\text{Spin}_{2l}}) \otimes \Lambda(T_{\text{PSO}_{2l}})$ and on $\Lambda(T_{\text{SO}_{2l}}) \otimes \Lambda(T_{\text{SO}_{2l}})$;
- $\alpha(-, -)$ is integral on $\Lambda(T_{\text{SO}_{2l}}) \otimes \Lambda(T_{\text{PSO}_{2l}})$ if and only if $2 \mid \alpha$;
- $\alpha(-, -)$ is integral on $\Lambda(T_{\text{PSO}_{2l}}) \otimes \Lambda(T_{\text{PSO}_{2l}})$ if and only if

$$\mathbb{Z} \ni \alpha \left(\epsilon_1, \frac{\sum_i \epsilon_i}{2} \right) = \frac{\alpha}{2} \text{ and } \mathbb{Z} \ni \alpha \left(\frac{\sum_i \epsilon_i}{2}, \frac{\sum_i \epsilon_i}{2} \right) = \frac{\alpha \cdot l}{4} \iff \begin{cases} 2 \mid \alpha & \text{if } 2 \mid l \\ 4 \mid \alpha & \text{if } 2 \nmid l; \end{cases}$$

- $\alpha(-, -)$ is integral on $\Lambda(T_{\Omega_{2l}^\pm}) \otimes \Lambda(T_{\Omega_{2l}^\pm})$ if and only if

$$\mathbb{Z} \ni \alpha \left(\omega_{l-\frac{1}{2} \pm \frac{1}{2}}, \omega_{l-\frac{1}{2} \pm \frac{1}{2}} \right) = \frac{\alpha \cdot l}{4} \iff \begin{cases} 1 \mid \alpha & \text{if } 4 \mid l \\ 2 \mid \alpha & \text{if } l \equiv 2 \pmod{4}; \end{cases}$$

- $\alpha(-, -)$ is integral on $\Lambda(T_{\Omega_{2l}^\pm}) \otimes \Lambda(T_{\text{PSO}_{2l}})$ if and only if

$$\mathbb{Z} \ni \alpha \left(\epsilon_1, \omega_{l-\frac{1}{2} \pm \frac{1}{2}} \right) = \frac{\alpha}{2} \text{ and } \mathbb{Z} \ni \alpha \left(\omega_{l-\frac{1}{2} \pm \frac{1}{2}}, \omega_{l-\frac{1}{2} \pm \frac{1}{2}} \right) = \frac{\alpha \cdot l}{4} \iff 2 \mid \alpha.$$

Combining the above equivalences, part (ii) follows.

Part (iii): from (7.5), we get that (for any α):

- $(-, -)$ is even on $\Lambda(T_{\text{Spin}_{2l}}) \otimes \Lambda(T_{\text{PSO}_{2l}})$;
- $\alpha(-, -)$ is even on $\Lambda(T_{\text{SO}_{2l}}) \otimes \Lambda(T_{\text{SO}_{2l}})$ if and only if $2 \mid \alpha$;
- $\alpha(-, -)$ is even on $\Lambda(T_{\text{PSO}_{2l}}) \otimes \Lambda(T_{\text{PSO}_{2l}})$ if and only if

$$2\mathbb{Z} \ni \alpha(\epsilon_1, \epsilon_1) = \alpha \quad \text{and}$$

$$2\mathbb{Z} \ni \alpha \left(\frac{\sum_i \epsilon_i}{2}, \frac{\sum_i \epsilon_i}{2} \right) = \frac{\alpha \cdot l}{4} \iff \begin{cases} 2 \mid \alpha & \text{if } 4 \mid l \\ 4 \mid \alpha & \text{if } l \equiv 2 \pmod{4} \\ 8 \mid \alpha & \text{if } 2 \nmid l; \end{cases}$$

- $\alpha(-, -)$ is even on $\Lambda(T_{\Omega_{2l}^\pm}) \otimes \Lambda(T_{\Omega_{2l}^\pm})$ if and only if

$$2\mathbb{Z} \ni \alpha \left(\frac{\sum_i \epsilon_i}{2}, \frac{\sum_i \epsilon_i}{2} \right) = \frac{\alpha \cdot l}{4} \iff \begin{cases} 2 \mid \alpha & \text{if } 4 \mid l \\ 4 \mid \alpha & \text{if } l \equiv 2 \pmod{4}. \end{cases}$$

Combining the above equivalences with part (ii), we get (iii). □

Lemma 7.6. *Let G be a reductive group as in the above set-up. Consider an element $\delta \in \pi_1(G)$ and denote by δ^{ss} its image in $\pi_1(G^{ss})$. Then we have that:*

$$(i) \operatorname{coker}(\tilde{\operatorname{ev}}_{\mathcal{D}(G)}^\delta) \cong \begin{cases} \mathbb{Z}/4\mathbb{Z} & \text{if } \mathcal{D}(G) = \operatorname{Spin}_{2l}, 2 \nmid l, \delta^{ss} = 0 \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} & \text{if } \mathcal{D}(G) = \operatorname{Spin}_{2l}, 2 \mid l, \delta^{ss} = 0 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } \mathcal{D}(G) = \operatorname{Spin}_{2l}, \operatorname{ord}(\delta^{ss}) = 2 \\ & \text{or } \mathcal{D}(G) = G^{ss} = \operatorname{SO}_{2l}, \delta^{ss} = 0 \\ & \text{or } \mathcal{D}(G) = \operatorname{SO}_{2l}, G^{ss} = \operatorname{PSO}_{2l} \\ & \quad \operatorname{ord}(\delta^{ss}) \neq 4 \\ & \text{or } \mathcal{D}(G) = G^{ss} = \Omega_{2l}^\pm, \delta^{ss} = 0 \\ & \text{or } \mathcal{D}(G) = \Omega_{2l}^\pm, G^{ss} = \operatorname{PSO}_{2l} \\ & \text{otherwise,} \\ 0 & \end{cases}$$

where $\operatorname{ord}(\delta^{ss})$ is the order of δ^{ss} inside the group $\pi_1(G^{ss})$;

$$(ii) \operatorname{coker}(\operatorname{ev}_{\mathcal{D}(G)}^\delta) \cong \begin{cases} \mathbb{Z}/4\mathbb{Z} & \text{if } \mathcal{D}(G) = \operatorname{Spin}_{2l}, 2 \nmid l, \delta^{ss} = 0 \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} & \text{if } \mathcal{D}(G) = \operatorname{Spin}_{2l}, 2 \mid l, \delta^{ss} = 0 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } \mathcal{D}(G) = \operatorname{Spin}_{2l}, \operatorname{ord}(\delta^{ss}) = 2 \\ & \text{or } \mathcal{D}(G) = G^{ss} = \operatorname{SO}_{2l} \\ & \text{or } \mathcal{D}(G) = \operatorname{SO}_{2l}, G^{ss} = \operatorname{PSO}_{2l} \\ & \quad \operatorname{ord}(\delta^{ss}) \neq 4 \\ & \text{or } \mathcal{D}(G) = G^{ss} = \Omega_{2l}^\pm \\ & \text{or } \mathcal{D}(G) = \Omega_{2l}^\pm, G^{ss} = \operatorname{PSO}_{2l} \\ & \text{otherwise.} \\ 0 & \end{cases}$$

In particular,

$$\left| \operatorname{coker}(\operatorname{ev}_{\mathcal{D}(G)}^\delta) \right| = \begin{cases} \left| \operatorname{coker}(\tilde{\operatorname{ev}}_{\mathcal{D}(G)}^\delta) \right| + 1 & \text{if } \mathcal{D}(G) = G^{ss} = \operatorname{SO}_{2l} \\ & \text{and } \delta^{ss} \neq 0 \\ & \text{or } \mathcal{D}(G) = G^{ss} = \Omega_{2l}^\pm \\ & \text{and } \delta^{ss} \neq 0 \\ \left| \operatorname{coker}(\tilde{\operatorname{ev}}_{\mathcal{D}(G)}^\delta) \right| & \text{otherwise.} \end{cases}$$

The computation of $\operatorname{coker}(\tilde{\operatorname{ev}}_{\mathcal{D}(G)}^\delta)$ for $G = \operatorname{Spin}_{2l}$ or SO_{2l} or PSO_{2l} or Ω_{2l}^\pm can be found in [6, Section 8.3].

Proof. We will distinguish several cases according to what $\mathcal{D}(G)$ and G^{ss} are. We will freely use (7.9) and (7.10).

(a) $\mathcal{D}(G) = \operatorname{PSO}_{2l}$. In this case, the codomains of $\tilde{\operatorname{ev}}_{\mathcal{D}(G)}^\delta$ and of $\operatorname{ev}_{\mathcal{D}(G)}^\delta$ are zero, hence their cokernels are zero.

(b) $\mathcal{D}(G) = G^{ss} = \operatorname{SO}_{2l}$. In this case, we have that

$$\Lambda^*(\mathcal{L}(\mathcal{D}(G))) = \{0, \epsilon_1\} \cong \mathbb{Z}/2\mathbb{Z} \quad \text{and} \quad \pi_1(G^{ss}) = \{0, \epsilon_1\} \cong \mathbb{Z}/2\mathbb{Z}.$$

Since $\text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}(G)}$ is generated by $(-, -)$ by Lemma 7.5(ii), we have that

$$\widetilde{\text{ev}}_{\mathcal{D}(G)}^{\delta}((-, -)) = \begin{cases} \epsilon_1 & \text{if } \delta^{\text{ss}} = \epsilon_1 \\ 0 & \text{if } \delta^{\text{ss}} = 0 \end{cases} \implies \text{coker}(\widetilde{\text{ev}}_{\mathcal{D}(G)}^{\delta}) = \begin{cases} 0 & \text{if } \delta^{\text{ss}} \neq 0 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } \delta^{\text{ss}} = 0. \end{cases}$$

On the other hand, since $\text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}(G)}$ is generated by $2(-, -)$ by Lemma 7.5(iii), the map $\text{ev}_{\mathcal{D}(G)}^{\delta}$ is zero for any δ , which implies that $\text{coker}(\text{ev}_{\mathcal{D}(G)}^{\delta}) = \mathbb{Z}/2\mathbb{Z}$.

(c) $\mathcal{D}(G) = \text{SO}_{2l}$ and $G^{\text{ss}} = \text{PSO}_{2l}$. In this case we have that

$$\Lambda^*(\mathcal{L}(\mathcal{D}(G))) = \{0, \epsilon_1\} \cong \mathbb{Z}/2\mathbb{Z} \quad \text{and}$$

$$\pi_1(G^{\text{ss}}) = \{0, \epsilon_1, \omega_l, \omega_{l-1}\} \cong \begin{cases} \mathbb{Z}/4\mathbb{Z} & \text{if } l \text{ is odd} \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} & \text{if } l \text{ is even.} \end{cases}$$

Since $\text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}(G)} = \text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}(G)}$ is generated by $2(-, -)$ by Lemma 7.5, we have that

$$\widetilde{\text{ev}}_{\mathcal{D}(G)}^{\delta}(2(-, -)) = \text{ev}_{\mathcal{D}(G)}^{\delta}(2(-, -)) = 2\delta^{\text{ss}} = \begin{cases} 0 & \text{if } \text{ord}(\delta^{\text{ss}}) \neq 4 \\ \epsilon_1 & \text{if } \text{ord}(\delta^{\text{ss}}) = 4. \end{cases}$$

Hence we get that

$$\text{coker}(\widetilde{\text{ev}}_{\mathcal{D}(G)}^{\delta}) = \text{coker}(\text{ev}_{\mathcal{D}(G)}^{\delta}) = \begin{cases} 0 & \text{if } \text{ord}(\delta^{\text{ss}}) = 4 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } \text{ord}(\delta^{\text{ss}}) \neq 4. \end{cases}$$

(d) $\mathcal{D}(G) = G^{\text{ss}} = \Omega_{2l}^{\pm}$ (and l is even). In this case, we have that

$$\Lambda^*(\mathcal{L}(\mathcal{D}(G))) = \{0, \omega_{l-\frac{1}{2} \pm \frac{1}{2}}\} \cong \mathbb{Z}/2\mathbb{Z} \quad \text{and}$$

$$\pi_1(G^{\text{ss}}) = \{0, \omega_{l-\frac{1}{2} \pm \frac{1}{2}}\} \cong \mathbb{Z}/2\mathbb{Z}.$$

Since $\text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}(G)}$ is generated by $(-, -)$ by Lemma 7.5(ii), we have that

$$\widetilde{\text{ev}}_{\mathcal{D}(G)}^{\delta}((-, -)) = \begin{cases} \omega_{l-\frac{1}{2} \pm \frac{1}{2}} & \text{if } \delta^{\text{ss}} = \omega_{l-\frac{1}{2} \pm \frac{1}{2}} \\ 0 & \text{if } \delta^{\text{ss}} = 0 \end{cases}$$

$$\implies \text{coker}(\widetilde{\text{ev}}_{\mathcal{D}(G)}^{\delta}) = \begin{cases} 0 & \text{if } \delta^{\text{ss}} \neq 0 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } \delta^{\text{ss}} = 0. \end{cases}$$

On the other hand, since $\text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}(G)}$ is generated by $2(-, -)$ by Lemma 7.5(iii), the map $\text{ev}_{\mathcal{D}(G)}^\delta$ is zero for any δ , which implies that $\text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta) = \mathbb{Z}/2\mathbb{Z}$.

(e) $\mathcal{D}(G) = \Omega_{2l}^\pm$ and $G^{\text{ss}} = \text{PSO}_{2l}$ (and l is even). In this case, we have that

$$\Lambda^*(\mathcal{L}(\mathcal{D}(G))) = \left\{0, \omega_{l-\frac{1}{2} \pm \frac{1}{2}}\right\} \cong \mathbb{Z}/2\mathbb{Z} \quad \text{and} \quad \pi_1(G^{\text{ss}}) = \{0, \epsilon_1, \omega_l, \omega_{l-1}\}.$$

Since $\text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}(G)}$ is generated by $2(-, -)$ by Lemma 7.5(ii), we have that $\tilde{\text{ev}}_{\mathcal{D}(G)}^\delta \equiv 0$, which implies that

$$\text{coker}(\tilde{\text{ev}}_{\mathcal{D}(G)}^\delta) = \text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta) \cong \mathbb{Z}/2\mathbb{Z}.$$

(f) $\mathcal{D}(G) = \text{Spin}_{2l}$. In this case we have that

$$\begin{aligned} \Lambda^*(\mathcal{L}(\mathcal{D}(G))) &= \Lambda^*(\mathcal{L}(\text{Spin}_{2l})) = \{0, \epsilon_1, \omega_l, \omega_{l-1}\} \\ &\cong \begin{cases} \mathbb{Z}/4\mathbb{Z} & \text{if } l \text{ is odd} \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} & \text{if } l \text{ is even,} \end{cases} \end{aligned}$$

where $\pi_1(G^{\text{ss}})$ is a subgroup of $\pi_1(G^{\text{ad}}) = \pi_1(\text{PSO}_{2l})$, which is canonically isomorphic to $\Lambda^*(\mathcal{L}(\text{Spin}_{2l}))$ via the standard scalar product $(-, -)$. Since $\text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}(G)} = \text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}(G)}$ is generated by $(-, -)$ by Lemma 7.5, we have that

$$\begin{aligned} \tilde{\text{ev}}_{\mathcal{D}(G)}^\delta((-, -)) &= \text{ev}_{\mathcal{D}(G)}^\delta((-, -)) = \delta^{\text{ss}} \in \pi_1(G^{\text{ss}}) \\ &\subseteq \pi_1(\text{PSO}_{2l}) \cong \Lambda^*(\mathcal{L}(\text{Spin}_{2l})). \end{aligned}$$

Hence we deduce that

$$\begin{aligned} \text{coker}(\tilde{\text{ev}}_{\mathcal{D}(G)}^\delta) &= \text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta) \\ &= \begin{cases} \mathbb{Z}/4\mathbb{Z} & \text{if } 2 \nmid l, \delta^{\text{ss}} = 0 \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} & \text{if } 2 \mid l, \delta^{\text{ss}} = 0 \\ \mathbb{Z}/2\mathbb{Z} & \text{if } \text{ord}(\delta^{\text{ss}}) = 2 \\ 0 & \text{if } \text{ord}(\delta^{\text{ss}}) = 4 \text{ (which can occur only if } 2 \nmid l). \end{cases} \quad \square \end{aligned}$$

7.4. Types E_6, E_7 and E_8

Let us first recall some properties of the root systems E_6, E_7 and E_8 .

Consider the vector space \mathbb{R}^8 endowed with the standard scalar product $(-, -)$ and with the canonical basis $\{\epsilon_1, \dots, \epsilon_l\}$. Inside \mathbb{R}^8 , we will consider the following subvector spaces:

$$\begin{aligned} V(E_7) &:= \{\xi = (\xi_1, \dots, \xi_8) \in \mathbb{R}^8 : \xi_8 = -\xi_7\}, \\ V(E_6) &:= \{\xi = (\xi_1, \dots, \xi_8) \in \mathbb{R}^8 : \xi_8 = -\xi_6, \xi_7 = \xi_6\}. \end{aligned}$$

We will freely identify $\mathbb{R}^8, V(E_7)$ and $V(E_6)$ with its dual vector spaces by means of the (restriction of the) standard scalar product $(-, -)$. The root (respectively coroot) lattices and the weight (respectively coweight) lattices of E_8, E_7 and E_6 are given by

$$\begin{aligned} Q(E_8) &= Q(E_8^\vee) = \{\xi \in \mathbb{Z}^l : (\xi, \xi) \text{ is even}\} + \left\langle \frac{\sum_i \epsilon_i}{2} \right\rangle \\ &= P(E_8) = P(E_8^\vee), \\ Q(E_7) &= Q(E_7^\vee) = Q(E_8) \cap V(E_7) \subset P(E_7) \\ &= P(E_7^\vee) = Q(E_7) + \left\langle \omega_7 := \frac{2\epsilon_6 + \epsilon_7 - \epsilon_8}{2} \right\rangle, \\ Q(E_6) &= Q(E_6^\vee) = Q(E_8) \cap V(E_6) \subset P(E_6) \\ &= P(E_6^\vee) = Q(E_6) + \left\langle \omega_1 := \frac{2}{3}(\epsilon_8 - \epsilon_7 - \epsilon_6) \right\rangle. \end{aligned} \tag{7.11}$$

In particular, the group $P(E_*)/Q(E_*)$ is equal to

$$\begin{aligned} P(E_8)/Q(E_8) &= \{0\}, \\ P(E_7)/Q(E_7) &= \langle \omega_7 \rangle \cong \mathbb{Z}/2\mathbb{Z}, \\ P(E_6)/Q(E_6) &= \langle \omega_1 \rangle \cong \mathbb{Z}/3\mathbb{Z}. \end{aligned}$$

Explicit bases of the lattices $Q(E_8), Q(E_7)$ and $Q(E_6)$ are given by

$$\begin{aligned} Q(E_8) &= \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7, \alpha_8 \rangle, \\ Q(E_7) &= \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7 \rangle, \\ Q(E_6) &= \langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6 \rangle, \end{aligned} \tag{7.12}$$

where the elements α_i are given by

$$\begin{aligned} \alpha_1 &= \frac{\epsilon_1 - \epsilon_2 - \epsilon_3 - \epsilon_4 - \epsilon_5 - \epsilon_6 - \epsilon_7 + \epsilon_8}{2}, \\ \alpha_2 &= \epsilon_1 + \epsilon_2, \alpha_3 = \epsilon_2 - \epsilon_1, \alpha_4 = \epsilon_3 - \epsilon_2, \\ \alpha_5 &= \epsilon_4 - \epsilon_3, \alpha_6 = \epsilon_5 - \epsilon_4, \alpha_7 = \epsilon_6 - \epsilon_5, \alpha_8 = \epsilon_7 - \epsilon_6. \end{aligned}$$

From (7.11), it follows that the semisimple algebraic groups which are almost-simple of type E_* are the following ones: a simply-connected and adjoint group $\mathbb{E}_8 = \mathbb{E}_8 = \mathbb{E}_8^{\text{ad}}$ of type E_8 , a simply-connected group \mathbb{E}_7 (respectively \mathbb{E}_6) of type E_7 (respectively E_6), an adjoint group \mathbb{E}_7^{ad} (respectively \mathbb{E}_6^{ad}) of type E_7 (respectively E_6). The (co)character lattices of the maximal tori of the above semisimple groups are therefore equal to

$$\begin{aligned} \Lambda(T_{\mathbb{E}_8}) &= \Lambda^*(T_{\mathbb{E}_8}) = Q(E_8), \\ \Lambda(T_{\mathbb{E}_7}) &= \Lambda^*(T_{\mathbb{E}_7^{\text{ad}}}) = Q(E_7) \subset \Lambda(T_{\mathbb{E}_7^{\text{ad}}}) \\ &= \Lambda^*(T_{\mathbb{E}_7}) = P(E_7) = Q(E_7) + \left\langle \omega_7 := \frac{2\epsilon_6 + \epsilon_7 - \epsilon_8}{2} \right\rangle, \\ \Lambda(T_{\mathbb{E}_6}) &= \Lambda^*(T_{\mathbb{E}_6^{\text{ad}}}) = Q(E_6) \subset \Lambda(T_{\mathbb{E}_6^{\text{ad}}}) \\ &= \Lambda^*(T_{\mathbb{E}_6}) = P(E_6) + \left\langle \omega_1 := \frac{2}{3}(\epsilon_8 - \epsilon_7 - \epsilon_6) \right\rangle. \end{aligned} \tag{7.13}$$

It follows that the fundamental group of a semisimple group H as above is equal to

$$\pi_1(H) = \begin{cases} \{0\} & \text{if } H = \mathbb{E}_8, \mathbb{E}_7, \mathbb{E}_6 \\ \frac{\Lambda(T_{\mathbb{E}_7^{\text{ad}}})}{\Lambda(T_{\mathbb{E}_7})} = \left\langle \omega_7 := \frac{2\epsilon_6 + \epsilon_7 - \epsilon_8}{2} \right\rangle \cong \mathbb{Z}/2\mathbb{Z} & \text{if } H = \mathbb{E}_7^{\text{ad}} \\ \frac{\Lambda(T_{\mathbb{E}_6^{\text{ad}}})}{\Lambda(T_{\mathbb{E}_6})} = \left\langle \omega_1 := \frac{2}{3}(\epsilon_8 - \epsilon_7 - \epsilon_6) \right\rangle \cong \mathbb{Z}/3\mathbb{Z} & \text{if } H = \mathbb{E}_6^{\text{ad}}, \end{cases} \tag{7.14}$$

while the character group of the center $\mathcal{Z}(H)$ is equal to

$$\begin{aligned} &\Lambda^*(\mathcal{Z}(H)) \\ &= \begin{cases} \{0\} & \text{if } H = \mathbb{E}_8, \mathbb{E}_7^{\text{ad}}, \mathbb{E}_6^{\text{ad}} \\ \frac{\Lambda^*(T_{\mathbb{E}_7})}{\Lambda^*(T_{\mathbb{E}_7^{\text{ad}}})} = \left\langle \omega_7 := \frac{2\epsilon_6 + \epsilon_7 - \epsilon_8}{2} \right\rangle \cong \mathbb{Z}/2\mathbb{Z} & \text{if } H = \mathbb{E}_7 \\ \frac{\Lambda^*(T_{\mathbb{E}_6})}{\Lambda^*(T_{\mathbb{E}_6^{\text{ad}}})} = \left\langle \omega_1 := \frac{2}{3}(\epsilon_8 - \epsilon_7 - \epsilon_6) \right\rangle \cong \mathbb{Z}/3\mathbb{Z} & \text{if } H = \mathbb{E}_6. \end{cases} \end{aligned} \tag{7.15}$$

From now on, we will consider the following:

Set-up: Let G be a reductive group such that $\mathcal{D}(G)$ (or equivalently G^{ss}) is almost-simple of type E_* . Equivalently, G is the product of a torus and one of the following reductive groups (see [7, Example 2.3])

$$\mathbb{E}_8, \mathbb{E}_7, \mathbb{E}_7^{\text{ad}}, C_{\mu_2}(\mathbb{E}_7), \mathbb{E}_6, \mathbb{E}_6^{\text{ad}}, C_{\mu_3}(\mathbb{E}_6).$$

Lemma 7.7. *Let G be a reductive group as in the above set-up. Then we have that:*

- (i) $\text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathscr{W}G} = \langle(-, -)\rangle;$
- (ii) $\text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathscr{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}G}$
 $= \begin{cases} \langle(-, -)\rangle & \text{if } \mathscr{D}(G) = \mathbb{E}_8 \text{ or } \mathscr{D}(G) = \mathbb{E}_7 \text{ or } \mathscr{D}(G) = \mathbb{E}_6 \\ \langle 2(-, -)\rangle & \text{if } \mathscr{D}(G) = \mathbb{E}_7^{\text{ad}} \\ \langle 3(-, -)\rangle & \text{if } \mathscr{D}(G) = \mathbb{E}_6^{\text{ad}}; \end{cases}$
- (iii) $\text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathscr{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}G}$
 $= \begin{cases} \langle(-, -)\rangle & \text{if } \mathscr{D}(G) = \mathbb{E}_8 \text{ or } \mathscr{D}(G) = \mathbb{E}_7 \text{ or } \mathscr{D}(G) = \mathbb{E}_6 \\ \langle 4(-, -)\rangle & \text{if } \mathscr{D}(G) = \mathbb{E}_7^{\text{ad}} \\ \langle 3(-, -)\rangle & \text{if } \mathscr{D}(G) = \mathbb{E}_6^{\text{ad}}. \end{cases}$

In particular,

$$\text{coker}(r_G) = \begin{cases} \mathbb{Z}/2\mathbb{Z} & \text{if } \mathscr{D}(G) = \mathbb{E}_7^{\text{ad}} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. First of all, the three lattices have dimension one by Corollary 2.7.

Part (i): the symmetric bilinear form $(-, -) \in \text{Bil}^s(\Lambda(T_{\mathbb{E}_*}))$ is $\mathscr{W}(E_*)$ -invariant and, from (7.12), it follows that $(-, -)$ is even and it generates $\text{Bil}^s(\Lambda(T_{\mathbb{E}_*}))^{\mathscr{W}(E_*)}$.

Parts (ii) and (iii) for $\mathscr{D}(G) = \mathbb{E}_8, \mathbb{E}_7, \mathbb{E}_6$ follow from part (7.7) and Lemma 2.9.

Parts (ii) and (iii) for $\mathscr{D}(G) = \mathbb{E}_7^{\text{ad}}$ follows, using (7.13), from

$$\alpha(-, -) \text{ is integral on } \Lambda(T_{\mathbb{E}_7^{\text{ad}}}) \otimes \Lambda(T_{\mathbb{E}_7^{\text{ad}}}) \iff \mathbb{Z} \ni \alpha\langle\omega_7, \omega_7\rangle = \frac{3\alpha}{2} \iff 2 \mid \alpha,$$

$$\alpha(-, -) \text{ is even on } \Lambda(T_{\mathbb{E}_7^{\text{ad}}}) \otimes \Lambda(T_{\mathbb{E}_7^{\text{ad}}}) \iff \mathbb{Z} \ni \alpha\langle\omega_7, \omega_7\rangle = \frac{3\alpha}{2} \iff 4 \mid \alpha.$$

Parts (7.7) and (7.7) for $\mathscr{D}(G) = \mathbb{E}_6^{\text{ad}}$ follows, using (7.13), from

$$\alpha(-, -) \text{ is integral on } \Lambda(T_{\mathbb{E}_6^{\text{ad}}}) \otimes \Lambda(T_{\mathbb{E}_6^{\text{ad}}}) \iff \mathbb{Z} \ni \alpha\langle\omega_1, \omega_1\rangle = \frac{4\alpha}{3} \iff 3 \mid \alpha,$$

$$\alpha(-, -) \text{ is even on } \Lambda(T_{\mathbb{E}_6^{\text{ad}}}) \otimes \Lambda(T_{\mathbb{E}_6^{\text{ad}}}) \iff \mathbb{Z} \ni \alpha\langle\omega_1, \omega_1\rangle = \frac{4\alpha}{3} \iff 3 \mid \alpha. \quad \square$$

Lemma 7.8. *Let G be a reductive group as in the above set-up. Consider an element $\delta \in \pi_1(G)$ with image $\delta^{\text{ss}} \in \pi_1(G^{\text{ss}})$. Then we have that*

$$\text{coker}\left(\tilde{\text{ev}}_{\mathscr{D}(G)}^\delta\right) = \text{coker}\left(\text{ev}_{\mathscr{D}(G)}^\delta\right) = \begin{cases} \mathbb{Z}/2\mathbb{Z} & \text{if either } G^{\text{ss}} = \mathbb{E}_7 \\ & \text{or } \mathscr{D}(G) = \mathbb{E}_7 \text{ and } G^{\text{ss}} = \mathbb{E}_7^{\text{ad}} \text{ and } \delta^{\text{ss}} = 0 \\ \mathbb{Z}/3\mathbb{Z} & \text{if either } G^{\text{ss}} = \mathbb{E}_6 \\ & \text{or } \mathscr{D}(G) = \mathbb{E}_6 \text{ and } G^{\text{ss}} = \mathbb{E}_6^{\text{ad}} \text{ and } \delta^{\text{ss}} = 0 \\ 0 & \text{otherwise.} \end{cases}$$

Proof. If $\mathcal{D}(G)$ is of type E_8 , then the conclusion is obvious since $\mathcal{D}(G) = G^{\text{ad}}$ by (7.13).

Assume that $\mathcal{D}(G)$ is of type E_7 or E_6 . The map $\tilde{\text{ev}}_{\mathcal{D}(G)}^\delta$ (and hence also $\text{ev}_{\mathcal{D}(G)}^\delta$) is obviously zero if either $\delta^{\text{ss}} = 0$ (which is always the case if $G^{\text{ss}} = \mathbb{E}_7$ or \mathbb{E}_6 by (7.14)) or $\mathcal{D}(G)$ is equal to \mathbb{E}_7^{ad} or \mathbb{E}_6^{ad} . In the remaining cases, i.e., $(\mathcal{D}(G), G^{\text{ss}}) = (\mathbb{E}_7, \mathbb{E}_7^{\text{ad}})$ or $(\mathbb{E}_6, \mathbb{E}_6^{\text{ad}})$ and $\delta^{\text{ss}} = \omega_7 \in \pi_1(\mathbb{E}_7^{\text{ad}})$ or $\delta^{\text{ss}} = m\omega_1 \in \pi_1(\mathbb{E}_6^{\text{ad}})$ with $1 \leq m \leq 2$ (by (7.14)), Lemma 7.7 implies that

$$\text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}G} = \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)})) = \langle (-, -) \rangle$$

and hence we get that $\tilde{\text{ev}}_{\mathcal{D}(G)}^\delta = \text{ev}_{\mathcal{D}(G)}^\delta$ is non-zero (and hence surjective), because

$$\begin{aligned} \text{ev}_{\mathcal{D}(G)}^\delta((-,-)(\omega_7) &= (\omega_7, \omega_7) = \left(\frac{2\epsilon_6 + \epsilon_7 - \epsilon_8}{2}, \frac{2\epsilon_6 + \epsilon_7 - \epsilon_8}{2} \right) = \frac{6}{4} \notin \mathbb{Z}, \\ \text{ev}_{\mathcal{D}(G)}^\delta((-,-)(\omega_1) &= (m\omega_1, \omega_1) = m \left(\frac{2}{3}(\epsilon_8 - \epsilon_7 - \epsilon_6), \frac{2}{3}(\epsilon_8 - \epsilon_7 - \epsilon_6) \right) \\ &= m \frac{4}{3} \notin \mathbb{Z}. \end{aligned}$$

We conclude using (7.15). □

7.5. Types G_2, F_4

For the root systems of type G_2 and F_4 , the (co)root and (co)weight lattices (whose explicit definition we will not need to recall) satisfy

$$\begin{aligned} Q(F_4) &= P(F_4) \text{ and } Q(F_4^\vee) = P(F_4^\vee), \\ Q(G_2) &= P(G_2) \text{ and } Q(G_2^\vee) = P(G_2^\vee). \end{aligned} \tag{7.16}$$

This implies that there is a unique semisimple group \mathbb{F}_4 (respectively \mathbb{G}_2) which is almost simple of type F_4 (respectively G_2), and these two groups are both simply-connected and adjoint. In particular, we have that

$$\begin{aligned} \Lambda(T_{\mathbb{F}_4}) &= Q(F_4^\vee) = P(F_4^\vee) \text{ and } \Lambda^*(T_{\mathbb{F}_4}) = Q(F_4) = P(F_4), \\ \Lambda(T_{\mathbb{G}_2}) &= Q(G_2^\vee) = P(G_2^\vee) \text{ and } \Lambda^*(T_{\mathbb{G}_2}) = Q(G_2) = P(G_2). \end{aligned} \tag{7.17}$$

From now on, we will consider the following

Set-up: Let G be a reductive group such that $\mathcal{D}(G) = \mathbb{F}_4$ or \mathbb{G}_2 , or equivalently G is the product of a torus and either \mathbb{F}_4 or \mathbb{G}_2 .

Lemma 7.9. *Let G be a reductive group as in the above set-up. Then we have that*

- (i) $\text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}G} = \text{Bil}^{s,-\text{ev}}(\Lambda(T_{\mathcal{D}(G)}) | \Lambda(T_{G^{\text{ss}}}))^{\mathcal{W}G} = \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathcal{W}G} \cong \mathbb{Z}$. In particular, $\text{coker}(r_G) = 0$;
- (ii) $\text{coker}(\tilde{\text{ev}}_{\mathcal{D}(G)}^\delta) = \text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta) = 0$ for any $\delta \in \pi_1(G)$.

Proof. Part (i): the equality of the three lattices follows from (7.17) and Lemma 2.9, while the fact that they are of rank one follows from Corollary 2.7.

Part (ii) follows from the fact that $\mathcal{D}(G) = G^{\text{ad}}$ by (7.17). \square

References

- [1] E. ARBARELLO, M. CORNALBA and P. A. GRIFFITHS, “Geometry of Algebraic Curves”, Vol. II, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], Vol. 268, with a contribution by Joseph Daniel Harris, Springer, Heidelberg, 2011.
- [2] J. ALPER, *Adequate moduli spaces and geometrically reductive group schemes*, *Algebr. Geom.* **1** (2014), 489–531.
- [3] I. BISWAS and N. HOFFMANN, *The line bundles on moduli stacks of principal bundles on a curve*, *Doc. Math.* **15** (2010), 35–72.
- [4] I. BISWAS and A. HOGADI, *Brauer group of moduli spaces of $\text{PGL}(r)$ -bundles over a curve*, *Adv. Math.* **225** (2010), 2317–2331.
- [5] I. BISWAS and N. HOFFMANN, *Poincaré families of G -bundles on a curve*, *Math. Ann.* **352** (2012), 133–154.
- [6] I. BISWAS and Y. I. HOLLA, *Brauer group of moduli of principal bundles over a curve*, *J. Reine Angew. Math.* **677** (2013), 225–249.
- [7] A. BEAUVILLE, Y. LASZLO and C. SORGER, *The Picard group of the moduli of G -bundles on a curve*, *Compositio Math.* **112** (1998), 183–216.
- [8] Á. L. MUÑOZ CASTAÑEDA, *A compactification of the universal moduli space of principal G -bundles*, *Mediterr. J. Math.* **19** (2022), Paper No. 54, 23 pp.
- [9] G. FALTINGS, *Stable G -bundles and projective connections*, *J. Algebraic Geom.* **2** (1993), 507–568.
- [10] R. FRINGUELLI and R. PIRISI, *The Brauer group of the universal moduli space of vector bundles over smooth curves*, *Int. Math. Res. Not. IMRN* **2021**, 13609–13644.
- [11] R. FRINGUELLI, *The Picard group of the universal moduli space of vector bundles on stable curves*, *Adv. Math.* **336** (2018), 477–557.
- [12] R. FRINGUELLI and F. VIVIANI, *The Brauer group of the universal moduli stack of principal bundles on pointed smooth curves*, work in progress.
- [13] R. FRINGUELLI and F. VIVIANI, *On the Picard group scheme of the moduli stack of stable pointed curves*, preprint arXiv:2005.06920.
- [14] R. FRINGUELLI and F. VIVIANI, *The Picard group of the universal moduli stack of principal bundles on pointed smooth curves*, *J. Topol.* **15** (2022), 2065–2142.
- [15] T. L. GÓMEZ, A. LANGER, A. H. W. SCHMITT and I. SOLS, *Moduli spaces for principal bundles in arbitrary characteristic*, *Adv. Math.* **219** (2008), 1177–1245.
- [16] T. L. GÓMEZ, A. LANGER, A. H. W. SCHMITT and I. SOLS, *Moduli spaces for principal bundles in large characteristic*, In: “Teichmüller Theory and Moduli Problem”, I. Biswas, S. Kulkarni and S. Mitra (eds.), *Ramanujan Math. Soc. Lect. Notes Ser.*, Vol. 10, Ramanujan Math. Soc., Mysore, 2010, 281–371.
- [17] S. KOIZUMI, *The ring of algebraic correspondences on a generic curve of genus g* , *Nagoya Math. J.* **60** (1976), 173–180.
- [18] A. KOUVIDAKIS, *The Picard group of the universal Picard varieties over the moduli space of curves*, *J. Differential Geom.* **34** (1991), 839–850.
- [19] A. KOUVIDAKIS, *On the moduli space of vector bundles on the fibers of the universal curve*, *J. Differential Geom.* **37** (1993), 505–522.
- [20] A. LANGER, *Moduli spaces of principal bundles on singular varieties*, *Kyoto J. Math.* **53** (2013), 3–23.

- [21] Y. LASZLO, *Linearization of group stack actions and the Picard group of the moduli of SL_r/μ_s -bundles on a curve*, Bull. Soc. Math. France **125** (1997), 529–545.
- [22] Y. LASZLO and C. SORGER, *The line bundles on the moduli of parabolic G -bundles over curves and their sections*, Ann. Sci. École Norm. Sup. (4) **30** (1997), 499–525.
- [23] J. MILNE, “Reductive Groups”, available at <https://www.jmilne.org/math/CourseNotes/RG.pdf>.
- [24] M. MELO and F. VIVIANI, *The Picard group of the compactified universal Jacobian*, Doc. Math. **19** (2014), 457–507.
- [25] R. PANDHARIPANDE, *A compactification over \overline{M}_g of the universal moduli space of slope-semistable vector bundles*, J. Amer. Math. Soc. **9** (1996), 425–471.
- [26] THE STACKS PROJECT AUTHORS, *Stacks Project*, <https://stacks.math.columbia.edu>, 2018.
- [27] C. TELEMAN and C. T. WOODWARD, *The index formula for the moduli of G -bundles on a curve*, Ann. of Math. (2) **170** (2009), 495–527.

Dipartimento di Matematica
Sapienza Università di Roma
Piazzale Aldo Moro 5
I-00185 Roma, Italia
r.fringuelli@uniroma1.it

Dipartimento di Matematica e Fisica
Università Roma Tre
Largo San Leonardo Murialdo
I-00146 Roma, Italia
viviani@mat.uniroma3.it